

# Crystal structure of 2-(4-chlorophenyl)-3-(4-methoxyphenyl)-3-(methylsulfonyl)acrylonitrile

Shamantha Kumar,<sup>a</sup> Amar A. Hosamani,<sup>b</sup> A. C. Vinayaka,<sup>c</sup> M. P. Sadashiva<sup>c</sup> and B. H. Doreswamy<sup>a\*</sup>

<sup>a</sup>Department of Physics, SJB Institute of Technology, Kengeri, Bangalore 560 060, India, <sup>b</sup>Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, India, and <sup>c</sup>Department of Studies in Chemistry, Manasagangotri, University of Mysore, Mysore 570 006, India. \*Correspondence e-mail: dorephy@gmail.com

Received 12 September 2014; accepted 30 September 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


In the title compound, C<sub>17</sub>H<sub>14</sub>ClNOS, the aromatic rings are inclined to one another by 64.22 (9)°. The acrylonitrile group (C=C—C≡N) is planar to within 0.003 (2) Å, with the S atom and the methyl C atom displaced from this plane by 0.2317 (6) and −0.637 (2) Å, respectively. In the crystal, molecules are linked *via* pairs of C—H⋯π interactions, forming inversion dimers. There are no other significant intermolecular interactions present.

**Keywords:** crystal structure; acrylonitrile; C—H⋯π interactions; biological activity; pharmacological activity.

**CCDC reference:** 1026843

## 1. Related literature

For the biological and pharmacological activities of acrylonitrile derivatives, see: Boëdec *et al.* (2008); Napolitano *et al.* (2001); Saczewski *et al.* (2004); Sommen *et al.* (2003). For related literature, see: Saufi & Ismail (2002); Urska *et al.* (2003).



## 2. Experimental

### 2.1. Crystal data

C <sub>17</sub> H <sub>14</sub> ClNOS	<i>V</i> = 1542.00 (14) Å <sup>3</sup>
<i>M<sub>r</sub></i> = 315.81	<i>Z</i> = 4
Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>c</i>	Mo <i>K</i> α radiation
<i>a</i> = 8.3060 (4) Å	<i>μ</i> = 0.38 mm <sup>−1</sup>
<i>b</i> = 10.5048 (6) Å	<i>T</i> = 293 K
<i>c</i> = 17.9795 (9) Å	0.30 × 0.25 × 0.20 mm
<i>β</i> = 100.598 (5)°	

### 2.2. Data collection

Bruker APEXII CCD area-detector diffractometer	3537 independent reflections
6889 measured reflections	2807 reflections with <i>I</i> > 2σ( <i>I</i> )
	<i>R</i> <sub>int</sub> = 0.028

### 2.3. Refinement

<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )] = 0.046	192 parameters
<i>wR</i> ( <i>F</i> <sup>2</sup> ) = 0.114	H-atom parameters constrained
<i>S</i> = 1.04	Δ <i>ρ</i> <sub>max</sub> = 0.26 e Å <sup>−3</sup>
3537 reflections	Δ <i>ρ</i> <sub>min</sub> = −0.26 e Å <sup>−3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 ring.

<i>D</i> —H⋯ <i>A</i>	<i>D</i> —H	H⋯ <i>A</i>	<i>D</i> ⋯ <i>A</i>	<i>D</i> —H⋯ <i>A</i>
C15—H15⋯Cg1 <sup>1</sup>	0.93	2.96	3.739 (2)	142

Symmetry code: (i)  $-x + 1, -y + 2, -z$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

## Acknowledgements

The authors are grateful to the SJB Institute of Technology, Kengeri, Bangalore, for their support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU2787).

## References

- Boëdec, A., Sicard, H., Dessolin, J., Herbette, G., Ingoure, S., Raymond, C., Belmont, C. & Kraus, J. L. (2008). *J. Med. Chem.* **51**, 1747–1754.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Napolitano, A., Bruno, I., Rovero, P., Lucas, R., Peris, M. P. & Riccio, R. (2001). *Tetrahedron*, **57**, 6249–6255.
- Saczewski, F., Reszka, P., Gdaniec, M., Grünert, R. & Bednarski, P. J. (2004). *J. Med. Chem.* **47**, 3438–3449.
- Saufi, S. M. & Ismail, A. F. (2002). *Songklanakarin J. Sci. Technol.* **24**, 843–854.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sommen, G., Comel, A. & Kirsch, G. (2003). *Tetrahedron*, **59**, 1557–1564.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Urska, B., Anton, M., Jurij, S. & Branko, S. (2003). *Arkivoc*, **5**, 77–86.

## supporting information

*Acta Cryst.* (2014). E70, o1155 [doi:10.1107/S1600536814021576]

## Crystal structure of 2-(4-chlorophenyl)-3-(4-methoxyphenyl)-3-(methylsulfanyl)acrylonitrile


Shamantha Kumar, Amar A. Hosamani, A. C. Vinayaka, M. P. Sadashiva and B. H. Doreswamy

### S1. Experimental

To a stirred suspension of NaH (0.45 g, 11.0 mmol, 60% suspension in oil) in dry THF (25 ml), a solution of (4-chlorophenyl)acetonitrile (0.75 g, 5.0 mmol) and 4-methoxy-dithiobenzoic acid methyl ester (0.99 g, 5.0 mmol) in dry THF (50 ml) was added drop wise at 273 K under a nitrogen atmosphere. The reaction mixture was stirred at room temperature for 4 h. It was again cooled to 273 K and methyl iodide (1.42 g, 10 mmol) was added drop wise. The reaction mixture was further stirred at room temperature for 4 h and poured into ice cold water (25 ml). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 10$  ml). The combined organic extracts were washed with water ( $1 \times 20$  ml), brine ( $1 \times 20$  ml), and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solvent was evaporated under reduced pressure to give a crude product which was purified by silica gel column using EtOAc:hexane as eluent. Colourless block-like crystals were grown by dissolving the product in absolute ethanol followed by slow evaporation at room temperature.

### S2. Refinement

The C-bound H atoms were positioned geometrically and allowed to ride on their parent atoms: C–H = 0.93 - 0.96 Å, with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and =  $1.2U_{\text{eq}}(\text{C})$  for other H atoms.



**Figure 1**

A view of the molecular structure of the title molecule, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

A view along the *b* axis of the crystal packing of the title compound.

### 2-(4-Chlorophenyl)-3-(4-methoxyphenyl)-3-(methylsulfanyl)acrylonitrile

#### Crystal data

$C_{17}H_{14}ClNO$

$M_r = 315.81$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P 2_1/c$

$a = 8.3060$  (4) Å

$b = 10.5048$  (6) Å

$c = 17.9795$  (9) Å

$\beta = 100.598$  (5)°

$V = 1542.00$  (14) Å<sup>3</sup>

$Z = 4$

$F(000) = 656$

$D_x = 1.360$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 3537 reflections

$\theta = 2.5$ – $27.5$ °

$\mu = 0.38$  mm<sup>-1</sup>

$T = 293$  K

Block, colourless

$0.30 \times 0.25 \times 0.20$  mm

#### Data collection

Bruker APEXII CCD area-detector  
diffractometer

$\omega$  and  $\varphi$  scans

6889 measured reflections

3537 independent reflections

2807 reflections with  $I > 2\sigma(I)$

$R_{int} = 0.028$

$\theta_{max} = 27.5$ °,  $\theta_{min} = 2.5$ °

$h = -10$ → $6$

$k = -13$ → $5$

$l = -23$ → $23$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.114$

$S = 1.04$

3537 reflections

192 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.045P)^2 + 0.3209P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$

$\Delta\rho_{max} = 0.26$  e Å<sup>-3</sup>

$\Delta\rho_{min} = -0.26$  e Å<sup>-3</sup>

*Special details*

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement on  $F^2$  for ALL reflections except those flagged by the user for potential systematic errors. Weighted  $R$ -factors  $wR$  and all goodnesses of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The observed criterion of  $F^2 > \sigma(F^2)$  is used only for calculating  $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl7	-0.19319 (7)	1.21573 (7)	-0.19619 (4)	0.0708 (2)
S12	0.39592 (6)	0.84311 (6)	0.00222 (3)	0.0463 (2)
O20	1.09261 (17)	0.59095 (16)	-0.06455 (9)	0.0603 (6)
N10	0.5710 (2)	0.9039 (2)	-0.25595 (10)	0.0575 (7)
C1	0.2687 (2)	1.09961 (19)	-0.18246 (10)	0.0396 (6)
C2	0.1284 (2)	1.1742 (2)	-0.19521 (11)	0.0463 (6)
C3	-0.0153 (2)	1.1236 (2)	-0.18024 (11)	0.0441 (6)
C4	-0.0223 (2)	1.0011 (2)	-0.15428 (12)	0.0496 (7)
C5	0.1178 (2)	0.9272 (2)	-0.14196 (11)	0.0452 (6)
C6	0.2658 (2)	0.97624 (18)	-0.15536 (9)	0.0347 (5)
C8	0.4179 (2)	0.89756 (18)	-0.14288 (10)	0.0346 (5)
C9	0.5065 (2)	0.8996 (2)	-0.20479 (10)	0.0401 (6)
C11	0.4820 (2)	0.83108 (18)	-0.07963 (10)	0.0353 (5)
C13	0.4464 (3)	0.6919 (2)	0.04764 (12)	0.0570 (8)
C14	0.6374 (2)	0.75884 (19)	-0.07423 (10)	0.0350 (5)
C15	0.7767 (2)	0.7996 (2)	-0.02329 (10)	0.0402 (6)
C16	0.9244 (2)	0.7402 (2)	-0.02164 (11)	0.0443 (6)
C17	0.9379 (2)	0.6388 (2)	-0.06941 (11)	0.0406 (6)
C18	0.8005 (2)	0.5942 (2)	-0.11771 (11)	0.0441 (6)
C19	0.6513 (2)	0.6553 (2)	-0.11980 (11)	0.0429 (6)
C21	1.1194 (3)	0.4982 (3)	-0.11813 (15)	0.0742 (10)
H1	0.36640	1.13300	-0.19230	0.0470*
H2	0.13140	1.25680	-0.21350	0.0550*
H4	-0.12060	0.96800	-0.14510	0.0600*
H5	0.11330	0.84410	-0.12460	0.0540*
H13A	0.40050	0.62460	0.01420	0.0860*
H13B	0.40220	0.68780	0.09330	0.0860*
H13C	0.56320	0.68250	0.05960	0.0860*
H15	0.76880	0.86710	0.00940	0.0480*
H16	1.01650	0.76820	0.01190	0.0530*
H18	0.80790	0.52400	-0.14840	0.0530*
H19	0.55900	0.62580	-0.15260	0.0520*
H21A	1.06080	0.42180	-0.11070	0.1110*
H21B	1.23430	0.47980	-0.11170	0.1110*
H21C	1.08110	0.53000	-0.16830	0.1110*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C17	0.0571 (3)	0.0760 (5)	0.0789 (4)	0.0308 (3)	0.0114 (3)	0.0093 (4)
S12	0.0507 (3)	0.0523 (3)	0.0385 (3)	0.0136 (2)	0.0150 (2)	0.0035 (2)
O20	0.0451 (8)	0.0666 (11)	0.0699 (10)	0.0192 (7)	0.0125 (7)	-0.0035 (9)
N10	0.0579 (11)	0.0707 (14)	0.0479 (10)	0.0082 (10)	0.0199 (8)	0.0068 (10)
C1	0.0436 (10)	0.0401 (11)	0.0362 (9)	-0.0011 (8)	0.0104 (8)	0.0014 (9)
C2	0.0594 (12)	0.0383 (11)	0.0421 (10)	0.0073 (9)	0.0120 (9)	0.0045 (9)
C3	0.0447 (10)	0.0491 (12)	0.0376 (10)	0.0149 (9)	0.0049 (8)	-0.0006 (9)
C4	0.0348 (9)	0.0585 (14)	0.0548 (12)	0.0010 (9)	0.0065 (8)	0.0081 (11)
C5	0.0393 (10)	0.0412 (12)	0.0532 (11)	-0.0006 (8)	0.0033 (8)	0.0103 (10)
C6	0.0369 (9)	0.0369 (10)	0.0293 (8)	0.0019 (8)	0.0033 (7)	-0.0004 (8)
C8	0.0344 (9)	0.0353 (10)	0.0336 (9)	-0.0007 (7)	0.0051 (7)	-0.0027 (8)
C9	0.0374 (9)	0.0437 (12)	0.0377 (10)	0.0035 (8)	0.0033 (8)	0.0013 (9)
C11	0.0360 (9)	0.0346 (10)	0.0351 (9)	-0.0002 (7)	0.0064 (7)	-0.0039 (8)
C13	0.0608 (13)	0.0651 (16)	0.0479 (12)	0.0059 (11)	0.0171 (10)	0.0161 (11)
C14	0.0368 (9)	0.0365 (10)	0.0321 (8)	0.0022 (7)	0.0078 (7)	0.0015 (8)
C15	0.0423 (10)	0.0408 (11)	0.0370 (9)	-0.0008 (8)	0.0063 (8)	-0.0078 (9)
C16	0.0358 (9)	0.0521 (13)	0.0437 (10)	-0.0014 (9)	0.0036 (8)	-0.0044 (10)
C17	0.0405 (10)	0.0420 (12)	0.0409 (10)	0.0071 (8)	0.0118 (8)	0.0077 (9)
C18	0.0552 (11)	0.0366 (11)	0.0405 (10)	0.0078 (9)	0.0091 (9)	-0.0052 (9)
C19	0.0434 (10)	0.0434 (12)	0.0396 (10)	0.0022 (9)	0.0013 (8)	-0.0049 (9)
C21	0.0776 (17)	0.0797 (19)	0.0691 (15)	0.0403 (15)	0.0239 (13)	0.0036 (15)

*Geometric parameters (Å, °)*

C17—C3	1.745 (2)	C15—C16	1.372 (3)
S12—C11	1.7550 (18)	C16—C17	1.386 (3)
S12—C13	1.800 (2)	C17—C18	1.383 (3)
O20—C17	1.368 (2)	C18—C19	1.390 (3)
O20—C21	1.416 (3)	C1—H1	0.9300
N10—C9	1.147 (2)	C2—H2	0.9300
C1—C2	1.388 (3)	C4—H4	0.9300
C1—C6	1.386 (3)	C5—H5	0.9300
C2—C3	1.378 (3)	C13—H13A	0.9600
C3—C4	1.374 (3)	C13—H13B	0.9600
C4—C5	1.382 (3)	C13—H13C	0.9600
C5—C6	1.394 (2)	C15—H15	0.9300
C6—C8	1.492 (2)	C16—H16	0.9300
C8—C9	1.443 (2)	C18—H18	0.9300
C8—C11	1.357 (3)	C19—H19	0.9300
C11—C14	1.485 (2)	C21—H21A	0.9600
C14—C15	1.404 (2)	C21—H21B	0.9600
C14—C19	1.379 (3)	C21—H21C	0.9600
C11—S12—C13	102.71 (10)	C2—C1—H1	119.00
C17—O20—C21	118.21 (17)	C6—C1—H1	119.00

C2—C1—C6	121.16 (16)	C1—C2—H2	121.00
C1—C2—C3	118.77 (19)	C3—C2—H2	121.00
C17—C3—C2	119.45 (16)	C3—C4—H4	120.00
C17—C3—C4	119.20 (14)	C5—C4—H4	120.00
C2—C3—C4	121.34 (17)	C4—C5—H5	120.00
C3—C4—C5	119.57 (17)	C6—C5—H5	120.00
C4—C5—C6	120.52 (19)	S12—C13—H13A	109.00
C1—C6—C5	118.62 (17)	S12—C13—H13B	109.00
C1—C6—C8	120.12 (15)	S12—C13—H13C	110.00
C5—C6—C8	121.24 (17)	H13A—C13—H13B	109.00
C6—C8—C9	114.44 (15)	H13A—C13—H13C	109.00
C6—C8—C11	127.02 (16)	H13B—C13—H13C	109.00
C9—C8—C11	118.48 (16)	C14—C15—H15	120.00
N10—C9—C8	176.9 (2)	C16—C15—H15	120.00
S12—C11—C8	120.54 (14)	C15—C16—H16	120.00
S12—C11—C14	117.73 (13)	C17—C16—H16	120.00
C8—C11—C14	121.34 (16)	C17—C18—H18	120.00
C11—C14—C15	119.27 (17)	C19—C18—H18	120.00
C11—C14—C19	122.06 (16)	C14—C19—H19	119.00
C15—C14—C19	118.62 (16)	C18—C19—H19	119.00
C14—C15—C16	120.12 (18)	O20—C21—H21A	109.00
C15—C16—C17	120.68 (17)	O20—C21—H21B	110.00
O20—C17—C16	115.16 (16)	O20—C21—H21C	109.00
O20—C17—C18	124.96 (18)	H21A—C21—H21B	109.00
C16—C17—C18	119.88 (17)	H21A—C21—H21C	109.00
C17—C18—C19	119.30 (19)	H21B—C21—H21C	109.00
C14—C19—C18	121.31 (17)		
C13—S12—C11—C8	152.50 (16)	C9—C8—C11—S12	170.97 (14)
C13—S12—C11—C14	-34.67 (17)	C9—C8—C11—C14	-1.6 (3)
C21—O20—C17—C18	7.3 (3)	C6—C8—C11—S12	-5.9 (3)
C21—O20—C17—C16	-172.2 (2)	C6—C8—C11—C14	-178.51 (17)
C2—C1—C6—C8	179.37 (17)	S12—C11—C14—C15	-60.3 (2)
C2—C1—C6—C5	0.9 (3)	C8—C11—C14—C19	-65.1 (3)
C6—C1—C2—C3	0.2 (3)	S12—C11—C14—C19	122.17 (18)
C1—C2—C3—C17	-179.80 (15)	C8—C11—C14—C15	112.5 (2)
C1—C2—C3—C4	-1.0 (3)	C11—C14—C19—C18	175.46 (18)
C2—C3—C4—C5	0.8 (3)	C15—C14—C19—C18	-2.1 (3)
C17—C3—C4—C5	179.59 (16)	C11—C14—C15—C16	-174.94 (18)
C3—C4—C5—C6	0.3 (3)	C19—C14—C15—C16	2.7 (3)
C4—C5—C6—C8	-179.57 (18)	C14—C15—C16—C17	-0.7 (3)
C4—C5—C6—C1	-1.1 (3)	C15—C16—C17—C18	-2.0 (3)
C1—C6—C8—C9	-47.0 (2)	C15—C16—C17—O20	177.50 (18)
C5—C6—C8—C11	-51.5 (3)	O20—C17—C18—C19	-176.86 (19)
C1—C6—C8—C11	130.0 (2)	C16—C17—C18—C19	2.6 (3)
C5—C6—C8—C9	131.45 (18)	C17—C18—C19—C14	-0.5 (3)

*Hydrogen-bond geometry (Å, °)*

Cg1 is the centroid of the C1–C6 ring.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C15—H15···Cg1 <sup>i</sup>	0.93	2.96	3.739 (2)	142

Symmetry code: (i)  $-x+1, -y+2, -z$ .