

Crystal structure of (Z)-3-(4-methoxybenzylidene)-2,3-dihydrobenzo[*b*][1,4]-thiazepin-4(5*H*)-one

V. Vinayagam,^a J. Mohan Raj,^b S. Murugavel,^{c*}
R. Selvakumar^a and M. Bakthadoss^{a,d*}

^aDepartment of Organic Chemistry, University of Madras, Maraimalai Campus, Chennai 600 025, India, ^bDepartment of Physics, C. Abdul Hakeem College of Engineering & Technology, Melvisharam, Vellore 632 509, India, ^cDepartment of Physics, Thanthai Periyar Government Institute of Technology, Vellore 632 002, India, and ^dDepartment of Chemistry, Pondicherry University, Puducherry 605 014, India. *Correspondence e-mail: smurugavel27@gmail.com, bhakthadoss@yahoo.com

Received 26 November 2014; accepted 30 November 2014

Edited by E. R. T. Tiekink, University of Malaya, Malaysia


In the title compound, C₁₇H₁₅NO₂S, the two C atoms linking the S and carbonyl C atoms of the seven-membered thiazepine ring are disordered over two sites, with occupancies of 0.511 (4) and 0.489 (4); both disorder components adopt distorted twist-boat conformations. In the crystal, N—H···O and C—H···O hydrogen bonds link inverted-related molecules into dimers, incorporating R₁²(6) and R₂²(8) ring motifs; the acceptor carbonyl O atom is bifurcated. These dimers are further linked by C—H···O hydrogen bonds, forming supra-molecular tapes running along the *a* axis.

Keywords: crystal structure; benzo[*b*][1,4]thiazepin-4(5*H*)-one; pharmaceutical properties; thiazepin derivatives; hydrogen bonding.

CCDC reference: 1036763

1. Related literature

For the pharmaceutical properties of thiazepin derivatives, see: Lončar-Tomasovic *et al.* (2000); Rajsner *et al.* (1971); Metys & Metysová (1965). For related structures, see: Lakshmanan *et al.* (2012); Selvakumar *et al.* (2012).



2. Experimental

2.1. Crystal data

C₁₇H₁₅NO₂S
M_r = 297.36
 Monoclinic, C₂/c
a = 21.434 (5) Å
b = 5.715 (4) Å
c = 23.870 (5) Å
 β = 101.091 (4)°

V = 2869 (2) Å³
Z = 8
 Mo *K*α radiation
 μ = 0.23 mm⁻¹
T = 293 K
 0.30 × 0.30 × 0.25 mm

2.2. Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
*T*_{min} = 0.934, *T*_{max} = 0.944

17286 measured reflections
 4099 independent reflections
 2744 reflections with *I* > 2σ(*I*)
*R*_{int} = 0.028

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.139$
S = 1.03
 4099 reflections
 210 parameters

4 restraints
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1···O1 ⁱ	0.86	2.07	2.9291 (18)	177
C6—H6···O1 ⁱ	0.93	2.45	3.263 (3)	146
C1B—H1C···O1 ⁱⁱ	0.97	2.52	3.377 (4)	147

Symmetry codes: (i) $-x, -y, -z$; (ii) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

Acknowledgements

The authors thank Dr Babu Vargheese, SAIF, IIT, Madras, India, for his help with the data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5350).

References

- Bruker (2004). *APEX2, SAINT and XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Lakshmanan, D., Murugavel, S., Selvakumar, R. & Bakthadoss, M. (2012). *Acta Cryst.* **E68**, o2130.
- Lončar-Tomašević, L., Šarac-Arneri, R., Hergold-Brundić, A., Nagl, A., Mintas, M. & Sandström, J. (2000). *HCA*, **83**, 479–494.
- Metys, J. & Metysová, J. (1965). *Acta Biol. Med. Ger.* **15**, 871–873.
- Rajsner, M., Protiva, M. & Metysova, J. (1971). Czech. Patent Appl. CS 143737.
- Selvakumar, R., Bakthadoss, M., Lakshmanan, D. & Murugavel, S. (2012). *Acta Cryst.* **E68**, o2126.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2015). E71, o21–o22 [https://doi.org/10.1107/S2056989014026267]

Crystal structure of (*Z*)-3-(4-methoxybenzylidene)-2,3-dihydrobenzo[*b*][1,4]thiazepin-4(5*H*)-one

V. Vinayagam, J. Mohan Raj, S. Murugavel, R. Selvakumar and M. Bakthadoss

S1. Structural commentary

The title compound is used as an intermediate for the synthesis of dosulepin, which is an antidepressant of the tricyclic family. Dosulepin prevents reabsorption of serotonin and noradrenaline in the brain, helps to prolong the mood lightening effect of any released noradrenaline and serotonin, thus relieving depression. Dibenzoc[e]thiazepin derivatives exhibit chiroptical properties (Tomascovic *et al.*, 2000). Dibenzo[*b,e*]thiazepin-5,5-dioxide derivatives possess antihistaminic and anti-allergenic activities (Rajsner *et al.*, 1971). Benzene thiazepin derivatives are identified as a type of effective anti-histaminic compounds (Metys *et al.*, 1965). In view of this biological importance, the crystal structure of the title compound has been carried out and the results are presented here.

Fig. 1 shows a displacement ellipsoid plot of (I), with the atom numbering scheme. The geometric parameters of the title molecule agree well with those reported for similar structures (Selvakumar *et al.*, 2012; Lakshmanan *et al.*, 2012). The sum of angles at N1 atom of the thiazepin ring (359.9°) is in accordance with sp^2 hybridization. Both the major and minor conformers of the disordered thiazepine ring adopt distorted twist-boat conformations.


In the crystal, intermolecular bifurcated acceptor N1—H1 \cdots O1ⁱ and C6—H6 \cdots O1ⁱ (Table 1) hydrogen bonds link inverted-related molecules into dimers, incorporating $R_1^2(6)$ and $R_2^2(8)$ ring motifs. These dimers are further linked by C1B—H1C \cdots O1ⁱⁱ (Table 1) hydrogen bonds forming supramolecular tapes running along the *a* axis (Fig. 2).

S2. Synthesis and crystallization

A mixture of (*Z*)-methyl 2-(bromomethyl)-3-(4-methoxyphenyl)acrylate (2 mmol) and *o*-aminothiophenol (2 mmol) in the presence of potassium *tert*-butoxide (4.8 mmol) in dry THF (10 ml) was stirred at room temperature for 1 h. After the completion of the reaction as indicated by TLC, the reaction mixture was concentrated and the resulting crude mass was diluted with water (20 ml) and extracted with ethyl acetate (3 x 20 ml). The organic layer was washed with brine (2 x 20 ml) and dried over anhydrous sodium sulfate. It was then concentrated to successfully provide the crude final product ((*Z*)-3-(4-methoxybenzylidene)-2,3-dihydrobenzo[*b*][1,4]thiazepin-4(5*H*)-one). This was purified by column chromatography on silica gel with ethylacetate/hexane 1:19 as eluent to afford the title compound in good yield (47 %). Single crystals suitable for X-ray diffraction were obtained by slow evaporation of its ethylacetate solution at room temperature.

S3. Refinement

Atoms C1 and C9 of the thiazepine ring are disordered over two positions (C1A/C1B and C9A/C9B) with refined occupancies of 0.511 (4) and 0.489 (4). The corresponding bond distances involving the disordered atoms were restrained to be equal. H atoms were positioned geometrically, (C—H = 0.93–0.97 Å and N—H = 0.86 Å) constrained to ride on their parent atom, with $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

**Figure 1**

Molecular structure of the title compound showing displacement ellipsoids at the 30% probability level. H atoms are presented as a small spheres of arbitrary radii.




Figure 2

Supramolecular tape formation in the crystal packing of the title compound whereby bifurcated hydrogen bonds link inverted molecules into dimers sustained by N—H \cdots O and C—H \cdots O (red dashed lines) contacts are linked *via* C—H \cdots O contacts (blue dashed lines) along *a* axis. [Symmetry code: (i) $-x, -y, -z$; (ii) $x, 1+y, z$].

(Z)-3-(4-Methoxybenzylidene)-2,3-dihydrobenzo[*b*][1,4]thiazepin-4(5*H*)-one

Crystal data

$C_{17}H_{15}NO_2S$

$M_r = 297.36$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 21.434 (5) \text{ \AA}$

$b = 5.715 (4) \text{ \AA}$

$c = 23.870 (5) \text{ \AA}$

$\beta = 101.091 (4)^\circ$

$V = 2869 (2) \text{ \AA}^3$

$Z = 8$

$F(000) = 1248$

$D_x = 1.377 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4132 reflections

$\theta = 1.7\text{--}29.9^\circ$

$\mu = 0.23 \text{ mm}^{-1}$

$T = 293$ K $0.30 \times 0.30 \times 0.25$ mm
 Block, colourless

Data collection

Bruker APEXII CCD diffractometer Radiation source: fine-focus sealed tube Graphite monochromator Detector resolution: 10.0 pixels mm^{-1} ω scans Absorption correction: multi-scan (SADABS; Sheldrick, 1996) $T_{\min} = 0.934$, $T_{\max} = 0.944$	17286 measured reflections 4099 independent reflections 2744 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.028$ $\theta_{\max} = 29.9^\circ$, $\theta_{\min} = 1.7^\circ$ $h = -30 \rightarrow 24$ $k = -7 \rightarrow 8$ $l = -33 \rightarrow 32$
---	--

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.045$ $wR(F^2) = 0.139$ $S = 1.03$ 4099 reflections 210 parameters 4 restraints Primary atom site location: structure-invariant direct methods	Secondary atom site location: difference Fourier map Hydrogen site location: inferred from neighbouring sites H-atom parameters constrained $w = 1/[\sigma^2(F_o^2) + (0.0647P)^2 + 1.1615P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$
--	---

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C2	-0.05513 (8)	0.5769 (3)	0.10567 (7)	0.0516 (4)	
C3	-0.09963 (11)	0.7568 (3)	0.10107 (9)	0.0689 (5)	
H3	-0.1001	0.8512	0.1327	0.083*	
C4	-0.14266 (12)	0.8004 (4)	0.05193 (10)	0.0790 (6)	
H4	-0.1718	0.9219	0.0503	0.095*	
C5	-0.14211 (10)	0.6629 (4)	0.00527 (10)	0.0722 (5)	
H5	-0.1708	0.6906	-0.0286	0.087*	
C6	-0.09941 (10)	0.4851 (4)	0.00854 (9)	0.0691 (5)	
H6	-0.0996	0.3927	-0.0236	0.083*	
C7	-0.05515 (8)	0.4357 (3)	0.05837 (8)	0.0525 (4)	
C8	0.03347 (9)	0.1194 (3)	0.08459 (7)	0.0549 (4)	
C10	0.10676 (9)	0.0681 (3)	0.17287 (7)	0.0528 (4)	
H10A	0.1284	-0.0130	0.1487	0.063*	0.489 (4)
H10B	0.1093	-0.0785	0.1566	0.063*	0.511 (4)

C11	0.14272 (8)	0.0921 (3)	0.23129 (7)	0.0474 (4)	
C12	0.13743 (8)	0.2801 (3)	0.26731 (7)	0.0534 (4)	
H12	0.1074	0.3957	0.2549	0.064*	
C13	0.17552 (8)	0.2999 (3)	0.32082 (7)	0.0525 (4)	
H13	0.1710	0.4276	0.3439	0.063*	
C14	0.22010 (8)	0.1302 (3)	0.33987 (7)	0.0492 (4)	
C15	0.22513 (8)	-0.0627 (3)	0.30585 (8)	0.0551 (4)	
H15	0.2542	-0.1806	0.3190	0.066*	
C16	0.18736 (8)	-0.0793 (3)	0.25292 (8)	0.0533 (4)	
H16	0.1915	-0.2093	0.2305	0.064*	
C17	0.26254 (13)	0.3437 (5)	0.42387 (10)	0.0895 (7)	
H17A	0.2716	0.4752	0.4017	0.134*	
H17B	0.2950	0.3305	0.4576	0.134*	
H17C	0.2220	0.3656	0.4346	0.134*	
N1	-0.01563 (7)	0.2418 (3)	0.05307 (6)	0.0610 (4)	
H1	-0.0258	0.1826	0.0194	0.073*	
O1	0.05465 (6)	-0.0461 (2)	0.06174 (5)	0.0605 (3)	
O2	0.26115 (6)	0.1372 (2)	0.39107 (6)	0.0691 (4)	
S1	-0.00374 (3)	0.55138 (8)	0.17212 (2)	0.06381 (18)	
C1A	0.00694 (16)	0.2393 (5)	0.18212 (13)	0.0438 (8)	0.489 (4)
H1A	0.0222	0.2080	0.2224	0.053*	0.489 (4)
H1B	-0.0340	0.1629	0.1710	0.053*	0.489 (4)
C9A	0.05207 (17)	0.1363 (7)	0.14902 (14)	0.0425 (8)	0.489 (4)
C1B	0.06528 (16)	0.4832 (6)	0.14859 (14)	0.0529 (9)	0.511 (4)
H1C	0.0661	0.5669	0.1134	0.063*	0.511 (4)
H1D	0.1016	0.5338	0.1768	0.063*	0.511 (4)
C9B	0.07040 (18)	0.2265 (6)	0.13852 (15)	0.0467 (8)	0.511 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C2	0.0612 (10)	0.0418 (8)	0.0559 (9)	-0.0078 (7)	0.0218 (8)	-0.0015 (7)
C3	0.0945 (15)	0.0501 (10)	0.0679 (12)	0.0085 (10)	0.0301 (11)	0.0005 (9)
C4	0.0937 (16)	0.0612 (12)	0.0861 (15)	0.0218 (11)	0.0275 (13)	0.0105 (11)
C5	0.0695 (12)	0.0685 (13)	0.0761 (13)	0.0072 (10)	0.0079 (10)	0.0050 (11)
C6	0.0674 (12)	0.0676 (12)	0.0677 (12)	0.0064 (10)	0.0018 (9)	-0.0141 (10)
C7	0.0505 (9)	0.0505 (9)	0.0570 (9)	-0.0039 (7)	0.0117 (7)	-0.0085 (7)
C8	0.0606 (10)	0.0570 (10)	0.0470 (9)	0.0012 (8)	0.0105 (8)	-0.0085 (7)
C10	0.0680 (11)	0.0419 (8)	0.0494 (9)	-0.0073 (8)	0.0134 (8)	-0.0044 (7)
C11	0.0530 (9)	0.0431 (8)	0.0481 (8)	-0.0056 (7)	0.0147 (7)	-0.0008 (6)
C12	0.0564 (10)	0.0456 (9)	0.0562 (9)	0.0089 (7)	0.0060 (8)	-0.0005 (7)
C13	0.0556 (10)	0.0487 (9)	0.0530 (9)	0.0055 (7)	0.0097 (7)	-0.0071 (7)
C14	0.0464 (8)	0.0509 (9)	0.0514 (9)	0.0014 (7)	0.0122 (7)	0.0047 (7)
C15	0.0538 (10)	0.0464 (9)	0.0658 (11)	0.0106 (7)	0.0135 (8)	0.0047 (8)
C16	0.0601 (10)	0.0418 (8)	0.0614 (10)	0.0026 (7)	0.0200 (8)	-0.0054 (7)
C17	0.1079 (18)	0.0866 (16)	0.0624 (12)	0.0135 (14)	-0.0125 (12)	-0.0123 (12)
N1	0.0660 (9)	0.0634 (9)	0.0499 (8)	0.0091 (7)	0.0020 (7)	-0.0183 (7)
O1	0.0781 (8)	0.0546 (7)	0.0481 (7)	0.0079 (6)	0.0104 (6)	-0.0085 (5)

O2	0.0680 (8)	0.0720 (9)	0.0597 (8)	0.0116 (7)	-0.0064 (6)	0.0002 (7)
S1	0.0906 (4)	0.0514 (3)	0.0512 (3)	-0.0002 (2)	0.0179 (2)	-0.00693 (19)
C1A	0.0543 (18)	0.0415 (16)	0.0383 (15)	-0.0025 (13)	0.0156 (13)	-0.0022 (12)
C9A	0.048 (2)	0.0385 (18)	0.0434 (17)	-0.0072 (15)	0.0143 (15)	-0.0046 (14)
C1B	0.063 (2)	0.0443 (17)	0.0482 (17)	-0.0116 (14)	0.0023 (15)	0.0032 (13)
C9B	0.0474 (19)	0.0451 (19)	0.0477 (18)	-0.0094 (15)	0.0093 (15)	-0.0068 (15)

Geometric parameters (Å, °)

C2—C7	1.388 (2)	C12—C13	1.382 (2)
C2—C3	1.392 (3)	C12—H12	0.9300
C2—S1	1.7545 (19)	C13—C14	1.376 (2)
C3—C4	1.368 (3)	C13—H13	0.9300
C3—H3	0.9300	C14—O2	1.362 (2)
C4—C5	1.365 (3)	C14—C15	1.386 (2)
C4—H4	0.9300	C15—C16	1.367 (2)
C5—C6	1.360 (3)	C15—H15	0.9300
C5—H5	0.9300	C16—H16	0.9300
C6—C7	1.400 (3)	C17—O2	1.413 (3)
C6—H6	0.9300	C17—H17A	0.9600
C7—N1	1.415 (2)	C17—H17B	0.9600
C8—O1	1.222 (2)	C17—H17C	0.9600
C8—N1	1.363 (2)	N1—H1	0.8600
C8—C9B	1.505 (4)	S1—C1B	1.725 (3)
C8—C9A	1.516 (4)	S1—C1A	1.808 (3)
C10—C9A	1.262 (4)	C1A—C9A	1.484 (4)
C10—C9B	1.361 (4)	C1A—H1A	0.9700
C10—C11	1.464 (2)	C1A—H1B	0.9700
C10—H10A	0.9300	C1B—C9B	1.494 (4)
C10—H10B	0.9300	C1B—H1C	0.9700
C11—C12	1.395 (2)	C1B—H1D	0.9700
C11—C16	1.397 (2)		
C7—C2—C3	118.18 (18)	C12—C13—H13	120.1
C7—C2—S1	126.08 (14)	O2—C14—C13	124.28 (16)
C3—C2—S1	115.74 (14)	O2—C14—C15	116.08 (15)
C4—C3—C2	122.73 (19)	C13—C14—C15	119.64 (16)
C4—C3—H3	118.6	C16—C15—C14	119.87 (15)
C2—C3—H3	118.6	C16—C15—H15	120.1
C5—C4—C3	119.0 (2)	C14—C15—H15	120.1
C5—C4—H4	120.5	C15—C16—C11	122.34 (16)
C3—C4—H4	120.5	C15—C16—H16	118.8
C6—C5—C4	119.7 (2)	C11—C16—H16	118.8
C6—C5—H5	120.2	O2—C17—H17A	109.5
C4—C5—H5	120.2	O2—C17—H17B	109.5
C5—C6—C7	122.6 (2)	H17A—C17—H17B	109.5
C5—C6—H6	118.7	O2—C17—H17C	109.5
C7—C6—H6	118.7	H17A—C17—H17C	109.5

C2—C7—C6	117.89 (17)	H17B—C17—H17C	109.5
C2—C7—N1	128.43 (17)	C8—N1—C7	139.85 (15)
C6—C7—N1	113.67 (16)	C8—N1—H1	110.1
O1—C8—N1	117.68 (15)	C7—N1—H1	110.1
O1—C8—C9B	121.1 (2)	C14—O2—C17	117.57 (15)
N1—C8—C9B	119.10 (19)	C1B—S1—C2	98.80 (12)
O1—C8—C9A	117.03 (19)	C1B—S1—C1A	74.06 (15)
N1—C8—C9A	123.48 (18)	C2—S1—C1A	104.06 (12)
C9B—C8—C9A	27.63 (14)	C9A—C1A—S1	113.6 (2)
C9A—C10—C9B	31.63 (17)	C9A—C1A—H1A	108.8
C9A—C10—C11	132.4 (2)	S1—C1A—H1A	108.8
C9B—C10—C11	130.31 (19)	C9A—C1A—H1B	108.8
C9A—C10—H10A	113.8	S1—C1A—H1B	108.8
C9B—C10—H10A	104.7	H1A—C1A—H1B	107.7
C11—C10—H10A	113.8	C10—C9A—C1A	121.8 (3)
C9A—C10—H10B	102.4	C10—C9A—C8	118.6 (2)
C9B—C10—H10B	114.8	C1A—C9A—C8	119.6 (3)
C11—C10—H10B	114.9	C9B—C1B—S1	111.6 (2)
H10A—C10—H10B	38.2	C9B—C1B—H1C	109.3
C12—C11—C16	116.35 (16)	S1—C1B—H1C	109.3
C12—C11—C10	124.67 (15)	C9B—C1B—H1D	109.3
C16—C11—C10	118.94 (15)	S1—C1B—H1D	109.3
C13—C12—C11	121.95 (16)	H1C—C1B—H1D	108.0
C13—C12—H12	119.0	C10—C9B—C1B	127.4 (3)
C11—C12—H12	119.0	C10—C9B—C8	113.0 (2)
C14—C13—C12	119.80 (16)	C1B—C9B—C8	119.6 (3)
C14—C13—H13	120.1		
C7—C2—C3—C4	-0.6 (3)	C7—C2—S1—C1B	39.04 (18)
S1—C2—C3—C4	179.50 (17)	C3—C2—S1—C1B	-141.03 (17)
C2—C3—C4—C5	-0.1 (3)	C7—C2—S1—C1A	-36.62 (18)
C3—C4—C5—C6	0.5 (3)	C3—C2—S1—C1A	143.31 (16)
C4—C5—C6—C7	-0.1 (4)	C1B—S1—C1A—C9A	-17.3 (2)
C3—C2—C7—C6	0.9 (3)	C2—S1—C1A—C9A	78.0 (2)
S1—C2—C7—C6	-179.14 (14)	C9B—C10—C9A—C1A	-109.5 (6)
C3—C2—C7—N1	-178.87 (18)	C11—C10—C9A—C1A	-8.6 (6)
S1—C2—C7—N1	1.1 (3)	C9B—C10—C9A—C8	68.3 (4)
C5—C6—C7—C2	-0.6 (3)	C11—C10—C9A—C8	169.3 (2)
C5—C6—C7—N1	179.2 (2)	S1—C1A—C9A—C10	107.7 (4)
C9A—C10—C11—C12	-30.8 (4)	S1—C1A—C9A—C8	-70.1 (4)
C9B—C10—C11—C12	11.7 (4)	O1—C8—C9A—C10	34.1 (4)
C9A—C10—C11—C16	151.4 (3)	N1—C8—C9A—C10	-161.6 (3)
C9B—C10—C11—C16	-166.1 (3)	C9B—C8—C9A—C10	-71.9 (4)
C16—C11—C12—C13	1.9 (2)	O1—C8—C9A—C1A	-148.0 (3)
C10—C11—C12—C13	-175.93 (16)	N1—C8—C9A—C1A	16.3 (4)
C11—C12—C13—C14	-0.2 (3)	C9B—C8—C9A—C1A	105.9 (6)
C12—C13—C14—O2	177.88 (16)	C2—S1—C1B—C9B	-85.8 (2)
C12—C13—C14—C15	-1.9 (3)	C1A—S1—C1B—C9B	16.4 (2)

O2—C14—C15—C16	-177.57 (15)	C9A—C10—C9B—C1B	118.0 (6)
C13—C14—C15—C16	2.2 (3)	C11—C10—C9B—C1B	10.1 (6)
C14—C15—C16—C11	-0.4 (3)	C9A—C10—C9B—C8	-63.2 (4)
C12—C11—C16—C15	-1.6 (2)	C11—C10—C9B—C8	-171.13 (19)
C10—C11—C16—C15	176.40 (16)	S1—C1B—C9B—C10	-102.3 (4)
O1—C8—N1—C7	-179.9 (2)	S1—C1B—C9B—C8	79.0 (4)
C9B—C8—N1—C7	-16.1 (4)	O1—C8—C9B—C10	-32.2 (4)
C9A—C8—N1—C7	15.9 (4)	N1—C8—C9B—C10	164.6 (2)
C2—C7—N1—C8	-1.1 (4)	C9A—C8—C9B—C10	57.2 (4)
C6—C7—N1—C8	179.1 (2)	O1—C8—C9B—C1B	146.6 (3)
C13—C14—O2—C17	-7.0 (3)	N1—C8—C9B—C1B	-16.6 (4)
C15—C14—O2—C17	172.7 (2)	C9A—C8—C9B—C1B	-123.9 (6)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 ⁱ	0.86	2.07	2.9291 (18)	177
C6—H6...O1 ⁱ	0.93	2.45	3.263 (3)	146
C1B—H1C...O1 ⁱⁱ	0.97	2.52	3.377 (4)	147

Symmetry codes: (i) $-x, -y, -z$; (ii) $x, y+1, z$.