

# Crystal structure of ethyl (2*S*)-9-methoxy-2-methyl-4-oxo-3,4,5,6-tetrahydro-2*H*-2,6-methanobenzo[*g*][1,3,5]oxadiazocine-11-carboxylate

A. Dhandapani,<sup>a</sup> S. Manivarman,<sup>a</sup> S. Subashchandrabose<sup>b</sup> and B. Gunasekaran<sup>c\*</sup>

<sup>a</sup>Post Graduate and Research Department of Chemistry, Government Arts College, C-Mutlur, Chidambaram 608 102, Tamil Nadu, India, <sup>b</sup>Centre for Research & Development, PRIST University, Vallam, Thanjavur 613 403, Tamil Nadu, India, and <sup>c</sup>Department of Physics & Nano Technology, SRM University, SRM Nagar, Kattankulathur, Kancheepuram Dist, Chennai 603 203, Tamil Nadu, India.

\*Correspondence e-mail: phdguna@gmail.com

Received 3 January 2015; accepted 11 January 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


In the title compound, C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>, the methoxyphenyl ring makes a dihedral angle of 84.70 (12)° with the mean plane of the tetrahydropyrimidin-2(1*H*)-one ring. Both the pyran and tetrahydropyrimidin-2(1*H*)-one rings have distorted envelope conformations with the carboxylate-substituted C atom as the flap. In the crystal, molecules are linked *via* pairs of N—H...O hydrogen bonds, forming zigzag chains propagating along [010], which enclose R<sub>2</sub><sup>2</sup>(8) ring motifs. The chains are linked by C—H...π interactions, forming a two-dimensional network parallel to (100).

**Keywords:** crystal structure; hydroypyrimidine; oxadiazocine; pyran; hydrogen bonding.

**CCDC reference:** 1043105

## 1. Related literature

For the biological activity of dihydropyrimidine derivatives, see: Hurst & Hull (1961); Ashok *et al.* (2007); Bahekar & Shinde (2004); Mayer *et al.* (1999); Kappe (2000); For the crystal structures of two very similar compounds, see: Jing *et al.* (2009); Yar *et al.* (2014); Liu *et al.* (2014).



## 2. Experimental

### 2.1. Crystal data

C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>  
*M<sub>r</sub>* = 306.31  
 Monoclinic, *P*2<sub>1</sub>  
*a* = 9.6982 (14) Å  
*b* = 7.4802 (12) Å  
*c* = 10.8293 (17) Å  
 $\beta$  = 111.252 (5)°

*V* = 732.2 (2) Å<sup>3</sup>  
*Z* = 2  
 Mo *K*α radiation  
 $\mu$  = 0.11 mm<sup>-1</sup>  
*T* = 295 K  
 0.25 × 0.20 × 0.20 mm

### 2.2. Data collection

Bruker APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)  
*T<sub>min</sub>* = 0.954, *T<sub>max</sub>* = 0.975

10886 measured reflections  
 3030 independent reflections  
 2029 reflections with *I* > 2σ(*I*)  
*R<sub>int</sub>* = 0.053

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$   
 $wR(F^2) = 0.175$   
*S* = 1.14  
 3030 reflections  
 203 parameters  
 1 restraint  
 H-atom parameters constrained

$\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$   
 Absolute structure: Flack (1983),  
 1371 Friedel pairs  
 Absolute structure parameter:  
 0.01 (4)

**Table 1**

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C1–C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1–H1 $\cdots$ O2 <sup>i</sup>	0.86	2.12	2.962 (5)	168
N2–H2A $\cdots$ O2 <sup>ii</sup>	0.86	2.11	2.936 (4)	162
C14–H14B $\cdots$ Cg1 <sup>iii</sup>	0.96	2.62	3.570 (6)	171

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z$ ; (ii)  $-x + 1, y - \frac{1}{2}, -z$ ; (iii)  $-x + 1, y - \frac{1}{2}, -z + 1$ .

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU5057).

## References

- Ashok, M., Holla, B. S. & Kumari, N. S. (2007). *Eur. J. Med. Chem.* **42**, 380–385.
- Bahekar, S. S. & Shinde, D. B. (2004). *Bioorg. Med. Chem. Lett.* **14**, 1733–1736.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Hurst, E. W. & Hull, R. (1961). *J. Med. Chem.* **3**, 215–229.
- Jing, X., Li, Z., Pan, X., Wang, Q., Yan, C. & Zhu, H. (2009). *Synth. Commun.* **39**, 3796–3803.
- Kappe, C. O. (2000). *Eur. J. Med. Chem.* **35**, 1043–1052.
- Liu, Q., Xu, J., Teng, F., Chen, A., Pan, N. & Zhang, W. (2014). *J. Heterocycl. Chem.* **51**, 741–746.
- Mayer, T. U., Kapoor, T. M., Haggarty, S. J., King, R. W., Schreiber, S. L. & Mitchison, T. J. (1999). *Science*, **286**, 971–974.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Yar, M., Bajda, M., Shahzadi, L., Shahzad, S. A., Ahmed, M., Ashraf, M., Alam, U., Khan, I. U. & Khan, A. F. (2014). *Bioorg. Chem.* **54**, 96–104.

## supporting information

*Acta Cryst.* (2015). E71, o117–o118 [doi:10.1107/S2056989015000559]

## Crystal structure of ethyl (2*S*)-9-methoxy-2-methyl-4-oxo-3,4,5,6-tetrahydro-2*H*-2,6-methanobenzo[*g*][1,3,5]oxadiazocine-11-carboxylate

A. Dhandapani, S. Manivarman, S. Subashchandrabose and B. Gunasekaran

### S1. Comment

Dihydropyrimidine derivatives have recently received great attention because of their wide range of therapeutic and pharmacological properties, such as antiviral (Hurst & Hull, 1961), antitumor, antibacterial and antifungal (Ashok *et al.*, 2007), anti-inflammatory (Bahekar & Shinde, 2004), antihypertensive agents, and neuropeptide Y (NPY) antagonists (Mayer *et al.*, 1999). The natural products containing these heterocyclic moieties have been studied as new leads for AIDS therapies (Kappe, 2000).

The molecular structure of the title compound is illustrated in Fig. 1. The methoxyphenyl ring (C1-C6) makes a dihedral angles of 84.70 (12) ° with the mean plane of the tetrahydropyrimidin-2(1*H*)-one ring (N1/N2/C7-C10). Both the pyran (O1/C3/C4/C7/C10) and tetrahydropyrimidin-2(1*H*)-one rings have distorted envelope conformations with atom C10 as the flap.

The geometrical parameters of the title molecule agree well with those reported for two very similar compounds, *viz.* ethyl 2-methyl-4-oxo-3,4,5,6-tetrahydro-2*H*-2,6-methanobenzo[*g*][1,3,5]oxadiazocine-11-carboxylate (Jing *et al.*, 2009; Yar *et al.*, 2014) and methyl 2-methyl-4-oxo-3,4,5,6-tetrahydro-2*H*-2,6-methanobenzo[*g*][1,3,5]oxadiazocine-11-carboxylate (Liu *et al.*, 2014)


In the crystal, molecules are linked via pairs of N—H...O hydrogen bonds forming zigzag chains propagating along [010], which enclose R<sup>2</sup><sub>2</sub>(8) ring motifs (Table 1 and Fig. 2). The chains are linked by C-H... $\pi$  interactions forming a two-dimensional network parallel to (100).

### S2. Experimental

2-hydroxy-4-methoxybenzaldehyde (0.76 g, 5 mmol) and urea (0.9 g, 15 mmol) were added to an ethanolic solution of ethyl acetoacetate (0.65 ml, 5 mmol). To this mixture CeCl<sub>3</sub>·7H<sub>2</sub>O (0.465 g, 25%) was added slowly with stirring. After the addition was complete the reaction mixture was reflux at 363 K. The reaction mixture was then cooled to room temperature, then poured onto crushed ice water and stirred for 10 min. The solid that separated was filtered under suction, washed with water, and then recrystallized from DMSO giving colourless block-like crystals (yield: 96%; m.p.: 463 K).


### S3. Refinement

H atoms were positioned geometrically and refined using a riding model: N—H = 0.86 Å, C—H = 0.93 - 98 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$  for methyl H atoms and = 1.2 $U_{\text{eq}}(\text{N,C})$  for other H atoms.



**Figure 1**

The molecular structure of the title compound, with atom labelling. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

The crystal packing of the title compound, viewed along the *b* axis. Hydrogen bonds are shown as dashed lines (see Table 1 for details).

**Ethyl (2*S*)-9-methoxy-2-methyl-4-oxo-3,4,5,6-tetrahydro-2*H*-2,6-methanobenzo[*g*][1,3,5]oxadiazocine-11-carboxylate**

*Crystal data*

$C_{15}H_{18}N_2O_5$

$M_r = 306.31$

Monoclinic,  $P2_1$

Hall symbol:  $P\ 2_1y$

$a = 9.6982\ (14)\ \text{\AA}$

$b = 7.4802\ (12)\ \text{\AA}$

$c = 10.8293\ (17)\ \text{\AA}$

$\beta = 111.252\ (5)^\circ$

$V = 732.2\ (2)\ \text{\AA}^3$

$Z = 2$

$F(000) = 324$

$D_x = 1.389\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3030 reflections

$\theta = 2.0\text{--}26.8^\circ$

$\mu = 0.11\ \text{mm}^{-1}$

$T = 295\ \text{K}$

Block, colourless

$0.25 \times 0.20 \times 0.20\ \text{mm}$

*Data collection*

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 0 pixels  $\text{mm}^{-1}$

$\omega$  and  $\varphi$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.954$ ,  $T_{\max} = 0.975$

10886 measured reflections

3030 independent reflections

2029 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.053$

$\theta_{\max} = 26.9^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -11 \rightarrow 12$

$k = -9 \rightarrow 9$

$l = -13 \rightarrow 13$

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.071$  $wR(F^2) = 0.175$  $S = 1.14$ 

3030 reflections

203 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0537P)^2 + 0.7506P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$ Extinction correction: *SHELXL97* (Sheldrick,  
2008)

Extinction coefficient: 0.031 (8)

Absolute structure: Flack (1983), 1371 Friedel  
pairs

Absolute structure parameter: 0.01 (4)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.9053 (5)	0.2379 (7)	0.5750 (5)	0.0437 (13)
C2	0.8132 (5)	0.0956 (7)	0.5239 (5)	0.0412 (12)
H2	0.8430	-0.0198	0.5535	0.049*
C3	0.6753 (5)	0.1257 (6)	0.4278 (4)	0.0325 (10)
C4	0.6293 (5)	0.2978 (6)	0.3837 (4)	0.0309 (10)
C5	0.7245 (6)	0.4362 (6)	0.4375 (5)	0.0375 (11)
H5	0.6948	0.5519	0.4085	0.045*
C6	0.8629 (6)	0.4105 (7)	0.5331 (5)	0.0455 (13)
H6	0.9257	0.5066	0.5682	0.055*
C7	0.4765 (5)	0.3254 (6)	0.2807 (4)	0.0299 (10)
H7	0.4353	0.4379	0.2984	0.036*
C8	0.4926 (4)	0.1772 (6)	0.0871 (4)	0.0302 (9)
C9	0.4544 (5)	0.0033 (6)	0.2670 (4)	0.0295 (10)
C10	0.3772 (4)	0.1706 (6)	0.2878 (4)	0.0308 (9)
H10	0.3722	0.1671	0.3765	0.037*
C11	0.2243 (5)	0.2001 (7)	0.1873 (5)	0.0412 (11)
C12	0.0178 (7)	0.3958 (9)	0.1308 (6)	0.0700 (19)
H12A	0.0121	0.3701	0.0413	0.084*
H12B	0.0123	0.5244	0.1396	0.084*
C13	-0.1071 (7)	0.3110 (11)	0.1540 (7)	0.088 (2)
H13A	-0.1119	0.1870	0.1299	0.132*
H13B	-0.1974	0.3693	0.1011	0.132*
H13C	-0.0937	0.3212	0.2460	0.132*
C14	0.3703 (5)	-0.1689 (6)	0.2594 (5)	0.0398 (12)
H14A	0.2826	-0.1685	0.1810	0.060*
H14B	0.3432	-0.1792	0.3361	0.060*
H14C	0.4316	-0.2683	0.2564	0.060*
C15	1.1301 (7)	0.3274 (10)	0.7503 (6)	0.074 (2)
H15A	1.1513	0.4157	0.6953	0.111*
H15B	1.2209	0.2756	0.8085	0.111*
H15C	1.0794	0.3828	0.8017	0.111*

N1	0.4778 (4)	0.3287 (5)	0.1464 (3)	0.0341 (9)
H1	0.4690	0.4288	0.1051	0.041*
N2	0.4908 (4)	0.0226 (4)	0.1509 (3)	0.0326 (9)
H2A	0.5136	-0.0733	0.1187	0.039*
O1	0.5875 (3)	-0.0211 (4)	0.3833 (3)	0.0343 (8)
O2	0.5039 (3)	0.1774 (4)	-0.0236 (3)	0.0383 (8)
O3	0.1709 (4)	0.1240 (6)	0.0862 (4)	0.0817 (15)
O4	0.1593 (4)	0.3304 (6)	0.2256 (4)	0.0650 (12)
O5	1.0392 (4)	0.1917 (6)	0.6693 (4)	0.0682 (12)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.033 (3)	0.059 (3)	0.037 (3)	0.002 (2)	0.010 (2)	-0.003 (2)
C2	0.044 (3)	0.040 (3)	0.039 (3)	0.006 (2)	0.014 (2)	0.003 (2)
C3	0.040 (3)	0.028 (2)	0.031 (2)	-0.001 (2)	0.016 (2)	-0.0004 (18)
C4	0.039 (3)	0.028 (2)	0.028 (2)	0.003 (2)	0.015 (2)	0.0012 (19)
C5	0.046 (3)	0.033 (3)	0.038 (3)	-0.003 (2)	0.020 (2)	0.000 (2)
C6	0.046 (3)	0.044 (3)	0.048 (3)	-0.015 (3)	0.018 (3)	-0.008 (2)
C7	0.042 (3)	0.023 (2)	0.028 (2)	0.000 (2)	0.017 (2)	-0.0006 (18)
C8	0.039 (2)	0.026 (2)	0.029 (2)	0.001 (2)	0.0161 (18)	0.000 (2)
C9	0.037 (2)	0.025 (2)	0.028 (2)	0.000 (2)	0.013 (2)	0.0020 (18)
C10	0.036 (2)	0.031 (2)	0.030 (2)	0.001 (2)	0.0173 (18)	-0.002 (2)
C11	0.042 (3)	0.040 (3)	0.042 (3)	0.002 (2)	0.016 (2)	-0.004 (2)
C12	0.060 (4)	0.077 (4)	0.065 (4)	0.028 (4)	0.014 (3)	0.009 (3)
C13	0.060 (4)	0.084 (5)	0.099 (5)	0.006 (4)	0.003 (4)	0.011 (5)
C14	0.051 (3)	0.030 (2)	0.047 (3)	-0.007 (2)	0.028 (2)	-0.002 (2)
C15	0.057 (4)	0.089 (5)	0.055 (3)	-0.020 (4)	-0.006 (3)	0.004 (4)
N1	0.058 (3)	0.0218 (18)	0.0261 (19)	0.0069 (18)	0.0197 (18)	0.0055 (15)
N2	0.050 (2)	0.0202 (19)	0.036 (2)	0.0038 (17)	0.0244 (19)	0.0013 (15)
O1	0.0418 (18)	0.0264 (16)	0.0315 (16)	0.0019 (14)	0.0092 (14)	0.0033 (13)
O2	0.062 (2)	0.0286 (16)	0.0309 (15)	-0.0006 (17)	0.0244 (14)	0.0015 (14)
O3	0.058 (3)	0.089 (3)	0.071 (3)	0.019 (2)	-0.011 (2)	-0.041 (3)
O4	0.060 (2)	0.074 (3)	0.051 (2)	0.037 (2)	0.0092 (19)	-0.004 (2)
O5	0.044 (2)	0.073 (3)	0.066 (2)	0.007 (2)	-0.0064 (19)	-0.008 (2)

*Geometric parameters (Å, °)*

C1—C2	1.371 (7)	C9—C10	1.516 (6)
C1—O5	1.373 (6)	C10—C11	1.504 (6)
C1—C6	1.381 (7)	C10—H10	0.9800
C2—C3	1.383 (6)	C11—O3	1.174 (5)
C2—H2	0.9300	C11—O4	1.308 (6)
C3—O1	1.366 (5)	C12—O4	1.468 (7)
C3—C4	1.389 (6)	C12—C13	1.468 (9)
C4—C5	1.369 (6)	C12—H12A	0.9700
C4—C7	1.511 (6)	C12—H12B	0.9700
C5—C6	1.379 (7)	C13—H13A	0.9600

C5—H5	0.9300	C13—H13B	0.9600
C6—H6	0.9300	C13—H13C	0.9600
C7—N1	1.459 (5)	C14—H14A	0.9600
C7—C10	1.526 (6)	C14—H14B	0.9600
C7—H7	0.9800	C14—H14C	0.9600
C8—O2	1.242 (5)	C15—O5	1.420 (7)
C8—N1	1.336 (5)	C15—H15A	0.9600
C8—N2	1.351 (5)	C15—H15B	0.9600
C9—N2	1.431 (6)	C15—H15C	0.9600
C9—O1	1.452 (5)	N1—H1	0.8600
C9—C14	1.510 (6)	N2—H2A	0.8600
C2—C1—O5	114.0 (5)	C7—C10—H10	109.1
C2—C1—C6	121.3 (4)	O3—C11—O4	123.8 (5)
O5—C1—C6	124.6 (5)	O3—C11—C10	126.1 (4)
C1—C2—C3	119.3 (4)	O4—C11—C10	110.0 (4)
C1—C2—H2	120.4	O4—C12—C13	110.9 (5)
C3—C2—H2	120.4	O4—C12—H12A	109.5
O1—C3—C2	116.3 (4)	C13—C12—H12A	109.5
O1—C3—C4	122.9 (4)	O4—C12—H12B	109.5
C2—C3—C4	120.8 (4)	C13—C12—H12B	109.5
C5—C4—C3	118.1 (4)	H12A—C12—H12B	108.1
C5—C4—C7	122.8 (4)	C12—C13—H13A	109.5
C3—C4—C7	119.1 (4)	C12—C13—H13B	109.5
C4—C5—C6	122.5 (4)	H13A—C13—H13B	109.5
C4—C5—H5	118.8	C12—C13—H13C	109.5
C6—C5—H5	118.8	H13A—C13—H13C	109.5
C5—C6—C1	118.1 (5)	H13B—C13—H13C	109.5
C5—C6—H6	121.0	C9—C14—H14A	109.5
C1—C6—H6	121.0	C9—C14—H14B	109.5
N1—C7—C4	112.2 (4)	H14A—C14—H14B	109.5
N1—C7—C10	107.3 (3)	C9—C14—H14C	109.5
C4—C7—C10	109.1 (3)	H14A—C14—H14C	109.5
N1—C7—H7	109.4	H14B—C14—H14C	109.5
C4—C7—H7	109.4	O5—C15—H15A	109.5
C10—C7—H7	109.4	O5—C15—H15B	109.5
O2—C8—N1	121.7 (4)	H15A—C15—H15B	109.5
O2—C8—N2	121.1 (4)	O5—C15—H15C	109.5
N1—C8—N2	117.2 (3)	H15A—C15—H15C	109.5
N2—C9—O1	110.5 (3)	H15B—C15—H15C	109.5
N2—C9—C14	109.8 (3)	C8—N1—C7	120.4 (4)
O1—C9—C14	103.5 (3)	C8—N1—H1	119.8
N2—C9—C10	109.8 (3)	C7—N1—H1	119.8
O1—C9—C10	107.8 (3)	C8—N2—C9	126.1 (3)
C14—C9—C10	115.2 (4)	C8—N2—H2A	117.0
C11—C10—C9	115.1 (4)	C9—N2—H2A	117.0
C11—C10—C7	109.1 (4)	C3—O1—C9	116.7 (3)
C9—C10—C7	105.3 (3)	C11—O4—C12	117.5 (4)



C11—C10—H10	109.1	C1—O5—C15	119.1 (5)
C9—C10—H10	109.1		
O5—C1—C2—C3	-179.8 (4)	N1—C7—C10—C9	64.6 (4)
C6—C1—C2—C3	0.4 (7)	C4—C7—C10—C9	-57.2 (4)
C1—C2—C3—O1	-178.5 (4)	C9—C10—C11—O3	-12.8 (7)
C1—C2—C3—C4	-0.5 (7)	C7—C10—C11—O3	105.3 (6)
O1—C3—C4—C5	178.2 (4)	C9—C10—C11—O4	170.0 (4)
C2—C3—C4—C5	0.4 (6)	C7—C10—C11—O4	-71.9 (5)
O1—C3—C4—C7	-0.9 (6)	O2—C8—N1—C7	-175.1 (4)
C2—C3—C4—C7	-178.7 (4)	N2—C8—N1—C7	6.6 (6)
C3—C4—C5—C6	-0.2 (7)	C4—C7—N1—C8	75.5 (5)
C7—C4—C5—C6	178.9 (4)	C10—C7—N1—C8	-44.2 (5)
C4—C5—C6—C1	0.0 (7)	O2—C8—N2—C9	-169.1 (4)
C2—C1—C6—C5	-0.2 (7)	N1—C8—N2—C9	9.1 (6)
O5—C1—C6—C5	-179.9 (5)	O1—C9—N2—C8	-103.2 (5)
C5—C4—C7—N1	87.0 (5)	C14—C9—N2—C8	143.3 (4)
C3—C4—C7—N1	-94.0 (5)	C10—C9—N2—C8	15.7 (6)
C5—C4—C7—C10	-154.3 (4)	C2—C3—O1—C9	-169.5 (4)
C3—C4—C7—C10	24.7 (5)	C4—C3—O1—C9	12.6 (6)
N2—C9—C10—C11	69.2 (5)	N2—C9—O1—C3	72.7 (4)
O1—C9—C10—C11	-170.3 (3)	C14—C9—O1—C3	-169.9 (4)
C14—C9—C10—C11	-55.3 (5)	C10—C9—O1—C3	-47.4 (5)
N2—C9—C10—C7	-50.9 (4)	O3—C11—O4—C12	-6.2 (8)
O1—C9—C10—C7	69.5 (4)	C10—C11—O4—C12	171.1 (5)
C14—C9—C10—C7	-175.5 (4)	C13—C12—O4—C11	96.3 (7)
N1—C7—C10—C11	-59.5 (4)	C2—C1—O5—C15	-166.0 (5)
C4—C7—C10—C11	178.8 (3)	C6—C1—O5—C15	13.8 (8)

*Hydrogen-bond geometry (Å, °)*

Cg1 is the centroid of the C1—C6 ring.

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...O2 <sup>i</sup>	0.86	2.12	2.962 (5)	168
N2—H2A...O2 <sup>ii</sup>	0.86	2.11	2.936 (4)	162
C14—H14B...Cg1 <sup>iii</sup>	0.96	2.62	3.570 (6)	171

Symmetry codes: (i)  $-x+1, y+1/2, -z$ ; (ii)  $-x+1, y-1/2, -z$ ; (iii)  $-x+1, y-1/2, -z+1$ .