

# Crystal structure of ethyl 6-methyl-2-sulfanylidene-4-(thiophen-2-yl)-1,2,3,4-tetrahydropyrimidine-5-carboxylate

M. Suresh,<sup>a</sup> M. Syed Ali Padusha,<sup>a</sup> J. Josephine Novina,<sup>b</sup> G. Vasuki,<sup>c\*</sup> Vijayan Viswanathan<sup>d</sup> and Devadasan Velmurugan<sup>d</sup>

<sup>a</sup>PG & Research Department of Chemistry, Jamal Mohamed College (Autonomous), Tiruchirappalli-20, India, <sup>b</sup>Department of Physics, Idhaya College for Women, Kumbakonam-1, India, <sup>c</sup>Department of Physics, Kunthavai Naachiar Government Arts College (W) (Autonomous), Thanjavur-7, India, and <sup>d</sup>Centre of Advanced study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai-25, India. \*Correspondence e-mail: vasuki.arasi@yahoo.com

Received 18 December 2014; accepted 19 December 2014

Edited by E. R. T. Tiekink, University of Malaya, Malaysia


In the title compound, C<sub>12</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>S<sub>2</sub>, the dihydropyrimidine ring adopts a sofa conformation, with the C atom bearing the thienyl ring lying above the plane of the five remaining approximately coplanar (r.m.s. deviation = 0.0405 Å) atoms of the ring. The dihedral angle between the five near coplanar atoms of the ring and the thienyl ring is 89.78 (11)°. In the crystal, molecules are linked into a supramolecular chain along [100] via N—H···O(carbonyl) hydrogen bonds. Inversion-related chains are linked into double chains via N—H···S(thione) hydrogen bonds. The three-dimensional architecture also features methyl–thienyl C—H···π interactions.

**Keywords:** crystal structure; pyrimidine; hydrogen bonding; C—H···π interactions; conformation.

**CCDC reference:** 1040426

## 1. Related literature

For general background and the biological activity of dihydropyrimidinones, see: Phucho *et al.* (2009); Patil *et al.* (2011).



## 2. Experimental

### 2.1. Crystal data

|  |   |
|--|---|
| C <sub>12</sub> H <sub>14</sub> N <sub>2</sub> O <sub>2</sub> S <sub>2</sub> | $\gamma = 101.276 (1)^\circ$              |
| $M_r = 282.37$   | $V = 668.18 (1) \text{ \AA}^3$            |
| Triclinic, $P\bar{1}$  | $Z = 2$                                   |
| $a = 7.3069 (1) \text{ \AA}$   | Mo $K\alpha$ radiation                    |
| $b = 8.3267 (1) \text{ \AA}$   | $\mu = 0.39 \text{ mm}^{-1}$              |
| $c = 11.2461 (1) \text{ \AA}$  | $T = 293 \text{ K}$                       |
| $\alpha = 90.109 (1)^\circ$  | $0.20 \times 0.15 \times 0.10 \text{ mm}$ |
| $\beta = 95.156 (1)^\circ$   |   |

### 2.2. Data collection

|  |  |
|--|--|
| Bruker APEXII CCD diffractometer                         | 10188 measured reflections             |
| Absorption correction: multi-scan (SADABS; Bruker, 2008) | 2762 independent reflections           |
| $T_{\min} = 0.925$ , $T_{\max} = 0.962$                  | 2325 reflections with $I > 2\sigma(I)$ |
|  | $R_{\text{int}} = 0.021$               |

### 2.3. Refinement

|                                 |  |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.060$ | 165 parameters                                       |
| $wR(F^2) = 0.189$               | H-atom parameters constrained                        |
| $S = 1.06$                      | $\Delta\rho_{\text{max}} = 0.64 \text{ e \AA}^{-3}$  |
| 2762 reflections                | $\Delta\rho_{\text{min}} = -0.40 \text{ e \AA}^{-3}$ |

**Table 1**

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the S1/C1–C4 thiophene ring.

| $D-H\cdots A$                 | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------------|-------|-------------|-------------|---------------|
| N1—H1N1···S2 <sup>i</sup>     | 0.86  | 2.63        | 3.408 (2)   | 151           |
| N2—H2N2···O1 <sup>ii</sup>    | 0.86  | 2.15        | 2.984 (3)   | 162           |
| C12—H12B···Cg1 <sup>iii</sup> | 0.96  | 2.81        | 3.664 (6)   | 149           |

Symmetry codes: (i)  $-x + 2, -y, -z + 1$ ; (ii)  $x + 1, y, z$ ; (iii)  $-x + 1, -y + 1, -z$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: APEX2 and SAINT (Bruker, 2008); data reduction: SAINT and XPREP (Bruker, 2008); program(s) used to solve structure: SIR92 (Altomare *et al.*, 1993); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: PLATON (Spek, 2009).

### Acknowledgements

The authors thank the TBI X-ray facility, CAS in Crystallography and Biophysics, University of Madras, India, for the data collection.

---

Supporting information for this paper is available from the IUCr electronic archives (Reference: TK5352).

---

### References

Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.

Bruker (2008). *APEX2, SADABS, SAINT and XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.

Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.

Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.

Patil, S., Jadhav, S. D. & Deshmukh, M. B. (2011). *Arch. Appl. Sci. Res.* **3**, 203–208.

Phucho, I. T., Nongpiur, A., Tumtin, S., Nongrum, R. & Nongkhlaw, R. L. (2009). *Rasayan J. Chem.* **2**, 662–676.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2015). E71, o81–o82 [doi:10.1107/S2056989014027741]

## Crystal structure of ethyl 6-methyl-2-sulfanylidene-4-(thiophen-2-yl)-1,2,3,4-tetrahydropyrimidine-5-carboxylate

M. Suresh, M. Syed Ali Padusha, J. Josephine Novina, G. Vasuki, Vijayan Viswanathan and Devadasan Velmurugan

### S1. Structural commentary

Pyrimidinones or dihydropyrimidinones (DHPMs) are well known for their wide range of bioactivities and their applications in the field of drug research have stimulated the invention of a wide range of synthetic methods for their preparation and chemical transformations (Phucho *et al.*, 2009). Several functionalized dihydropyrimidinones are used as calcium channel modulators, Ca-antagonists, vasodilative and anti-hypertensive agents (Patil *et al.*, 2011). Against this background and in order to obtain detailed information on its molecular conformation, the structure of the title compound has been determined and the results are presented herein.

The asymmetric unit of the title compound is illustrated in Fig. 1. The dihydropyrimidine ring adopts a sofa conformation, with puckering parameters  $q_2 = 0.283 \text{ \AA}$ ,  $q_3 = -0.102 \text{ \AA}$ ,  $Q = 0.301 \text{ \AA}$ ,  $\Theta = 109.8^\circ$  and  $\Phi = 232.1^\circ$ . The dihedral angle between the mean plane of the five essentially planar atoms (N1/C9/N2/C7/C6) of the dihydropyrimidine ring [maximum deviation  $0.1944 (18) \text{ \AA}$  for C5] and the thiophene ring (C1—C4/S1) is  $89.78 (11)^\circ$ .


In the crystal, the molecules are linked *via* a pair of N—H $\cdots$ S hydrogen bonds forming an inversion dimers with  $R^2_2(8)$  ring motif and form a chain parallel to the *bc* plane *via* N—H $\cdots$ O hydrogen bonds (Fig. 2). Additional stabilization to the structure is afforded by C—H $\cdots$  $\pi$  contacts (Table 1, Figs 3 and 4).

### S2. Synthesis and crystallization


A mixture of ethyl acetoacetate (0.13 ml, 0.001 mol), thiophene-2-carboxaldehyde (0.1 ml, 0.001 mol) and thiourea (0.228 g, 0.003 mol) in ethanol (5 ml) was heated under reflux in the presence of cerium chloride heptahydrate (25%) for 1 h (monitored by TLC). After the completion of the reaction, the reaction mixture was cooled to room temperature and poured onto crushed ice and stirred for 5–10 min. The solid was separated and filtered under suction, washed with ice-cold water (50 ml), and then recrystallized from hot ethanol to afford pure product [m.pt: 421 K; yield: 98%].

### S3. Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances fixed in the range  $0.93\text{--}0.98 \text{ \AA}$  and N—H =  $0.86 \text{ \AA}$  with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{CH}_3)$  and  $1.2U_{\text{eq}}(\text{CH}_2, \text{CH}, \text{NH})$ .


**Figure 1**

The molecular structure of the title compound, with the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.



**Figure 2**

Partial crystal packing of the title compound, showing the  $R^2_2(8)$  ring motif, viewed along the *b* axis. Hydrogen bonds are shown as dashed lines.



**Figure 3**

Part of the crystal packing of the title compound, showing C—H... $\pi$  interactions. Viewed along the *a* axis.




Figure 4

Image showing the C—H $\cdots$  $\pi$  interactions.

### Ethyl 6-methyl-2-sulfanylidene-4-(thiophen-2-yl)-1,2,3,4-tetrahydropyrimidine-5-carboxylate

#### Crystal data

$C_{12}H_{14}N_2O_2S_2$

$M_r = 282.37$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 7.3069$  (1) Å

$b = 8.3267$  (1) Å

$c = 11.2461$  (1) Å

$\alpha = 90.109$  (1)°

$\beta = 95.156$  (1)°

$\gamma = 101.276$  (1)°

$V = 668.18$  (1) Å<sup>3</sup>

$Z = 2$

$F(000) = 296$

$D_x = 1.403$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2762 reflections

$\theta = 1.8$ – $26.5$ °

$\mu = 0.39$  mm<sup>-1</sup>

$T = 293$  K  $0.20 \times 0.15 \times 0.10$  mm  
 Block, colourless

*Data collection*

|  |  |
|--|--|
| Bruker APEXII CCD diffractometer                         | 10188 measured reflections   |
| Radiation source: fine-focus sealed tube                 | 2762 independent reflections   |
| Graphite monochromator                                   | 2325 reflections with $I > 2\sigma(I)$                                 |
| $\omega$ and $\varphi$ scan                              | $R_{\text{int}} = 0.021$   |
| Absorption correction: multi-scan (SADABS; Bruker, 2008) | $\theta_{\text{max}} = 26.5^\circ$ , $\theta_{\text{min}} = 1.8^\circ$ |
| $T_{\text{min}} = 0.925$ , $T_{\text{max}} = 0.962$      | $h = -9 \rightarrow 9$   |
|  | $k = -10 \rightarrow 10$   |
|  | $l = -14 \rightarrow 14$   |

*Refinement*

|  |  |
|--|--|
| Refinement on $F^2$  | Secondary atom site location: difference Fourier map         |
| Least-squares matrix: full                                     | Hydrogen site location: inferred from neighbouring sites     |
| $R[F^2 > 2\sigma(F^2)] = 0.060$                                | H-atom parameters constrained                                |
| $wR(F^2) = 0.189$  | $w = 1/[\sigma^2(F_o^2) + (0.1181P)^2 + 0.5067P]$            |
| $S = 1.06$   | where $P = (F_o^2 + 2F_c^2)/3$                               |
| 2762 reflections   | $(\Delta/\sigma)_{\text{max}} < 0.001$                       |
| 165 parameters   | $\Delta\rho_{\text{max}} = 0.64 \text{ e } \text{\AA}^{-3}$  |
| 0 restraints   | $\Delta\rho_{\text{min}} = -0.40 \text{ e } \text{\AA}^{-3}$ |
| Primary atom site location: structure-invariant direct methods |  |

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | $x$        | $y$        | $z$         | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|------------|------------|-------------|----------------------------------|
| C1  | 0.8629 (5) | 0.6238 (4) | 0.4026 (3)  | 0.0584 (9)                       |
| H1  | 0.9131     | 0.7349     | 0.4141      | 0.070*                           |
| C2  | 0.7502 (5) | 0.5376 (4) | 0.4750 (3)  | 0.0534 (8)                       |
| H2  | 0.7143     | 0.5833     | 0.5425      | 0.064*                           |
| C3  | 0.6877 (4) | 0.3688 (3) | 0.4414 (2)  | 0.0341 (5)                       |
| H3  | 0.6069     | 0.2922     | 0.4819      | 0.041*                           |
| C4  | 0.7703 (3) | 0.3372 (3) | 0.3350 (2)  | 0.0338 (5)                       |
| C5  | 0.7577 (3) | 0.1734 (3) | 0.2747 (2)  | 0.0336 (6)                       |
| H5  | 0.6314     | 0.1088     | 0.2803      | 0.040*                           |
| C9  | 1.0680 (4) | 0.0949 (3) | 0.3047 (2)  | 0.0335 (5)                       |
| C7  | 0.9646 (4) | 0.1925 (4) | 0.1110 (2)  | 0.0384 (6)                       |
| C8  | 1.0343 (5) | 0.2269 (5) | -0.0103 (3) | 0.0581 (9)                       |
| H8A | 0.9320     | 0.1960     | -0.0709     | 0.087*                           |
| H8B | 1.1295     | 0.1651     | -0.0216     | 0.087*                           |



|      |              |              |              |             |
|------|--------------|--------------|--------------|-------------|
| H8C  | 1.0856       | 0.3417       | -0.0157      | 0.087*      |
| C6   | 0.7909 (4)   | 0.1904 (4)   | 0.1437 (2)   | 0.0371 (6)  |
| C10  | 0.6280 (4)   | 0.2135 (4)   | 0.0657 (2)   | 0.0439 (7)  |
| C11  | 0.5131 (5)   | 0.2937 (7)   | -0.1262 (3)  | 0.0817 (14) |
| H11A | 0.4303       | 0.1894       | -0.1465      | 0.098*      |
| H11B | 0.4413       | 0.3655       | -0.0918      | 0.098*      |
| C12  | 0.5900 (7)   | 0.3646 (9)   | -0.2304 (4)  | 0.115 (2)   |
| H12A | 0.6581       | 0.4738       | -0.2117      | 0.172*      |
| H12B | 0.4906       | 0.3682       | -0.2916      | 0.172*      |
| H12C | 0.6732       | 0.2996       | -0.2582      | 0.172*      |
| N1   | 0.8939 (3)   | 0.0838 (3)   | 0.33419 (19) | 0.0359 (5)  |
| H1N1 | 0.8586       | 0.0204       | 0.3916       | 0.043*      |
| N2   | 1.1028 (3)   | 0.1610 (3)   | 0.1964 (2)   | 0.0402 (5)  |
| H2N2 | 1.2174       | 0.1848       | 0.1796       | 0.048*      |
| O1   | 0.4710 (3)   | 0.1889 (4)   | 0.0948 (2)   | 0.0673 (8)  |
| O2   | 0.6667 (3)   | 0.2699 (4)   | -0.0409 (2)  | 0.0718 (9)  |
| S1   | 0.90646 (12) | 0.51126 (11) | 0.28681 (8)  | 0.0557 (3)  |
| S2   | 1.23669 (10) | 0.02700 (10) | 0.38956 (6)  | 0.0442 (3)  |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$    | $U^{13}$     | $U^{23}$     |
|-----|-------------|-------------|-------------|-------------|--------------|--------------|
| C1  | 0.0543 (19) | 0.0410 (17) | 0.077 (2)   | 0.0109 (14) | -0.0128 (17) | 0.0012 (16)  |
| C2  | 0.0529 (19) | 0.059 (2)   | 0.0520 (17) | 0.0243 (15) | -0.0017 (14) | -0.0055 (15) |
| C3  | 0.0345 (13) | 0.0374 (13) | 0.0329 (12) | 0.0136 (10) | 0.0015 (10)  | 0.0007 (10)  |
| C4  | 0.0287 (12) | 0.0411 (14) | 0.0324 (12) | 0.0089 (10) | 0.0026 (9)   | 0.0086 (10)  |
| C5  | 0.0271 (12) | 0.0441 (14) | 0.0303 (12) | 0.0075 (10) | 0.0045 (9)   | 0.0077 (10)  |
| C9  | 0.0328 (13) | 0.0345 (13) | 0.0340 (12) | 0.0074 (10) | 0.0051 (10)  | 0.0073 (10)  |
| C7  | 0.0320 (13) | 0.0538 (16) | 0.0308 (12) | 0.0105 (11) | 0.0058 (10)  | 0.0089 (11)  |
| C8  | 0.0421 (16) | 0.101 (3)   | 0.0375 (15) | 0.0257 (17) | 0.0135 (12)  | 0.0224 (16)  |
| C6  | 0.0310 (13) | 0.0497 (16) | 0.0311 (13) | 0.0083 (11) | 0.0044 (10)  | 0.0047 (11)  |
| C10 | 0.0327 (14) | 0.0662 (19) | 0.0343 (13) | 0.0121 (12) | 0.0058 (11)  | 0.0096 (12)  |
| C11 | 0.0403 (18) | 0.156 (5)   | 0.0490 (19) | 0.023 (2)   | -0.0035 (15) | 0.034 (2)    |
| C12 | 0.061 (3)   | 0.214 (7)   | 0.077 (3)   | 0.041 (3)   | 0.010 (2)    | 0.066 (4)    |
| N1  | 0.0350 (11) | 0.0430 (12) | 0.0333 (11) | 0.0128 (9)  | 0.0101 (9)   | 0.0120 (9)   |
| N2  | 0.0272 (11) | 0.0596 (15) | 0.0360 (11) | 0.0117 (10) | 0.0077 (9)   | 0.0138 (10)  |
| O1  | 0.0319 (11) | 0.126 (2)   | 0.0474 (12) | 0.0213 (13) | 0.0070 (9)   | 0.0226 (14)  |
| O2  | 0.0369 (12) | 0.138 (3)   | 0.0425 (12) | 0.0217 (14) | 0.0055 (9)   | 0.0377 (14)  |
| S1  | 0.0519 (5)  | 0.0531 (5)  | 0.0613 (5)  | 0.0053 (4)  | 0.0108 (4)   | 0.0143 (4)   |
| S2  | 0.0372 (4)  | 0.0567 (5)  | 0.0429 (4)  | 0.0181 (3)  | 0.0060 (3)   | 0.0170 (3)   |

*Geometric parameters (Å, °)*

|       |           |        |           |
|-------|-----------|--------|-----------|
| C1—C2 | 1.322 (5) | C7—C8  | 1.508 (4) |
| C1—S1 | 1.691 (4) | C8—H8A | 0.9600    |
| C1—H1 | 0.9300    | C8—H8B | 0.9600    |
| C2—C3 | 1.429 (4) | C8—H8C | 0.9600    |
| C2—H2 | 0.9300    | C6—C10 | 1.458 (4) |

|             |             |               |            |
|-------------|-------------|---------------|------------|
| C3—C4       | 1.434 (3)   | C10—O1        | 1.200 (3)  |
| C3—H3       | 0.9300      | C10—O2        | 1.323 (3)  |
| C4—C5       | 1.505 (4)   | C11—C12       | 1.425 (6)  |
| C4—S1       | 1.709 (3)   | C11—O2        | 1.454 (4)  |
| C5—N1       | 1.472 (3)   | C11—H11A      | 0.9700     |
| C5—C6       | 1.516 (3)   | C11—H11B      | 0.9700     |
| C5—H5       | 0.9800      | C12—H12A      | 0.9600     |
| C9—N1       | 1.329 (3)   | C12—H12B      | 0.9600     |
| C9—N2       | 1.363 (3)   | C12—H12C      | 0.9600     |
| C9—S2       | 1.677 (3)   | N1—H1N1       | 0.8600     |
| C7—C6       | 1.350 (4)   | N2—H2N2       | 0.8600     |
| C7—N2       | 1.393 (3)   |               |            |
|             |             |               |            |
| C2—C1—S1    | 113.1 (3)   | H8A—C8—H8C    | 109.5      |
| C2—C1—H1    | 123.5       | H8B—C8—H8C    | 109.5      |
| S1—C1—H1    | 123.5       | C7—C6—C10     | 126.4 (2)  |
| C1—C2—C3    | 114.9 (3)   | C7—C6—C5      | 119.0 (2)  |
| C1—C2—H2    | 122.6       | C10—C6—C5     | 114.5 (2)  |
| C3—C2—H2    | 122.6       | O1—C10—O2     | 121.5 (3)  |
| C2—C3—C4    | 108.7 (3)   | O1—C10—C6     | 124.0 (3)  |
| C2—C3—H3    | 125.7       | O2—C10—C6     | 114.5 (2)  |
| C4—C3—H3    | 125.7       | C12—C11—O2    | 108.3 (3)  |
| C3—C4—C5    | 126.9 (2)   | C12—C11—H11A  | 110.0      |
| C3—C4—S1    | 111.2 (2)   | O2—C11—H11A   | 110.0      |
| C5—C4—S1    | 121.73 (19) | C12—C11—H11B  | 110.0      |
| N1—C5—C4    | 110.9 (2)   | O2—C11—H11B   | 110.0      |
| N1—C5—C6    | 109.0 (2)   | H11A—C11—H11B | 108.4      |
| C4—C5—C6    | 111.9 (2)   | C11—C12—H12A  | 109.5      |
| N1—C5—H5    | 108.3       | C11—C12—H12B  | 109.5      |
| C4—C5—H5    | 108.3       | H12A—C12—H12B | 109.5      |
| C6—C5—H5    | 108.3       | C11—C12—H12C  | 109.5      |
| N1—C9—N2    | 115.7 (2)   | H12A—C12—H12C | 109.5      |
| N1—C9—S2    | 123.97 (19) | H12B—C12—H12C | 109.5      |
| N2—C9—S2    | 120.33 (19) | C9—N1—C5      | 124.1 (2)  |
| C6—C7—N2    | 119.0 (2)   | C9—N1—H1N1    | 118.0      |
| C6—C7—C8    | 128.0 (2)   | C5—N1—H1N1    | 118.0      |
| N2—C7—C8    | 113.1 (2)   | C9—N2—C7      | 124.1 (2)  |
| C7—C8—H8A   | 109.5       | C9—N2—H2N2    | 117.9      |
| C7—C8—H8B   | 109.5       | C7—N2—H2N2    | 117.9      |
| H8A—C8—H8B  | 109.5       | C10—O2—C11    | 118.6 (2)  |
| C7—C8—H8C   | 109.5       | C1—S1—C4      | 92.16 (16) |
|             |             |               |            |
| S1—C1—C2—C3 | -0.2 (4)    | C5—C6—C10—O1  | -14.5 (5)  |
| C1—C2—C3—C4 | 1.0 (4)     | C7—C6—C10—O2  | -12.6 (5)  |
| C2—C3—C4—C5 | 174.0 (2)   | C5—C6—C10—O2  | 163.4 (3)  |
| C2—C3—C4—S1 | -1.3 (3)    | N2—C9—N1—C5   | -16.3 (4)  |
| C3—C4—C5—N1 | -81.0 (3)   | S2—C9—N1—C5   | 166.1 (2)  |
| S1—C4—C5—N1 | 93.8 (2)    | C4—C5—N1—C9   | -89.9 (3)  |

|              |            |                |            |
|--------------|------------|----------------|------------|
| C3—C4—C5—C6  | 157.1 (2)  | C6—C5—N1—C9    | 33.7 (3)   |
| S1—C4—C5—C6  | -28.1 (3)  | N1—C9—N2—C7    | -10.0 (4)  |
| N2—C7—C6—C10 | -177.2 (3) | S2—C9—N2—C7    | 167.7 (2)  |
| C8—C7—C6—C10 | 3.3 (5)    | C6—C7—N2—C9    | 14.3 (4)   |
| N2—C7—C6—C5  | 7.0 (4)    | C8—C7—N2—C9    | -166.2 (3) |
| C8—C7—C6—C5  | -172.5 (3) | O1—C10—O2—C11  | -3.4 (6)   |
| N1—C5—C6—C7  | -27.6 (4)  | C6—C10—O2—C11  | 178.7 (4)  |
| C4—C5—C6—C7  | 95.4 (3)   | C12—C11—O2—C10 | 176.3 (5)  |
| N1—C5—C6—C10 | 156.1 (2)  | C2—C1—S1—C4    | -0.5 (3)   |
| C4—C5—C6—C10 | -80.9 (3)  | C3—C4—S1—C1    | 1.1 (2)    |
| C7—C6—C10—O1 | 169.5 (3)  | C5—C4—S1—C1    | -174.5 (2) |

*Hydrogen-bond geometry (Å, °)*

Cg1 is the centroid of the S1/C1—C4 thiophene ring.

| <i>D</i> —H... <i>A</i>       | <i>D</i> —H | H... <i>A</i> | <i>D</i> ... <i>A</i> | <i>D</i> —H... <i>A</i> |
|-------------------------------|-------------|---------------|-----------------------|-------------------------|
| N1—H1N1...S2 <sup>i</sup>     | 0.86        | 2.63          | 3.408 (2)             | 151                     |
| N2—H2N2...O1 <sup>ii</sup>    | 0.86        | 2.15          | 2.984 (3)             | 162                     |
| C12—H12B...Cg1 <sup>iii</sup> | 0.96        | 2.81          | 3.664 (6)             | 149                     |

Symmetry codes: (i)  $-x+2, -y, -z+1$ ; (ii)  $x+1, y, z$ ; (iii)  $-x+1, -y+1, -z$ .