

Crystal structures of two hydrazinecarbothioamide derivatives: (*E*)-*N*-ethyl-2-[(4-oxo-4*H*-chromen-3-yl)methylidene]hydrazinecarbothioamide hemihydrate and (*E*)-2-[(4-chloro-2*H*-chromen-3-yl)methylidene]-*N*-phenylhydrazinecarbothioamide

Received 4 February 2015

Accepted 17 February 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Rajeswari Gangadharan,^a Jebiti Haribabu,^b Ramasamy Karvembu^b and K. Sethusankar^{c*}

^aDepartment of Physics, Ethiraj College for Women (Autonomous), Chennai 600 008, India, ^bDepartment of Chemistry, National Institute of Technology, Tiruchirappalli 620 015, India, and ^cDepartment of Physics, RKM Vivekananda College (Autonomous), Chennai 600 004, India. *Correspondence e-mail: ksethusankar@yahoo.co.in

Keywords: crystal structure; hydrazinecarbothioamide; thiourea derivatives; α -*N*-heterocycle; hydrogen bonding


CCDC references: 1016440; 1049914

Supporting information: this article has supporting information at journals.iucr.org/e


The title compounds, $C_{13}H_{13}N_3O_2S \cdot 0.5H_2O$, (I), and $C_{17}H_{14}ClN_3OS$, (II), are hydrazinecarbothioamide derivatives. Compound (I) crystallizes with two independent molecules (*A* and *B*) and a water molecule of crystallization in the asymmetric unit. The chromene moiety is essentially planar in molecules *A* and *B*, with maximum deviations of 0.028 (3) and 0.016 (3) Å, respectively, for the carbonyl C atoms. In (II), the pyran ring of the chromene moiety adopts a screw-boat conformation and the phenyl ring is inclined by 61.18 (9)° to its mean plane. In the crystal of (I), bifurcated N—H...O and C—H...O hydrogen bonds link the two independent molecules forming *A*–*B* dimers with two $R_2^2(6)$ ring motifs, and $R_2^2(10)$ and $R_2^2(14)$ ring motifs. In addition to these, the water molecule forms tetrafurcated hydrogen bonds which alternately generate $R_4^4(12)$ and $R_6^6(22)$ graph-set ring motifs. There are also π – π [inter-centroid distances = 3.5648 (14) and 3.6825 (15) Å] interactions present, leading to the formation of columns along the *c*-axis direction. In the crystal of (II), molecules are linked by pairs of N—H...S hydrogen bonds, forming inversion dimers with an $R_2^2(8)$ ring motif. The dimers are linked by C—H... π interactions, forming ribbons lying parallel to (210).

1. Chemical context

Thiosemicarbazones belong to a large group of thiourea derivatives which are derived from parent aldehydes and ketones. The biological activity of these compounds depends on the parent aldehyde and ketone (Beraldo & Gambino, 2004). Derivatives of hydrazinecarbothioamide constitute an important group of multidentate ligands with potential binding sites available for a wide variety of metal ions. The chemistry of thiosemicarbazone complexes has received much attention owing to their significant biological activities and medicinal properties. Presently, the areas in which thiosemicarbazones are receiving the most attention are based on their antitumour, antiprotozoal, antibacterial and antiviral activities (Finch *et al.*, 1999; Antholine *et al.*, 1977). α -*N*-heterocyclic thiosemicarbazones possess antitumour properties partially related to their ability to inhibit ribonucleoside reductase (RR), an iron-containing enzyme which is essential in DNA synthesis (Sartorelli *et al.*, 1970). The medicinal action of these thiosemicarbazones appears to be directly related to their ability to chelate the iron atom of the active site of RR or by destroying the tyrosinase radical present in a subunit of this protein (Thelander & Graslund, 1983). The structures of the



title compounds were determined in order to investigate the extent of electron delocalization, ligand conformations and to illustrate their biological implications.



2. Structural commentary

In compound (I) (Fig. 1), the chromene moieties of molecules *A* and *B* are essentially planar, with maximum deviations of 0.028 (3) and 0.016 (3) Å for atoms C7 and C7', respectively. However, in compound (II) (Fig. 2), the chromene moiety is not quite planar with a dihedral angle of 5.67 (12)° between the mean planes of the fused six-membered rings. In compound (II), the pyran ring of the chromene moiety adopts a screw-boat conformation [puckering amplitudes and smallest displacement parameters are $q = 0.314$ (2) Å, $\theta = 116.4$ (4)°, $\varphi = 147.5$ (5)° and $\Delta C_2 = 0.7$ (3)]. In compound (II), the dihedral angle between the chromene moiety and the phenyl ring is 61.18 (9)°. The deviation of the carbonyl O




Figure 1
The molecular structure of the two independent molecules (*A* and *B*) of compound (I), showing the atom labelling. Displacement ellipsoids are drawn at the 30% probability level.




Figure 2
The molecular structure of compound (II), showing the atom labelling. Displacement ellipsoids are drawn at the 30% probability level.

atoms (O2 and O2') from the mean plane of the pyran ring in compound (I) are 0.0838 (18) and 0.0386 (19) Å in molecules *A* and *B*, respectively, while the deviation of the Cl atom in compound (II) is 0.312 (1) Å.

The hydrazinecarbothioamide backbone is almost planar, with the maximum deviation being exhibited by atom N2 in both compounds; 0.025 (2) and 0.051 (2) Å, respectively, in molecules *A* and *B* of compound (I) and 0.072 (2) Å in compound (II).

Thiosemicarbazones exist in the thione form in the solid state and in solution they exist as an equilibrium mixture of thione and thiol forms (Kurup & Joseph, 2003). The fact that the compounds exist in the thione form is confirmed by the N–N, N–C and C=S bond lengths. The C–S bond lengths are 1.681 (2) and 1.673 (2) Å in molecules *A* and *B*, respectively, of compound (I), and 1.668 (2) Å in compound (II). These bond lengths are intermediate between normal S–C sp^2 single-bond and S=C sp^2 double-bond distances of *ca* 1.75 and 1.59 Å, respectively, indicating the presence of partial double-bond character (Kumbhar *et al.*, 1997). The N1–N2 bond lengths [varying between 1.367 (2) and 1.369 (2) Å] are very close to that reported for a similar substituted hydrazinecarbothioamide compound (Joseph *et al.*, 2004). The resonance involving the pyran ring would account for the shortening of the N–N distance through extensive delocalization. The C–N bond lengths [varying between 1.324 (3) and 1.361 (3) Å] are shorter than the normal C–N single bond length (*ca* 1.48 Å), also indicating some degree of delocalization in both compounds. The S1=C11–N2–N1 torsion angles are 177.31 (16) and 174.29 (16)°, respectively, in molecules *A* and *B* of compound (I) and –172.62 (17)° in compound (II). This indicates that the thionyl atom S1 is positioned *trans* to the azomethane nitrogen atom N1 in both compounds.

3. Supramolecular features

The water molecule of crystallization plays an important role in the hydrogen-bonding patterns of the three-dimensional network in compound (I). In the crystal packing of compound (I), bifurcated N–H⋯O and C–H⋯O hydrogen bonds involving carbonyl oxygens O2 and O2' in adjacent molecules,

Table 1
 Hydrogen-bond geometry (Å, °) for (I).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2'-H2'A\cdots O2^i$	0.86	2.09	2.900 (3)	158
$N2-H2A\cdots O2^i$	0.86	2.14	2.938 (2)	155
$N3-H3A\cdots O1W$	0.86	2.31	3.131 (3)	161
$O1W-H1WB\cdots S1^i$	0.85 (3)	2.47 (3)	3.322 (2)	178 (4)
$O1W-H1WA\cdots S1^{ii}$	0.87 (2)	2.52 (2)	3.370 (3)	167 (4)
$C9-H9\cdots O1W$	0.93	2.30	3.213 (4)	169
$C10-H10\cdots O2^i$	0.93	2.51	3.297 (3)	143
$C10'-H10'\cdots O2^i$	0.93	2.52	3.302 (3)	142

 Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$.



Table 2
 Hydrogen-bond geometry (Å, °) for (II).

Cg1 is the centroid of the C12–C17 phenyl ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H2A\cdots S1^i$	0.86	2.61	3.456 (2)	167
$C2-H2\cdots Cg1^{ii}$	0.93	2.86	3.697 (3)	151

 Symmetry codes: (i) $-x+2, -y+2, -z$; (ii) $-x+1, -y, -z$.

interconnect them to form $A-B$ dimers with two $R_2^1(6)$ ring motifs, and $R_2^2(10)$ and $R_2^2(14)$ ring motifs (Table 1 and Fig. 3). Similar bifurcated hydrogen bonds between molecule A and the water O atom form an $R_2^1(10)$ ring motif. In addition to these, the water molecule forms tetrafurcated hydrogen bonds which alternately generate $R_4^1(12)$ and $R_6^0(22)$ graph-set ring motifs. The supramolecular aggregation in the crystal of compound (I) is completed by the presence of slipped parallel $\pi-\pi$ interactions, forming columns along the c -axis direction. The most significant interactions are $Cg1\cdots Cg1^i = 3.5648$ (14) Å [inter-planar distance = 3.3154 (10) Å, slippage = 1.310 Å, where $Cg1$ is the centroid of the O1/C1/C6–C9 ring; symmetry code: (i) = $-x+1, -y+1, -z+1$] and $Cg5\cdots Cg5^{ii} = 3.6825$ (15) Å [inter-planar distance = 3.5441 (11) Å, slippage = 0.999 Å, where $Cg5$ is the centroid of the C1'–C6' ring; symmetry code: (ii) = $-x+2, -y+1, -z$].


Figure 3
 A partial view along the c axis of the crystal packing of compound (I), showing the $N-H\cdots O$, $C-H\cdots O$ and $OW-H\cdots S$ hydrogen bonds (dashed lines; see Table 1), which result in the formation of two $R_2^1(6)$ ring motifs and $R_2^2(10)$, $R_2^2(14)$, $R_4^1(12)$ and $R_6^0(22)$ ring motifs. H atoms not involved in hydrogen bonding have been omitted for clarity.

Figure 4
 A partial view along the b axis of the crystal packing of compound (II), showing the $N-H\cdots S$ hydrogen bonds (dashed lines; see Table 2), which result in the formation of inversion dimers with an $R_2^2(8)$ ring motif. H atoms not involved in hydrogen bonding have been omitted for clarity.

In the crystal of compound (II), molecules are linked by pairs of $N-H\cdots S$ hydrogen bonds, forming inversion dimers with an $R_2^2(8)$ ring motif (Table 2 and Fig. 4). The dimers are linked by $C-H\cdots\pi$ interactions (Table 2), forming ribbons lying parallel to plane (210).

4. Database survey

A search of the Cambridge Structural Database (Version 5.36; last update Nov. 2014; Groom & Allen, 2014) for similar structures gave 3 hits, one of which is a copper(II) complex, dibromo-(2-[[6-methyl-4-(oxo)-4*H*-chromen-3-yl]methylene]-*N*-phenylhydrazinecarbothioamide)copper (Ilies *et al.*, 2014). The other two include, *N*-methyl-2-[[4-oxo-4*H*-chromen-3-yl]methylene] hydrazinecarbothioamide (III) (Vimala *et al.*, 2014), which is the *N*-methyl derivative of compound (I), and (*E*)-2-[[4-chloro-2*H*-chromen-3-yl]methylene]-*N*-cyclohexylhydrazine carbothioamide (IV) (Gangadharan *et al.*, 2014), which is the *N*-cyclohexane derivative of compound (II). The bond distances and angles in compounds (I) and (III) are very similar, as are those in compounds (II) and (IV).

5. Synthesis and crystallization

Compound (I): 1.19 g (0.01 mol) of *N*-ethylhydrazinecarbothioamide was dissolved in 20 ml of hot ethanol and to this was added 1.74 g (0.01 mol) of 4-oxo-4*H*-chromene-3-carbaldehyde in 10 cm³ of ethanol over a period of 10 min with continuous stirring. The reaction mixture was refluxed for 2 h and allowed to cool whereby a shiny white compound began to separate; this was filtered and washed thoroughly with ethanol and then dried *in vacuo*. The compound was recrystallized from hot ethanol (yield: 96%), giving colourless block-like crystals.

Compound (II): 1.67 g (0.01 mol) of 4(*N*)-phenylthiosemicarbazide was dissolved in 20 ml of hot ethanol and to this was added 1.94 g (0.01 mol) of 4-chloro-2*H*-chromene-3-carbaldehyde in 10 ml of ethanol over a period of 10 min with

Table 3
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C ₁₃ H ₁₃ N ₃ O ₂ S·0.5H ₂ O	C ₁₇ H ₁₄ ClN ₃ OS
<i>M_r</i>	284.33	343.82
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$	Monoclinic, <i>P</i> 2 ₁ / <i>c</i>
Temperature (K)	296	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	8.2858 (2), 12.5422 (4), 14.3520 (5)	10.3176 (3), 5.7589 (2), 27.0364 (7)
α , β , γ (°)	114.379 (2), 95.751 (3), 94.200 (2)	90, 96.564 (2), 90
<i>V</i> (Å ³)	1340.81 (7)	1595.92 (8)
<i>Z</i>	4	4
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.25	0.38
Crystal size (mm)	0.35 × 0.30 × 0.25	0.30 × 0.25 × 0.20
Data collection		
Diffractometer	Bruker Kappa APEXII CCD	Bruker Kappa APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2008)	Multi-scan (<i>SADABS</i> ; Bruker, 2008)
<i>T</i> _{min} , <i>T</i> _{max}	0.917, 0.940	0.893, 0.927
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	19142, 5579, 2764	14667, 3902, 2089
<i>R</i> _{int}	0.046	0.050
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.631	0.668
Refinement		
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> [<i>F</i> ²], <i>S</i>	0.048, 0.137, 0.94	0.047, 0.124, 0.99
No. of reflections	5579	3902
No. of parameters	362	208
No. of restraints	2	0
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.26, -0.24	0.23, -0.20

Computer programs: *APEX2* and *SAINT* (Bruker, 2008), *SHELXS97* and *SHELXL97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia, 2012), *Mercury* (Macrae et al., 2008) and *PLATON* Spek, 2009).

continuous stirring. The reaction mixture was refluxed for 2 h and allowed to cool whereby a shiny yellow compound began to separate. It was filtered and washed thoroughly with ethanol and then dried *in vacuo*. The compound was recrystallized from hot ethanol (yield: 89%), giving colourless block-like crystals.

6. Refinement

Crystal data, data collection and structure refinement details for compounds (I) and (II) are summarized in Table 3. For compound (I), the positions of the water H atoms were located from difference electron density maps and freely refined. In compounds (I) and (II), the NH H atoms were included in calculated positions and treated as riding atoms: N–H = 0.86 Å with *U*_{iso}(H) = 1.2*U*_{eq}(N). The C-bound H atoms in both molecules were included in calculated positions and treated as riding atoms: C–H = 0.93–0.97 Å with *U*_{iso}(H) = 1.5*U*_{eq}(C) for methyl H atoms and = 1.2*U*_{eq}(C) for other H atoms.

Acknowledgements

The authors thank Professor D. Velmurugan and Mr T. Srinivasan, CAS in Crystallography and Biophysics, University of Madras, Chennai, India, for the X-ray intensity data collection.

References

- Antholine, W., Knight, J., Whelan, H. & Petering, D. H. (1977). *Mol. Pharmacol.* **13**, 89–98.
- Beraldo, H. & Gambino, D. (2004). *Mini. Rev. Med. Chem.* **4**, 31–39.
- Bruker (2008). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Finch, R. A., Liu, M. C., Cory, A. H., Cory, J. G. & Sartorelli, A. C. (1999). *Adv. Enzyme Regul.* **39**, 3–12.
- Gangadharan, R., Haribabu, J., Karvembu, R. & Sethusankar, K. (2014). *Acta Cryst. E* **70**, o1039–o1040.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Ilies, D.-C., Pahontu, E., Shova, S., Georgescu, R., Stanica, N., Olar, R., Gulea, A. & Rosu, T. (2014). *Polyhedron*, **81**, 123–131.
- Joseph, M., Suni, V., Nayar, C. R., Kurup, M. R. P. & Fun, H.-K. (2004). *J. Mol. Struct.* **705**, 63–70.
- Kumbhar, A., Sonawane, P., Padhye, S., West, D. X. & Butcher, R. J. (1997). *J. Chem. Crystallogr.* **27**, 533–539.
- Kurup, M. R. P. & Joseph, M. (2003). *Synth. React. Inorg. Met. Org. Chem.* **33**, 1275–1287.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Sartorelli, A. C., Moore, E. C., Zedeck, M. S. & Agrawal, K. C. (1970). *Biochemistry*, **9**, 4492–4498.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Thelander, L. & Gräslund, A. (1983). *J. Biol. Chem.* **258**, 4063–4066.
- Vimala, G., Govindaraj, J., Haribabu, J., Karvembu, R. & Subbiah-Pandi, A. (2014). *Acta Cryst. E* **70**, o1151.

supporting information

Acta Cryst. (2015). E71, 305-308 [doi:10.1107/S2056989015003369]

Crystal structures of two hydrazinecarbothioamide derivatives: (*E*)-*N*-ethyl-2-[(4-oxo-4*H*-chromen-3-yl)methylidene]hydrazinecarbothioamide hemihydrate and (*E*)-2-[(4-chloro-2*H*-chromen-3-yl)methylidene]-*N*-phenyl-hydrazinecarbothioamide

Rajeswari Gangadharan, Jebiti Haribabu, Ramasamy Karvembu and K. Sethusankar

Computing details

For both compounds, data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINTE* (Bruker, 2008); data reduction: *SAINTE* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *Mercury* (Macrae *et al.*, 2008). Software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009) for (I); *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009) for (II).

(I) (*E*)-*N*-Ethyl-2-[(4-oxo-4*H*-chromen-3-yl)methylidene]hydrazinecarbothioamide hemihydrate

Crystal data

$C_{13}H_{13}N_3O_2S \cdot 0.5H_2O$

$M_r = 284.33$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.2858$ (2) Å

$b = 12.5422$ (4) Å

$c = 14.3520$ (5) Å

$\alpha = 114.379$ (2)°

$\beta = 95.751$ (3)°

$\gamma = 94.200$ (2)°

$V = 1340.81$ (7) Å³

$Z = 4$

$F(000) = 596$

$D_x = 1.409$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5579 reflections

$\theta = 1.6$ – 26.6 °

$\mu = 0.25$ mm⁻¹

$T = 296$ K

Block, colourless

$0.35 \times 0.30 \times 0.25$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and ϕ scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.917$, $T_{\max} = 0.940$

19142 measured reflections

5579 independent reflections

2764 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 26.6$ °, $\theta_{\min} = 1.6$ °

$h = -10 \rightarrow 10$

$k = -14 \rightarrow 15$

$l = -18 \rightarrow 18$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.137$
 $S = 0.94$
 5579 reflections
 362 parameters
 2 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0658P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.26 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.24 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1'	0.7753 (3)	0.4656 (2)	-0.08664 (19)	0.0444 (6)
C1	0.8004 (3)	0.5298 (2)	0.46364 (19)	0.0425 (6)
C2'	0.8576 (3)	0.5059 (2)	-0.1473 (2)	0.0557 (7)
H2'	0.8603	0.4568	-0.2163	0.067*
C2	0.8858 (3)	0.5665 (2)	0.4025 (2)	0.0557 (7)
H2	0.8976	0.5138	0.3360	0.067*
C3	0.9530 (3)	0.6821 (2)	0.4413 (2)	0.0581 (7)
H3	1.0107	0.7083	0.4008	0.070*
C3'	0.9351 (3)	0.6189 (2)	-0.1045 (2)	0.0593 (8)
H3'	0.9903	0.6469	-0.1448	0.071*
C4'	0.9321 (3)	0.6925 (2)	-0.0009 (2)	0.0572 (7)
H4'	0.9859	0.7690	0.0278	0.069*
C4	0.9357 (3)	0.7602 (2)	0.5402 (2)	0.0568 (7)
H4	0.9835	0.8382	0.5664	0.068*
C5	0.8487 (3)	0.7233 (2)	0.6000 (2)	0.0487 (6)
H5	0.8357	0.7768	0.6660	0.058*
C5'	0.8498 (3)	0.6520 (2)	0.0583 (2)	0.0503 (7)
H5'	0.8475	0.7017	0.1273	0.060*
C6	0.7792 (3)	0.6059 (2)	0.56270 (18)	0.0404 (6)
C6'	0.7691 (3)	0.5370 (2)	0.01658 (19)	0.0408 (6)
C7	0.6827 (3)	0.5632 (2)	0.62307 (19)	0.0420 (6)
C7'	0.6784 (3)	0.4911 (2)	0.07702 (19)	0.0429 (6)
C8'	0.6037 (3)	0.3694 (2)	0.02285 (18)	0.0409 (6)
C8	0.6228 (3)	0.43815 (19)	0.57379 (18)	0.0394 (6)

C9	0.6516 (3)	0.3722 (2)	0.47741 (19)	0.0488 (7)
H9	0.6102	0.2926	0.4477	0.059*
C9'	0.6189 (3)	0.3082 (2)	-0.0777 (2)	0.0535 (7)
H9'	0.5693	0.2305	-0.1106	0.064*
C10	0.5298 (3)	0.3872 (2)	0.62971 (18)	0.0430 (6)
H10	0.5125	0.4350	0.6965	0.052*
C10'	0.5137 (3)	0.3172 (2)	0.07880 (19)	0.0456 (6)
H10'	0.5090	0.3620	0.1485	0.055*
C11	0.3128 (3)	0.1280 (2)	0.61075 (18)	0.0412 (6)
C11'	0.2875 (3)	0.0607 (2)	0.05910 (19)	0.0420 (6)
C12	0.2363 (3)	-0.0610 (2)	0.45867 (19)	0.0538 (7)
H12A	0.1249	-0.0630	0.4739	0.065*
H12B	0.2923	-0.1109	0.4842	0.065*
C12'	0.1908 (3)	-0.1203 (2)	-0.09642 (19)	0.0524 (7)
H12C	0.0750	-0.1186	-0.0915	0.063*
H12D	0.2316	-0.1696	-0.0645	0.063*
C13'	0.2150 (4)	-0.1715 (2)	-0.2081 (2)	0.0732 (9)
H13D	0.1711	-0.1241	-0.2402	0.110*
H13E	0.1596	-0.2505	-0.2426	0.110*
H13F	0.3296	-0.1725	-0.2128	0.110*
C13	0.2336 (3)	-0.1086 (2)	0.3439 (2)	0.0631 (8)
H13A	0.1773	-0.0598	0.3182	0.095*
H13B	0.1782	-0.1877	0.3115	0.095*
H13C	0.3437	-0.1086	0.3285	0.095*
N1	0.4719 (2)	0.27888 (16)	0.58919 (15)	0.0431 (5)
N1'	0.4412 (2)	0.21203 (17)	0.03521 (15)	0.0447 (5)
N2'	0.3664 (2)	0.17184 (16)	0.09748 (16)	0.0499 (5)
H2'A	0.3695	0.2178	0.1618	0.060*
N2	0.3884 (2)	0.24022 (16)	0.64915 (15)	0.0454 (5)
H2A	0.3837	0.2878	0.7121	0.055*
N3'	0.2766 (2)	-0.00132 (17)	-0.04207 (16)	0.0511 (6)
H3'A	0.3224	0.0303	-0.0773	0.061*
N3	0.3184 (2)	0.06031 (16)	0.51214 (15)	0.0475 (5)
H3A	0.3728	0.0889	0.4779	0.057*
O1	0.7355 (2)	0.41240 (14)	0.42089 (13)	0.0534 (5)
O1'	0.6999 (2)	0.35106 (14)	-0.13347 (13)	0.0575 (5)
O2	0.6529 (2)	0.62856 (14)	0.70913 (13)	0.0603 (5)
O2'	0.6649 (2)	0.55142 (14)	0.16840 (14)	0.0644 (5)
O1W	0.4956 (3)	0.10193 (19)	0.34506 (17)	0.0689 (6)
S1'	0.21192 (9)	0.00898 (6)	0.13833 (5)	0.0591 (2)
S1	0.21930 (9)	0.08415 (6)	0.68995 (5)	0.0587 (2)
H1WA	0.569 (4)	0.054 (3)	0.326 (4)	0.18 (2)*
H1WB	0.422 (4)	0.080 (4)	0.293 (2)	0.16 (2)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1'	0.0496 (15)	0.0441 (15)	0.0419 (16)	0.0038 (12)	0.0133 (13)	0.0194 (13)

C1	0.0458 (15)	0.0384 (14)	0.0450 (16)	0.0025 (12)	0.0109 (13)	0.0186 (13)
C2'	0.0640 (17)	0.0635 (18)	0.0423 (16)	0.0010 (15)	0.0162 (14)	0.0244 (15)
C2	0.0649 (18)	0.0576 (18)	0.0497 (17)	0.0082 (15)	0.0259 (15)	0.0238 (15)
C3	0.0645 (18)	0.0609 (18)	0.0604 (19)	0.0022 (15)	0.0229 (15)	0.0348 (16)
C3'	0.0595 (18)	0.0645 (19)	0.063 (2)	-0.0021 (15)	0.0179 (15)	0.0363 (17)
C4'	0.0574 (17)	0.0525 (17)	0.064 (2)	-0.0023 (14)	0.0111 (15)	0.0284 (16)
C4	0.0595 (17)	0.0492 (16)	0.064 (2)	-0.0055 (14)	0.0125 (15)	0.0278 (16)
C5	0.0547 (16)	0.0430 (15)	0.0467 (16)	-0.0020 (13)	0.0092 (13)	0.0181 (13)
C5'	0.0566 (16)	0.0463 (15)	0.0475 (16)	0.0018 (13)	0.0113 (13)	0.0193 (13)
C6	0.0421 (14)	0.0411 (14)	0.0404 (15)	0.0030 (11)	0.0084 (12)	0.0194 (12)
C6'	0.0441 (14)	0.0392 (14)	0.0422 (15)	0.0051 (11)	0.0096 (12)	0.0196 (12)
C7	0.0483 (15)	0.0415 (14)	0.0366 (15)	0.0032 (12)	0.0064 (12)	0.0172 (13)
C7'	0.0481 (15)	0.0405 (14)	0.0411 (16)	0.0068 (12)	0.0129 (13)	0.0167 (13)
C8'	0.0460 (14)	0.0385 (14)	0.0389 (15)	0.0040 (11)	0.0113 (12)	0.0162 (12)
C8	0.0462 (14)	0.0349 (13)	0.0372 (15)	0.0031 (11)	0.0099 (12)	0.0150 (12)
C9	0.0612 (17)	0.0384 (14)	0.0463 (16)	-0.0024 (13)	0.0167 (14)	0.0170 (13)
C9'	0.0697 (18)	0.0418 (15)	0.0472 (17)	-0.0033 (13)	0.0186 (14)	0.0163 (14)
C10	0.0531 (15)	0.0389 (14)	0.0365 (14)	0.0013 (12)	0.0115 (12)	0.0151 (12)
C10'	0.0574 (16)	0.0398 (14)	0.0401 (15)	0.0045 (13)	0.0167 (13)	0.0156 (13)
C11	0.0483 (15)	0.0353 (14)	0.0417 (15)	0.0055 (12)	0.0119 (12)	0.0169 (12)
C11'	0.0468 (15)	0.0358 (14)	0.0432 (16)	0.0027 (12)	0.0100 (12)	0.0160 (12)
C12	0.0696 (18)	0.0405 (14)	0.0489 (17)	-0.0031 (13)	0.0164 (14)	0.0164 (13)
C12'	0.0598 (17)	0.0452 (15)	0.0488 (17)	-0.0027 (13)	0.0110 (14)	0.0174 (13)
C13'	0.094 (2)	0.0668 (19)	0.0442 (18)	-0.0098 (17)	0.0102 (17)	0.0126 (15)
C13	0.0741 (19)	0.0524 (17)	0.0506 (18)	-0.0046 (15)	0.0149 (15)	0.0105 (14)
N1	0.0539 (13)	0.0379 (12)	0.0395 (12)	-0.0002 (10)	0.0126 (10)	0.0182 (10)
N1'	0.0547 (13)	0.0400 (12)	0.0417 (12)	0.0007 (10)	0.0154 (10)	0.0184 (10)
N2'	0.0680 (14)	0.0413 (12)	0.0389 (12)	-0.0031 (11)	0.0175 (11)	0.0150 (10)
N2	0.0614 (13)	0.0371 (11)	0.0368 (12)	-0.0003 (10)	0.0166 (10)	0.0136 (10)
N3'	0.0663 (14)	0.0437 (12)	0.0412 (13)	-0.0063 (11)	0.0148 (11)	0.0164 (11)
N3	0.0602 (13)	0.0403 (12)	0.0422 (13)	-0.0013 (10)	0.0180 (11)	0.0166 (10)
O1	0.0716 (12)	0.0426 (10)	0.0428 (11)	-0.0008 (9)	0.0246 (9)	0.0125 (9)
O1'	0.0788 (13)	0.0477 (11)	0.0407 (11)	-0.0064 (9)	0.0234 (10)	0.0127 (9)
O2	0.0948 (14)	0.0418 (10)	0.0390 (11)	-0.0025 (10)	0.0261 (10)	0.0101 (9)
O2'	0.0966 (14)	0.0463 (11)	0.0427 (12)	-0.0063 (10)	0.0302 (11)	0.0094 (9)
O1W	0.0823 (16)	0.0636 (14)	0.0546 (14)	-0.0030 (13)	0.0150 (13)	0.0196 (11)
S1'	0.0820 (5)	0.0521 (4)	0.0441 (4)	-0.0073 (4)	0.0147 (4)	0.0227 (4)
S1	0.0800 (5)	0.0487 (4)	0.0507 (4)	-0.0004 (4)	0.0288 (4)	0.0214 (3)

Geometric parameters (Å, °)

C1'—O1'	1.376 (3)	C9'—O1'	1.338 (3)
C1'—C2'	1.382 (3)	C9'—H9'	0.9300
C1'—C6'	1.388 (3)	C10—N1	1.269 (3)
C1—C2	1.376 (3)	C10—H10	0.9300
C1—O1	1.380 (3)	C10'—N1'	1.273 (3)
C1—C6	1.385 (3)	C10'—H10'	0.9300
C2'—C3'	1.366 (3)	C11—N3	1.324 (3)

C2'—H2'	0.9300	C11—N2	1.356 (3)
C2—C3	1.370 (3)	C11—S1	1.681 (2)
C2—H2	0.9300	C11'—N3'	1.324 (3)
C3—C4	1.382 (4)	C11'—N2'	1.354 (3)
C3—H3	0.9300	C11'—S1'	1.673 (2)
C3'—C4'	1.392 (4)	C12—N3	1.465 (3)
C3'—H3'	0.9300	C12—C13	1.500 (3)
C4'—C5'	1.367 (3)	C12—H12A	0.9700
C4'—H4'	0.9300	C12—H12B	0.9700
C4—C5	1.369 (3)	C12'—N3'	1.454 (3)
C4—H4	0.9300	C12'—C13'	1.501 (3)
C5—C6	1.396 (3)	C12'—H12C	0.9700
C5—H5	0.9300	C12'—H12D	0.9700
C5'—C6'	1.398 (3)	C13'—H13D	0.9600
C5'—H5'	0.9300	C13'—H13E	0.9600
C6—C7	1.461 (3)	C13'—H13F	0.9600
C6'—C7'	1.458 (3)	C13—H13A	0.9600
C7—O2	1.230 (3)	C13—H13B	0.9600
C7—C8	1.452 (3)	C13—H13C	0.9600
C7'—O2'	1.236 (3)	N1—N2	1.367 (2)
C7'—C8'	1.451 (3)	N1'—N2'	1.369 (2)
C8'—C9'	1.350 (3)	N2'—H2'A	0.8600
C8'—C10'	1.453 (3)	N2—H2A	0.8600
C8—C9	1.344 (3)	N3'—H3'A	0.8600
C8—C10	1.457 (3)	N3—H3A	0.8600
C9—O1	1.339 (2)	O1W—H1WA	0.866 (19)
C9—H9	0.9300	O1W—H1WB	0.847 (19)
O1'—C1'—C2'	116.8 (2)	N1—C10—C8	121.4 (2)
O1'—C1'—C6'	121.5 (2)	N1—C10—H10	119.3
C2'—C1'—C6'	121.7 (2)	C8—C10—H10	119.3
C2—C1—O1	116.5 (2)	N1'—C10'—C8'	121.9 (2)
C2—C1—C6	122.3 (2)	N1'—C10'—H10'	119.0
O1—C1—C6	121.3 (2)	C8'—C10'—H10'	119.0
C3'—C2'—C1'	119.2 (2)	N3—C11—N2	116.8 (2)
C3'—C2'—H2'	120.4	N3—C11—S1	124.54 (18)
C1'—C2'—H2'	120.4	N2—C11—S1	118.62 (18)
C3—C2—C1	118.8 (2)	N3'—C11'—N2'	116.0 (2)
C3—C2—H2	120.6	N3'—C11'—S1'	123.86 (18)
C1—C2—H2	120.6	N2'—C11'—S1'	120.18 (19)
C2—C3—C4	120.5 (2)	N3—C12—C13	111.8 (2)
C2—C3—H3	119.8	N3—C12—H12A	109.2
C4—C3—H3	119.8	C13—C12—H12A	109.2
C2'—C3'—C4'	120.6 (2)	N3—C12—H12B	109.2
C2'—C3'—H3'	119.7	C13—C12—H12B	109.2
C4'—C3'—H3'	119.7	H12A—C12—H12B	107.9
C5'—C4'—C3'	119.9 (2)	N3'—C12'—C13'	110.4 (2)
C5'—C4'—H4'	120.1	N3'—C12'—H12C	109.6

C3'—C4'—H4'	120.1	C13'—C12'—H12C	109.6
C5—C4—C3	120.3 (2)	N3'—C12'—H12D	109.6
C5—C4—H4	119.8	C13'—C12'—H12D	109.6
C3—C4—H4	119.8	H12C—C12'—H12D	108.1
C4—C5—C6	120.5 (2)	C12'—C13'—H13D	109.5
C4—C5—H5	119.7	C12'—C13'—H13E	109.5
C6—C5—H5	119.7	H13D—C13'—H13E	109.5
C4'—C5'—C6'	120.8 (2)	C12'—C13'—H13F	109.5
C4'—C5'—H5'	119.6	H13D—C13'—H13F	109.5
C6'—C5'—H5'	119.6	H13E—C13'—H13F	109.5
C1—C6—C5	117.6 (2)	C12—C13—H13A	109.5
C1—C6—C7	120.0 (2)	C12—C13—H13B	109.5
C5—C6—C7	122.3 (2)	H13A—C13—H13B	109.5
C1'—C6'—C5'	117.9 (2)	C12—C13—H13C	109.5
C1'—C6'—C7'	119.7 (2)	H13A—C13—H13C	109.5
C5'—C6'—C7'	122.4 (2)	H13B—C13—H13C	109.5
O2—C7—C8	122.2 (2)	C10—N1—N2	116.40 (19)
O2—C7—C6	122.6 (2)	C10'—N1'—N2'	116.1 (2)
C8—C7—C6	115.2 (2)	C11'—N2'—N1'	120.9 (2)
O2'—C7'—C8'	121.7 (2)	C11'—N2'—H2'A	119.6
O2'—C7'—C6'	122.7 (2)	N1'—N2'—H2'A	119.6
C8'—C7'—C6'	115.7 (2)	C11—N2—N1	120.95 (19)
C9'—C8'—C7'	119.3 (2)	C11—N2—H2A	119.5
C9'—C8'—C10'	122.0 (2)	N1—N2—H2A	119.5
C7'—C8'—C10'	118.7 (2)	C11'—N3'—C12'	123.2 (2)
C9—C8—C7	119.7 (2)	C11'—N3'—H3'A	118.4
C9—C8—C10	121.4 (2)	C12'—N3'—H3'A	118.4
C7—C8—C10	119.0 (2)	C11—N3—C12	123.1 (2)
O1—C9—C8	125.0 (2)	C11—N3—H3A	118.5
O1—C9—H9	117.5	C12—N3—H3A	118.5
C8—C9—H9	117.5	C9—O1—C1	118.76 (18)
O1'—C9'—C8'	124.9 (2)	C9'—O1'—C1'	118.85 (19)
O1'—C9'—H9'	117.5	H1WA—O1W—H1WB	106 (4)
C8'—C9'—H9'	117.5		
O1'—C1'—C2'—C3'	-179.8 (2)	C6'—C7'—C8'—C10'	179.0 (2)
C6'—C1'—C2'—C3'	0.1 (4)	O2—C7—C8—C9	176.4 (2)
O1—C1—C2—C3	-179.9 (2)	C6—C7—C8—C9	-2.9 (3)
C6—C1—C2—C3	0.5 (4)	O2—C7—C8—C10	-2.5 (4)
C1—C2—C3—C4	0.2 (4)	C6—C7—C8—C10	178.11 (19)
C1'—C2'—C3'—C4'	0.3 (4)	C7—C8—C9—O1	0.9 (4)
C2'—C3'—C4'—C5'	-0.6 (4)	C10—C8—C9—O1	179.8 (2)
C2—C3—C4—C5	-1.1 (4)	C7'—C8'—C9'—O1'	0.3 (4)
C3—C4—C5—C6	1.3 (4)	C10'—C8'—C9'—O1'	180.0 (2)
C3'—C4'—C5'—C6'	0.4 (4)	C9—C8—C10—N1	-0.4 (4)
C2—C1—C6—C5	-0.3 (4)	C7—C8—C10—N1	178.5 (2)
O1—C1—C6—C5	-179.9 (2)	C9'—C8'—C10'—N1'	-1.7 (4)
C2—C1—C6—C7	177.9 (2)	C7'—C8'—C10'—N1'	177.9 (2)

O1—C1—C6—C7	-1.7 (4)	C8—C10—N1—N2	179.22 (18)
C4—C5—C6—C1	-0.6 (4)	C8'—C10'—N1'—N2'	177.44 (19)
C4—C5—C6—C7	-178.8 (2)	N3'—C11'—N2'—N1'	-5.3 (3)
O1'—C1'—C6'—C5'	179.6 (2)	S1'—C11'—N2'—N1'	174.29 (16)
C2'—C1'—C6'—C5'	-0.2 (4)	C10'—N1'—N2'—C11'	-179.2 (2)
O1'—C1'—C6'—C7'	-1.1 (4)	N3—C11—N2—N1	-2.7 (3)
C2'—C1'—C6'—C7'	179.0 (2)	S1—C11—N2—N1	177.31 (16)
C4'—C5'—C6'—C1'	0.0 (4)	C10—N1—N2—C11	175.7 (2)
C4'—C5'—C6'—C7'	-179.3 (2)	N2'—C11'—N3'—C12'	-177.6 (2)
C1—C6—C7—O2	-176.0 (2)	S1'—C11'—N3'—C12'	2.8 (3)
C5—C6—C7—O2	2.1 (4)	C13'—C12'—N3'—C11'	-174.1 (2)
C1—C6—C7—C8	3.3 (3)	N2—C11—N3—C12	-177.1 (2)
C5—C6—C7—C8	-178.5 (2)	S1—C11—N3—C12	2.8 (3)
C1'—C6'—C7'—O2'	-178.0 (2)	C13—C12—N3—C11	168.4 (2)
C5'—C6'—C7'—O2'	1.2 (4)	C8—C9—O1—C1	1.0 (4)
C1'—C6'—C7'—C8'	1.7 (3)	C2—C1—O1—C9	179.8 (2)
C5'—C6'—C7'—C8'	-179.1 (2)	C6—C1—O1—C9	-0.5 (3)
O2'—C7'—C8'—C9'	178.4 (2)	C8'—C9'—O1'—C1'	0.4 (4)
C6'—C7'—C8'—C9'	-1.3 (3)	C2'—C1'—O1'—C9'	179.9 (2)
O2'—C7'—C8'—C10'	-1.3 (3)	C6'—C1'—O1'—C9'	0.1 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2'—H2'A \cdots O2 ⁱ	0.86	2.09	2.900 (3)	158
N2—H2A \cdots O2 ⁱⁱ	0.86	2.14	2.938 (2)	155
N3—H3A \cdots O1W	0.86	2.31	3.131 (3)	161
O1W—H1WB \cdots S1'	0.85 (3)	2.47 (3)	3.322 (2)	178 (4)
O1W—H1WA \cdots S1 ⁱⁱ	0.87 (2)	2.52 (2)	3.370 (3)	167 (4)
C9—H9 \cdots O1W	0.93	2.30	3.213 (4)	169
C10—H10 \cdots O2 ⁱⁱ	0.93	2.51	3.297 (3)	143
C10'—H10' \cdots O2 ⁱ	0.93	2.52	3.302 (3)	142

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$.

(II) (E)-2-[(4-Chloro-2H-chromen-3-yl)methylidene]-N-phenylhydrazinecarbothioamide

Crystal data

 $C_{17}H_{14}ClN_3OS$ $M_r = 343.82$ Monoclinic, $P2_1/c$ Hall symbol: $-P\ 2ybc$ $a = 10.3176$ (3) \AA $b = 5.7589$ (2) \AA $c = 27.0364$ (7) \AA $\beta = 96.564$ (2) $^\circ$ $V = 1595.92$ (8) \AA^3 $Z = 4$ $F(000) = 712$ $D_x = 1.431$ Mg m^{-3} Mo $K\alpha$ radiation, $\lambda = 0.71073$ \AA

Cell parameters from 3902 reflections

 $\theta = 1.5\text{--}28.4^\circ$ $\mu = 0.38$ mm^{-1} $T = 296$ K

Block, colourless

 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2008)
 $T_{\min} = 0.893$, $T_{\max} = 0.927$

14667 measured reflections
3902 independent reflections
2089 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$
 $\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -13 \rightarrow 13$
 $k = -7 \rightarrow 7$
 $l = -35 \rightarrow 35$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.124$
 $S = 0.99$
3902 reflections
208 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.053P)^2 + 0.0466P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4918 (3)	-0.2996 (5)	-0.10347 (12)	0.0665 (8)
H1	0.4591	-0.3879	-0.0789	0.080*
C2	0.4637 (3)	-0.3582 (5)	-0.15313 (13)	0.0749 (9)
H2	0.4120	-0.4872	-0.1621	0.090*
C3	0.5116 (3)	-0.2267 (6)	-0.18911 (12)	0.0759 (9)
H3	0.4921	-0.2669	-0.2224	0.091*
C4	0.5882 (3)	-0.0365 (5)	-0.17654 (10)	0.0622 (7)
H4	0.6193	0.0518	-0.2015	0.075*
C5	0.6198 (2)	0.0260 (4)	-0.12710 (8)	0.0457 (6)
C6	0.5690 (2)	-0.1087 (4)	-0.09079 (10)	0.0531 (6)
C7	0.6397 (3)	0.1634 (4)	-0.02714 (9)	0.0563 (7)
H7A	0.6881	0.1612	0.0058	0.068*
H7B	0.5608	0.2529	-0.0251	0.068*
C8	0.7205 (2)	0.2839 (4)	-0.06223 (8)	0.0456 (6)
C9	0.7056 (2)	0.2157 (4)	-0.11002 (8)	0.0446 (6)
C10	0.7994 (2)	0.4764 (4)	-0.04404 (9)	0.0490 (6)

H10	0.8449	0.5619	-0.0656	0.059*
C11	0.8930 (2)	0.7961 (4)	0.06460 (8)	0.0458 (6)
C12	0.8439 (2)	0.6739 (4)	0.14825 (8)	0.0430 (6)
C13	0.8894 (2)	0.4948 (4)	0.17950 (9)	0.0497 (6)
H13	0.9269	0.3642	0.1667	0.060*
C14	0.8788 (2)	0.5110 (5)	0.22994 (9)	0.0538 (7)
H14	0.9088	0.3902	0.2510	0.065*
C15	0.8245 (2)	0.7033 (5)	0.24904 (9)	0.0563 (7)
H15	0.8189	0.7146	0.2831	0.068*
C16	0.7784 (2)	0.8787 (5)	0.21777 (9)	0.0580 (7)
H16	0.7407	1.0086	0.2307	0.070*
C17	0.7870 (2)	0.8659 (4)	0.16725 (9)	0.0520 (6)
H17	0.7547	0.9856	0.1463	0.062*
N1	0.80705 (19)	0.5309 (3)	0.00252 (7)	0.0496 (5)
N2	0.87785 (19)	0.7257 (4)	0.01625 (7)	0.0533 (5)
H2A	0.9130	0.8041	-0.0058	0.064*
N3	0.84862 (19)	0.6451 (3)	0.09615 (7)	0.0520 (5)
H3A	0.8196	0.5154	0.0836	0.062*
O1	0.60439 (18)	-0.0669 (3)	-0.04098 (6)	0.0662 (5)
Cl1	0.79000 (7)	0.35034 (13)	-0.15374 (2)	0.0649 (2)
S1	0.96413 (7)	1.05093 (12)	0.07927 (2)	0.0565 (2)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0604 (16)	0.0548 (19)	0.086 (2)	-0.0045 (14)	0.0171 (15)	-0.0053 (17)
C2	0.0523 (16)	0.070 (2)	0.101 (3)	-0.0045 (15)	0.0005 (16)	-0.030 (2)
C3	0.0651 (18)	0.092 (3)	0.068 (2)	0.0024 (18)	-0.0059 (15)	-0.0280 (19)
C4	0.0606 (16)	0.074 (2)	0.0509 (16)	0.0048 (15)	0.0005 (13)	-0.0125 (15)
C5	0.0467 (13)	0.0493 (15)	0.0410 (13)	0.0075 (11)	0.0046 (10)	-0.0025 (12)
C6	0.0534 (14)	0.0502 (17)	0.0564 (16)	0.0037 (12)	0.0089 (12)	-0.0055 (13)
C7	0.0773 (17)	0.0502 (17)	0.0437 (14)	-0.0077 (14)	0.0160 (12)	0.0006 (13)
C8	0.0527 (13)	0.0474 (15)	0.0375 (13)	0.0009 (12)	0.0093 (10)	0.0022 (12)
C9	0.0500 (13)	0.0473 (15)	0.0374 (12)	0.0027 (11)	0.0096 (10)	0.0037 (11)
C10	0.0584 (15)	0.0506 (16)	0.0387 (13)	-0.0022 (12)	0.0088 (11)	0.0028 (12)
C11	0.0528 (14)	0.0441 (15)	0.0416 (13)	0.0011 (11)	0.0103 (11)	0.0004 (12)
C12	0.0496 (13)	0.0409 (15)	0.0397 (13)	-0.0090 (11)	0.0099 (10)	-0.0015 (11)
C13	0.0582 (15)	0.0412 (15)	0.0522 (15)	0.0008 (12)	0.0167 (12)	-0.0038 (12)
C14	0.0631 (16)	0.0518 (17)	0.0479 (15)	-0.0023 (13)	0.0123 (12)	0.0063 (13)
C15	0.0646 (16)	0.0636 (19)	0.0428 (14)	-0.0140 (14)	0.0156 (12)	-0.0079 (14)
C16	0.0673 (16)	0.0512 (17)	0.0589 (17)	-0.0020 (13)	0.0211 (13)	-0.0143 (14)
C17	0.0623 (16)	0.0413 (15)	0.0539 (15)	0.0029 (12)	0.0128 (12)	0.0021 (13)
N1	0.0624 (13)	0.0435 (13)	0.0433 (12)	-0.0043 (10)	0.0083 (9)	-0.0006 (10)
N2	0.0713 (13)	0.0527 (14)	0.0368 (11)	-0.0131 (11)	0.0101 (9)	0.0021 (10)
N3	0.0789 (14)	0.0406 (12)	0.0386 (11)	-0.0136 (11)	0.0148 (10)	-0.0063 (10)
O1	0.0928 (13)	0.0550 (13)	0.0525 (11)	-0.0160 (10)	0.0162 (10)	0.0034 (10)
Cl1	0.0802 (5)	0.0762 (5)	0.0411 (4)	-0.0098 (4)	0.0189 (3)	0.0059 (3)
S1	0.0796 (5)	0.0433 (4)	0.0488 (4)	-0.0101 (3)	0.0162 (3)	-0.0023 (3)

Geometric parameters (Å, °)

C1—C6	1.378 (3)	C10—H10	0.9300
C1—C2	1.383 (4)	C11—N3	1.335 (3)
C1—H1	0.9300	C11—N2	1.361 (3)
C2—C3	1.369 (4)	C11—S1	1.668 (2)
C2—H2	0.9300	C12—C17	1.378 (3)
C3—C4	1.371 (4)	C12—C13	1.381 (3)
C3—H3	0.9300	C12—N3	1.425 (3)
C4—C5	1.387 (3)	C13—C14	1.384 (3)
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.399 (3)	C14—C15	1.369 (3)
C5—C9	1.448 (3)	C14—H14	0.9300
C6—O1	1.375 (3)	C15—C16	1.367 (3)
C7—O1	1.414 (3)	C15—H15	0.9300
C7—C8	1.503 (3)	C16—C17	1.381 (3)
C7—H7A	0.9700	C16—H16	0.9300
C7—H7B	0.9700	C17—H17	0.9300
C8—C9	1.343 (3)	N1—N2	1.367 (3)
C8—C10	1.429 (3)	N2—H2A	0.8600
C9—C11	1.730 (2)	N3—H3A	0.8600
C10—N1	1.291 (3)		
C6—C1—C2	119.2 (3)	N1—C10—H10	120.2
C6—C1—H1	120.4	C8—C10—H10	120.2
C2—C1—H1	120.4	N3—C11—N2	114.2 (2)
C3—C2—C1	120.1 (3)	N3—C11—S1	126.51 (18)
C3—C2—H2	119.9	N2—C11—S1	119.31 (18)
C1—C2—H2	119.9	C17—C12—C13	120.0 (2)
C2—C3—C4	120.7 (3)	C17—C12—N3	121.7 (2)
C2—C3—H3	119.6	C13—C12—N3	118.1 (2)
C4—C3—H3	119.6	C12—C13—C14	119.5 (2)
C3—C4—C5	120.8 (3)	C12—C13—H13	120.2
C3—C4—H4	119.6	C14—C13—H13	120.2
C5—C4—H4	119.6	C15—C14—C13	120.6 (2)
C4—C5—C6	117.8 (2)	C15—C14—H14	119.7
C4—C5—C9	124.8 (2)	C13—C14—H14	119.7
C6—C5—C9	117.3 (2)	C16—C15—C14	119.5 (2)
O1—C6—C1	117.7 (3)	C16—C15—H15	120.2
O1—C6—C5	120.8 (2)	C14—C15—H15	120.2
C1—C6—C5	121.3 (3)	C15—C16—C17	121.0 (2)
O1—C7—C8	114.2 (2)	C15—C16—H16	119.5
O1—C7—H7A	108.7	C17—C16—H16	119.5
C8—C7—H7A	108.7	C12—C17—C16	119.3 (2)
O1—C7—H7B	108.7	C12—C17—H17	120.3
C8—C7—H7B	108.7	C16—C17—H17	120.3
H7A—C7—H7B	107.6	C10—N1—N2	115.8 (2)
C9—C8—C10	123.8 (2)	C11—N2—N1	120.31 (19)

C9—C8—C7	117.5 (2)	C11—N2—H2A	119.8
C10—C8—C7	118.4 (2)	N1—N2—H2A	119.8
C8—C9—C5	121.8 (2)	C11—N3—C12	127.5 (2)
C8—C9—C11	121.0 (2)	C11—N3—H3A	116.3
C5—C9—C11	117.22 (17)	C12—N3—H3A	116.3
N1—C10—C8	119.6 (2)	C6—O1—C7	117.04 (19)
C6—C1—C2—C3	-0.3 (4)	C9—C8—C10—N1	179.3 (2)
C1—C2—C3—C4	0.2 (4)	C7—C8—C10—N1	5.2 (3)
C2—C3—C4—C5	0.6 (4)	C17—C12—C13—C14	-0.8 (3)
C3—C4—C5—C6	-1.2 (4)	N3—C12—C13—C14	-176.0 (2)
C3—C4—C5—C9	176.4 (2)	C12—C13—C14—C15	-0.4 (4)
C2—C1—C6—O1	-174.5 (2)	C13—C14—C15—C16	1.1 (4)
C2—C1—C6—C5	-0.3 (4)	C14—C15—C16—C17	-0.7 (4)
C4—C5—C6—O1	175.1 (2)	C13—C12—C17—C16	1.2 (3)
C9—C5—C6—O1	-2.7 (3)	N3—C12—C17—C16	176.3 (2)
C4—C5—C6—C1	1.1 (3)	C15—C16—C17—C12	-0.5 (4)
C9—C5—C6—C1	-176.8 (2)	C8—C10—N1—N2	-176.1 (2)
O1—C7—C8—C9	27.7 (3)	N3—C11—N2—N1	8.2 (3)
O1—C7—C8—C10	-157.8 (2)	S1—C11—N2—N1	-172.62 (17)
C10—C8—C9—C5	-177.1 (2)	C10—N1—N2—C11	-179.3 (2)
C7—C8—C9—C5	-2.9 (3)	N2—C11—N3—C12	-176.3 (2)
C10—C8—C9—C11	3.5 (3)	S1—C11—N3—C12	4.7 (4)
C7—C8—C9—C11	177.74 (17)	C17—C12—N3—C11	52.3 (3)
C4—C5—C9—C8	172.2 (2)	C13—C12—N3—C11	-132.6 (2)
C6—C5—C9—C8	-10.1 (3)	C1—C6—O1—C7	-157.2 (2)
C4—C5—C9—C11	-8.4 (3)	C5—C6—O1—C7	28.6 (3)
C6—C5—C9—C11	169.29 (17)	C8—C7—O1—C6	-40.3 (3)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C12–C17 phenyl ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2A...S1 ⁱ	0.86	2.61	3.456 (2)	167
C2—H2...Cg1 ⁱⁱ	0.93	2.86	3.697 (3)	151

Symmetry codes: (i) $-x+2, -y+2, -z$; (ii) $-x+1, -y, -z$.