

Crystal structure of diethyl 2,2'-[[(1*E*,1'*E*)-[[*(1R,4R)*-cyclohexane-1,4-diyl]bis(azanilylidene)]]bis(methanylylidene))bis(1*H*-pyrrole-2,1-diyl)]diacetate

Jasim Alshawi,^a Muoayed Yousif,^a Karzan H. Zangana,^b Inigo J. Vitorica Yrezabal,^b Richard Winpenny^b and Mohamad J. Al-Jeboori^{c*}

^aDepartment of Chemistry, College of Education for Pure Science, University of Basrah, Iraq, ^bSchool of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, England, and ^cDepartment of Chemistry, College of Education (Ibn Al-Haitham) for Pure Science, University of Baghdad, Iraq. *Correspondence e-mail: mohamad.aljeboori@yahoo.com

Received 6 February 2015; accepted 8 February 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


The whole molecule of the title compound, C₂₄H₃₂N₄O₄, is generated by inversion symmetry. The cyclohexane ring adopts a chair conformation and the conformation about the C=N bonds is *E*. The pyrrole rings have an *anti* confirmation with respect to the cyclohexane moiety and the ethyl acetate groups have extended conformations. In the crystal, molecules are linked by pairs of C—H...O hydrogen bonds forming chains, enclosing R₂²(10) ring motifs with inversion symmetry, propagating parallel to the (101) plane.

Keywords: crystal structure; Schiff base; bispyrrole; C—H...O hydrogen bonding.

CCDC reference: 1048163

1. Related literature

For general background on the applications of Schiff bases and the use of pyrrole compounds, see: Köse *et al.* (2015); Trofimov *et al.* (2015). For the synthesis of dipyrrole Schiff bases ligands, see: Meghdadi *et al.* (2010); Munro *et al.* (2004). For the synthesis of pyrrole ester precursors, see: Koriatopoulou *et al.* (2008); Singh & Pal (2010). For the preparation of Schiff bases, see: Yang *et al.* (2004); Ourari *et al.* (2013).



2. Experimental

2.1. Crystal data

C ₂₄ H ₃₂ N ₄ O ₄	$\gamma = 113.288 (8)^\circ$
$M_r = 440.54$	$V = 587.68 (10) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 8.5531 (6) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.8379 (7) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 9.6492 (9) \text{ \AA}$	$T = 150 \text{ K}$
$\alpha = 115.166 (9)^\circ$	$0.4 \times 0.3 \times 0.3 \text{ mm}$
$\beta = 92.105 (7)^\circ$	

2.2. Data collection

Agilent SuperNova (single source at offset, Atlas) diffractometer	4657 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2013)	2734 independent reflections
$T_{\min} = 0.933$, $T_{\max} = 1.000$	1827 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.028$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	146 parameters
$wR(F^2) = 0.124$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
2734 reflections	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C14—H14A...O16 ⁱ	0.97	2.50	3.317 (3)	142

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS2014* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2014* and *PLATON*.

Acknowledgements

The authors are grateful to the Iraqi Ministry for Higher Education for providing six months funding for JA's PhD scholarship.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU5080).

References

Agilent (2013). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.

- Koriatopoulou, K., Karousis, N. & Varvounis, G. (2008). *Tetrahedron*, **64**, 10009–10013.
- Köse, M., Ceyhan, G., Tümer, M., Demirtaş, İ., Gönül, İ. & McKee, V. (2015). *Spectrochim. Acta Part A*, **137**, 477–485.
- Meghdadi, S., Amirnasr, M., Mereiter, K. & Karimi Abdolmaleki, M. (2010). *Acta Cryst. E* **66**, m332–m333.
- Munro, O., Strydom, S. & Grimmer, C. (2004). *New J. Chem.* **28**, 34–42.
- Ourari, A., Aggoun, D. & Ouahab, L. (2013). *Inorg. Chem. Commun.* **33**, 118–124.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Singh, K. & Pal, D. (2010). *J. Serb. Chem. Soc.* **75**, 917–927.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Trofimov, B. A., Mikhaleva, A., Schmidt, E. Y. & Sobenina, L. N. (2015). In *Chemistry of Pyrroles*. Boca Raton: CRC Press.
- Yang, L., Shan, X., Chen, Q., Wang, Z. & Ma, J. S. (2004). *Eur. J. Inorg. Chem.* **2004**, 1474–1477.

supporting information

Acta Cryst. (2015). E71, o165–o166 [doi:10.1107/S2056989015002674]

Crystal structure of diethyl 2,2'-[((1*E*,1'*E*)-{[(1*R*,4*R*)-cyclohexane-1,4-diyl]bis-(azanylylidene)}bis(methanylylidene))bis(1*H*-pyrrole-2,1-diyl)]diacetate

Jasim Alshawi, Muoayed Yousif, Karzan H. Zangana, Inigo J. Vitorica Yrezabal, Richard Winpenny and Mohamad J. Al-Jeboori

S1. Synthesis

The title compound was prepared in a two step procedure:

Synthesis of ethyl (2-formyl-1*H*-pyrrole-1-yl)-acetate (L): prepared by reported procedures (Koriatopoulou *et al.*, 2008; Singh & Pal, 2010) as follows: To a mixture of 1*H*-pyrrole-2-carbaldehyde (1.00 g, 10.51 mmol), K₂CO₃ (2.90g, 21.02 mmol) and (2.64 g, 10.51 mmol) of 18-crown-6 in dry 1,4-dioxane (20ml), was added drop wise a solution of ethyl bromoacetate (2.00 g, 12 mmol) in dry 1,4-dioxane (20 ml), over a period of 30 min. The reaction mixture was allowed to reflux under a nitrogen atmosphere for 6 h, and then the solvent was removed under reduced pressure. Water (50ml) was added to the residue, and the mixture was extracted with ethyl acetate (3 × 15ml). The combined organic layers were washed with brine (15 ml), and then dried over Na₂SO₄. The solvent was removed under reduced pressure, and the oily residue was purified by flash chromatography with an eluent mixture (33% ethyl acetate / hexane), giving compound (L) as a yellow oil product (yield: 0.75 g, 75%). NMR data (p.p.m), δ H: (500 MHz, CDCl₃): 1.20 (3H, t, C12—H), 4.15 (2H, q, C11—H), 4.97 (2H, s, C8—H), 6.21 (1H, t, C3—H), 6.84 (1H, d, C4—H), 6.90 (1H, d, C2—H) and 9.45 (1H, s, C6—H); δ C (125.75 MHz, CDCl₃), 14.13 C12, 50.25 C8, 61.63 C11, 110.20 C3, 124.61 C4, 131.71 C5 and 132.10 C2. C=O for the carboxylate moiety at 168.37 (C12) and at 179.74 for C6. The positive ES mass spectrum at $m/z = 182.4$ (M+H)⁺ (62%) for C₉H₁₁NO₃, requires = 181.1. The other peaks detected at $m/z = 153.4$ (100%), 109.3 (6%), 95 (9%) and 67 (4%) correspond to [M—CH₂CH₃]⁺, [M-(CH₂CH₃+CO₂)]⁺, [M-(CH₂CH₃+CO₂+CH₂)]⁺ and [M-(CH₂CH₃+CO₂+CH₂+CO)]⁺, respectively. IR (ATR cm⁻¹): 1650 ν (C=O) aldehyde moiety. 1710 ν (C=O) ester group.

Synthesis of the title Schiff-base: performed using conventional procedures (Yang *et al.*, 2004; Ourari *et al.*, 2013). To a mixture of L (1.81 g, 10 mmol) in ethanol (20 ml) with 3 drops of glacial acetic acid, a solution of 1,4-diaminocyclohexan (0.57 g, 5 mmol) in ethanol (20ml) was added drop wise over a period of 20 min. The reaction mixture was allowed to reflux for 3h, and then cooled to room temperature. A white precipitate was collected by filtration and recrystallised from ethanol (yield: 1.09g, 60%). Crystals were obtained by slow evaporation of a solution in methanol/acetone. NMR data (p.p.m), δ H (500 MHz, CDCl₃): 1.19 (6H, t, C15, 15—H), 1.47 (C10, 10-, 4H, q), δ H = 1.67 (C9, 9-, 4H, q), 2.94 (2H, p, C8, 8—H), 4.10 (4H, q, C14, 14—H), 5.03 (C11, 11—H, 4H, s), 6.11 (2H, t, C3, 3—H), 6.38 (2H, d, C4, 4—H), 6.61 (2H, d, C2, 2—H) and 8.07 (2H, s, C6, 6—H); δ C (125.75 MHz, CDCl₃), 14.28 (C15, 15-), 32.61 (C10, 10- and C19, 9-), 51.13 (C11, 11-), 61.07 (C14, 14-), 68.8 (C8, 8-), 108.53 (C2, 2-), 116.58 (C5, 5-), 127.69 (C3, 3-), 129.93 (C4, 4-), 150.04 (C6, 6-), C=O 169.37 (C12, 12-). The positive ES mass spectrum at $m/z = 441.52$ (M+H)⁺ (100%) for C₂₄H₃₂N₄O₄, requires = 440.24. The other peaks detected at $m/z = 412.42$ (5%), 383 (3%), 295.19 (9%) and 267.1 (4%) correspond to [M—CH₂CH₃]⁺, [M-(2CH₂CH₃)]⁺, [M-(2CH₂CH₃+2CO₂)]⁺ and [M(2CH₂CH₃+2CO₂+2CH₂)]⁺, respectively. IR (ATR, cm⁻¹): 1580 (C=N), 1630 (C=O).

S2. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were placed in calculated positions and treated as riding atoms: C—H = 0.95 - 0.99 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and = $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

S3. Comment

The whole molecule of the title compound, Fig.1, is generated by inversion symmetry. The cyclohexane ring adopts a chair conformation and the conformation about the C=N bonds is *E*. The pyrrole rings crystallize in the anti-conformation with respect to the cyclohexane moiety and the ethyl acetate moieties have extended conformations.

In the crystal, molecules are linked by pairs of C—H···O hydrogen bonds forming chains, enclosing $R^2_2(10)$ ring motifs with inversion symmetry, propagating parallel to plane (101); see Table 1 and Fig. 2.

Infrared spectrum indicated typical absorbance bands of the functional —C=N and carbonyl —C=O at 1580 and 1630 cm^{-1} , respectively. The positive ES mass spectrum of the bis Schiff-base showed a parent ion peak at $m/z = 441.52$ ($M+H$)⁺, corresponding to $\text{C}_{26}\text{H}_{32}\text{N}_4\text{O}_4$, for which the required value is 440.24. The N7=C6 bond distance [1.270 (2) Å] is shorter than the N2—C8 bond distance [1.458 (2) Å], indicating a double bond order. However, the N1—C5 bond distance [1.384 (2) Å] indicates resonance has occurred in the pyrrole system between the lone pair electron of the nitrogen atom and the pyrrole ring.





Figure 1

A view of the molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A view along the *b* axis of the crystal packing of the title compound. The C—H...O hydrogen bonds are drawn as dashed lines (see Table 1 for details; H atom not involved in hydrogen bonding have been omitted for clarity).

Diethyl 2,2'-[[(1*E*,1'*E*)-[(1*R*,4*R*)-cyclohexane-1,4-diyl]bis(azanylylidene)]bis(methanylylidene)]bis(1*H*-pyrrole-2,1-diyl)]diacetate

Crystal data

$C_{24}H_{32}N_4O_4$

$M_r = 440.54$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 8.5531\ (6)\ \text{\AA}$

$b = 8.8379\ (7)\ \text{\AA}$

$c = 9.6492\ (9)\ \text{\AA}$

$\alpha = 115.166\ (9)^\circ$

$\beta = 92.105\ (7)^\circ$

$\gamma = 113.288\ (8)^\circ$

$V = 587.68\ (10)\ \text{\AA}^3$

$Z = 1$

$F(000) = 236$

$D_x = 1.245\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1515 reflections

$\theta = 3.3\text{--}26.7^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 150\ \text{K}$

Block, colourless

$0.4 \times 0.3 \times 0.3\ \text{mm}$

Data collection

Agilent SuperNova (single source at offset, Atlas)

diffractometer

Radiation source: SuperNova (Mo) X-ray Source

Mirror monochromator

Detector resolution: $10.37\ \text{pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.933$, $T_{\max} = 1.000$

4657 measured reflections

2734 independent reflections

1827 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 29.4^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -11 \rightarrow 11$

$k = -10 \rightarrow 12$

$l = -13 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.124$
 $S = 1.07$
 2734 reflections
 146 parameters
 0 restraints

Primary atom site location: structure-invariant
 direct methods
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0363P)^2 + 0.1039P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O13	0.46898 (16)	0.46998 (17)	0.19574 (15)	0.0317 (3)
O16	0.63475 (17)	0.71083 (17)	0.43494 (15)	0.0338 (4)
N1	0.67517 (19)	0.9650 (2)	0.32094 (18)	0.0274 (4)
N7	0.87176 (19)	0.7539 (2)	0.17968 (18)	0.0313 (4)
C12	0.5599 (2)	0.6517 (3)	0.3013 (2)	0.0270 (4)
C6	0.9477 (2)	0.9265 (3)	0.2821 (2)	0.0297 (5)
H6	1.0695	0.9854	0.3163	0.036*
C5	0.8559 (2)	1.0361 (2)	0.3483 (2)	0.0275 (4)
C10	0.9066 (2)	0.4733 (2)	0.1181 (2)	0.0311 (5)
H10A	0.7851	0.3992	0.0575	0.037*
H10B	0.9087	0.4935	0.2252	0.037*
C11	0.5474 (2)	0.7689 (2)	0.2290 (2)	0.0279 (4)
H11A	0.5653	0.7192	0.1233	0.033*
H11B	0.4302	0.7597	0.2201	0.033*
C2	0.6355 (3)	1.1083 (3)	0.4039 (2)	0.0317 (5)
H2	0.5229	1.0970	0.4061	0.038*
C9	1.0097 (2)	0.3654 (3)	0.0461 (2)	0.0307 (5)
H9A	0.9545	0.2438	0.0420	0.037*
H9B	1.1280	0.4334	0.1127	0.037*
C8	0.9819 (2)	0.6620 (2)	0.1201 (2)	0.0304 (5)
H8	1.1007	0.7412	0.1897	0.036*
C4	0.9269 (3)	1.2260 (3)	0.4484 (2)	0.0340 (5)
H4	1.0459	1.3098	0.4864	0.041*
C3	0.7880 (3)	1.2712 (3)	0.4833 (2)	0.0377 (5)
H3	0.7980	1.3898	0.5483	0.045*
C14	0.4577 (3)	0.3383 (3)	0.2513 (2)	0.0400 (5)
H14A	0.4025	0.3576	0.3392	0.048*
H14B	0.5743	0.3562	0.2869	0.048*
C15	0.3512 (3)	0.1451 (3)	0.1175 (3)	0.0544 (7)

H15A	0.4111	0.1242	0.0342	0.082*
H15B	0.2388	0.1314	0.0787	0.082*
H15C	0.3349	0.0551	0.1534	0.082*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O13	0.0352 (8)	0.0259 (7)	0.0293 (8)	0.0091 (6)	0.0017 (6)	0.0141 (6)
O16	0.0378 (8)	0.0349 (7)	0.0245 (8)	0.0133 (6)	0.0036 (6)	0.0140 (6)
N1	0.0295 (8)	0.0254 (8)	0.0269 (9)	0.0133 (7)	0.0060 (6)	0.0114 (7)
N7	0.0275 (8)	0.0302 (8)	0.0281 (9)	0.0150 (7)	0.0050 (7)	0.0054 (7)
C12	0.0238 (10)	0.0303 (10)	0.0253 (10)	0.0116 (8)	0.0079 (8)	0.0124 (8)
C6	0.0255 (10)	0.0337 (10)	0.0262 (10)	0.0119 (9)	0.0061 (8)	0.0126 (9)
C5	0.0291 (10)	0.0282 (10)	0.0231 (10)	0.0120 (8)	0.0062 (8)	0.0115 (8)
C10	0.0250 (10)	0.0340 (10)	0.0249 (10)	0.0115 (9)	0.0054 (8)	0.0083 (8)
C11	0.0250 (10)	0.0305 (10)	0.0265 (10)	0.0120 (8)	0.0052 (8)	0.0129 (8)
C2	0.0412 (11)	0.0355 (10)	0.0299 (11)	0.0252 (10)	0.0127 (9)	0.0178 (9)
C9	0.0259 (10)	0.0280 (10)	0.0300 (11)	0.0104 (8)	0.0039 (8)	0.0089 (8)
C8	0.0204 (9)	0.0302 (10)	0.0293 (11)	0.0116 (8)	0.0034 (7)	0.0050 (8)
C4	0.0361 (11)	0.0269 (10)	0.0308 (11)	0.0105 (9)	0.0062 (9)	0.0106 (9)
C3	0.0519 (13)	0.0269 (10)	0.0341 (12)	0.0207 (10)	0.0109 (10)	0.0118 (9)
C14	0.0480 (13)	0.0336 (11)	0.0396 (13)	0.0130 (10)	0.0065 (10)	0.0240 (10)
C15	0.0690 (16)	0.0316 (11)	0.0498 (15)	0.0101 (12)	−0.0021 (12)	0.0217 (11)

Geometric parameters (Å, °)

O13—C12	1.336 (2)	C6—C5	1.441 (2)
O13—C14	1.448 (2)	C5—C4	1.375 (2)
O16—C12	1.203 (2)	C10—C9	1.522 (2)
N1—C5	1.384 (2)	C10—C8	1.522 (3)
N1—C11	1.452 (2)	C2—C3	1.365 (3)
N1—C2	1.363 (2)	C9—C8 ⁱ	1.526 (3)
N7—C6	1.270 (2)	C8—C9 ⁱ	1.526 (3)
N7—C8	1.458 (2)	C4—C3	1.405 (3)
C12—C11	1.506 (3)	C14—C15	1.490 (3)
C12—O13—C14	116.15 (15)	C4—C5—C6	127.83 (17)
C5—N1—C11	126.32 (15)	C8—C10—C9	111.74 (14)
C2—N1—C5	108.70 (15)	N1—C11—C12	112.58 (15)
C2—N1—C11	124.84 (15)	N1—C2—C3	108.91 (17)
C6—N7—C8	117.71 (15)	C10—C9—C8 ⁱ	111.47 (17)
O13—C12—C11	109.47 (16)	N7—C8—C10	109.64 (14)
O16—C12—O13	124.75 (19)	N7—C8—C9 ⁱ	109.49 (17)
O16—C12—C11	125.74 (17)	C10—C8—C9 ⁱ	110.20 (15)
N7—C6—C5	123.74 (17)	C5—C4—C3	107.99 (18)
N1—C5—C6	124.91 (15)	C2—C3—C4	107.14 (17)
C4—C5—N1	107.27 (16)	O13—C14—C15	107.88 (18)

O13—C12—C11—N1	-166.07 (14)	C11—N1—C5—C6	3.5 (3)
O16—C12—C11—N1	16.3 (3)	C11—N1—C5—C4	-176.36 (17)
N1—C5—C4—C3	0.4 (2)	C11—N1—C2—C3	176.43 (17)
N1—C2—C3—C4	-0.3 (2)	C2—N1—C5—C6	179.33 (19)
N7—C6—C5—N1	6.4 (3)	C2—N1—C5—C4	-0.6 (2)
N7—C6—C5—C4	-173.7 (2)	C2—N1—C11—C12	-110.7 (2)
C12—O13—C14—C15	179.68 (16)	C9—C10—C8—N7	176.06 (15)
C6—N7—C8—C10	134.27 (18)	C9—C10—C8—C9 ⁱ	55.5 (2)
C6—N7—C8—C9 ⁱ	-104.71 (19)	C8—N7—C6—C5	178.46 (18)
C6—C5—C4—C3	-179.5 (2)	C8—C10—C9—C8 ⁱ	-56.2 (2)
C5—N1—C11—C12	64.5 (2)	C14—O13—C12—O16	2.0 (3)
C5—N1—C2—C3	0.6 (2)	C14—O13—C12—C11	-175.70 (15)
C5—C4—C3—C2	0.0 (2)		

Symmetry code: (i) $-x+2, -y+1, -z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C14—H14A...O16 ⁱⁱ	0.97	2.50	3.317 (3)	142

Symmetry code: (ii) $-x+1, -y+1, -z+1$.