

Crystal structures of dichloridopalladium(II),  
-platinum(II) and -rhodium(III) complexes  
containing 8-(diphenylphosphanyl)quinolineTakayoshi Suzuki,<sup>a\*</sup> Hiroshi Yamaguchi,<sup>b</sup> Masayuki Fujiki,<sup>b</sup> Akira Hashimoto<sup>b</sup> and  
Hideo D. Takagi<sup>b</sup>

Received 7 March 2015

Accepted 25 March 2015


Edited by H. Ishida, Okayama University, Japan

<sup>a</sup>Department of Chemistry, Okayama University, Okayama 700-8530, Japan, and <sup>b</sup>Graduate School of Science and Research Center for Material Science, Nagoya University, Chikusa, Nagoya 464-8602, Japan. \*Correspondence e-mail: suzuki@okayama-u.ac.jp**Keywords:** crystal structure; 8-quinolylphosphane; stacking interaction; geometrical structure; *trans* influence**CCDC references:** 1056041; 1056040; 1056039**Supporting information:** this article has supporting information at journals.iucr.org/e


The crystal structures of dichloridopalladium(II), -platinum(II) and -rhodium(III) complexes containing 8-(diphenylphosphanyl)quinoline, (*SP*-4)-[PdCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)], (1) [systematic name: dichlorido(8-diphenylphosphanylquinoline)palladium(II)], (*SP*-4)-[PtCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)]·CH<sub>2</sub>Cl<sub>2</sub>, (2) [systematic name: dichlorido(8-diphenylphosphanylquinoline)platinum(II) dichloromethane monosolvate], and (*OC*-6–32)-[RhCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)<sub>2</sub>]PF<sub>6</sub>·0.5CH<sub>2</sub>Cl<sub>2</sub>·0.5CH<sub>3</sub>OH, (3) [systematic name: *cis*-dichloridobis(8-diphenylphosphanylquinoline)rhodium(III) hexafluoridophosphate dichloromethane/methanol hemisolvate] are reported. In these complexes, the phosphanylquinoline acts as a bidentate ligand, forming a planar asymmetrical five-membered chelate ring. The palladium(II) and platinum(II) complex molecules in (1) and (2), respectively, show a typical square-planar coordination geometry and form a dimeric structure through an intermolecular  $\pi$ – $\pi$  stacking interaction between the quinolyl rings. The centroid–centroid distances between the stacked six-membered rings in (1) and (2) are 3.633 (2) and 3.644 (2) Å, respectively. The cationic rhodium(III) complex in (3) has a *cis(Cl)*, *cis(P)*, *cis(N)* (*OC*-6–32) configuration of the ligands, in which two kinds of intramolecular  $\pi$ – $\pi$  stacking interactions are observed: between the quinolyl and phenyl rings and between two phenyl rings, the centroid–centroid distances being 3.458 (2) and 3.717 (2) Å, respectively. The PF<sub>6</sub><sup>−</sup> anion in (3) is rotationally disordered, the site occupancies of each F atom being 0.613 (14) and 0.387 (14). The CH<sub>2</sub>Cl<sub>2</sub> and CH<sub>3</sub>OH solvent molecules are also disordered and equal site occupancies of 0.5 are assumed.

## 1. Chemical context

8-Quinolylphosphanes are an interesting class of ligands because they form a planar asymmetrical five-membered chelate ring *via* coordination through quinoline-*N* and phosphane-*P* atoms (Issleib & Hörnig, 1972; Salem & Wild, 1992; Wehman *et al.*, 1997). The electronic differentiation of the donor groups, in particular their  $\pi$ -bonding natures, may stabilize unusual electronic states of their transition metal complexes (Espinet & Soulantica, 1999). In addition, the steric requirement from the quinolyl moiety often has a strong influence on the properties of their metal complexes. For example, the nickel(II) and palladium(II) complexes containing two 8-(diphenylphosphanyl)quinoline (Ph<sub>2</sub>Pqn) molecules with a *cis(P)* configuration showed a severe distortion of the square-planar coordination geometry around Ni<sup>II</sup> and Pd<sup>II</sup> as a result of the steric hindrance between mutually *cis*-positioned quinolyl groups (Suzuki, 2004; Hashimoto *et al.*, 2010). Several crystallographic studies have been performed for other Ph<sub>2</sub>Pqn complexes, as described in




§4, but not for the platinum(II) and rhodium(III) complexes. In 1979, the preparation and spectroscopic characterization of  $[MCl_2(Ph_2Pqn)]$  ( $M = Pd^{II}$ ,  $Pt^{II}$ , and  $Rh^{III}$ ) was reported (Hudali *et al.*, 1979), but the crystal structures of these complexes were not confirmed, except for  $[PdCl_2(Ph_2Pqn)] \cdot CH_2Cl_2$  (Bastanov *et al.*, 2009). In particular, it is worthwhile to reinvestigate the rhodium(III) complex because it was prepared from  $RhCl_3 \cdot 3H_2O$  and  $Ph_2Pqn$  in acetone (Hudali *et al.*, 1979).




## 2. Structural commentary

A yellow block-shaped crystal of the  $Pd^{II}$  complex, (*SP-4*)- $[PdCl_2(Ph_2Pqn)]$ , (1), recrystallized from hot acetonitrile, was used for the X-ray diffraction analysis. The complex molecule (Fig. 1) has a typical square-planar coordination geometry




**Figure 1**  
 An ORTEP of the molecular structure of  $[PdCl_2(Ph_2Pqn)]$ , (1), showing the atom-numbering scheme, with displacement ellipsoids drawn at the 50% probability level.




**Figure 2**  
 An ORTEP of the molecular structure of  $[PtCl_2(Ph_2Pqn)] \cdot CH_2Cl_2$ , (2), showing the atom-numbering scheme, with displacement ellipsoids drawn at the 50% probability level.

with a chelating  $Ph_2Pqn$  ligand, whose  $P1-Pd1-N1$  bite angle is  $84.75(6)^\circ$ . The quinolyl plane is almost co-planar to the  $Pd^{II}$  coordination plane; the dihedral angle between these planes is only  $8.58(3)^\circ$ . The two  $Pd-Cl$  bonds show a significant difference in length [ $Pd1-Cl1$  2.3716 (6) vs  $Pd1-Cl2$  2.2885 (7) Å], indicating a strong *trans* influence of the phosphane donor group. The corresponding  $Pd-Cl$  bond in *cis*(*P*)- $[PdCl(Ph_2Pqn)_2]BF_4$ , which is also *trans* to the phosphane donor of  $Ph_2Pqn$ , was similarly long at 2.375 (2) Å (Suzuki, 2004). On the other hand, the  $Pd1-P1$  bond [2.2026 (6) Å] in (1) is slightly shorter than those in *cis*(*P*)- $[PdCl(Ph_2Pqn)_2]BF_4$  and *cis*(*P*)- $[Pd(Ph_2Pqn)_2]X_2$  ( $X = Cl$  or  $Br$ ) [2.229 (2)–2.267 (2) Å], presumably due to the steric congestion in the above bis( $Ph_2Pqn$ )-type complexes. The  $Pd1-N1$  bond length in (1) is 2.065 (2) Å. The dihedral angles



**Figure 3**  
 An ORTEP of the complex molecule in (*OC-6-32*)- $[RhCl_2(Ph_2Pqn)_2]PF_6 \cdot 0.5CH_2Cl_2 \cdot 0.5CH_3OH$ , (3), showing the atom-numbering scheme, with displacement ellipsoids drawn at 30% probability level. Hydrogen atoms are omitted for clarity.




**Figure 4**  
Possible configurations and notation for the  $[\text{RhCl}_2(\text{P-N})_2]^+$  complex cation.


between the quinolyl ring system and the two phenyl rings of the coordinated  $\text{Ph}_2\text{Pqn}$  are  $72.34$  (8) and  $74.79$  (8)°.

When the platinum(II) complex was recrystallized from dichloromethane, the resulting crystals contained a  $\text{CH}_2\text{Cl}_2$  molecule per a complex molecule:  $[\text{PtCl}_2(\text{Ph}_2\text{Pqn})]\cdot\text{CH}_2\text{Cl}_2$  (2). The X-ray analysis revealed that it was isomorphous with the  $\text{Pd}^{\text{II}}$  analogue,  $[\text{PdCl}_2(\text{Ph}_2\text{Pqn})]\cdot\text{CH}_2\text{Cl}_2$ , which has been deposited in the Cambridge Structural Database (Bastanov *et al.*, 2009). The molecular structure of the  $\text{Pt}^{\text{II}}$  complex moiety with a square-planar coordination geometry (Fig. 2) is very similar to the above  $\text{Pd}^{\text{II}}$  complex in (1). The  $\text{Pt1-P1}$  and  $\text{Pt1-N1}$  bond lengths are  $2.1963$  (6) and  $2.051$  (2) Å, respectively, and the  $\text{Ph}_2\text{Pqn}$  bite angle ( $\text{P1-Pt1-N1}$ ) is  $85.44$  (6)°. The  $\text{Pt1-Cl1}$  and  $\text{Pt1-Cl2}$  bond lengths are  $2.3747$  (6) and  $2.3002$  (7) Å, respectively, also indicative of a strong *trans* influence of the phosphane donor group.

Pale yellow prismatic crystals of  $[\text{RhCl}_2(\text{Ph}_2\text{Pqn})_2]\text{PF}_6\cdot 0.5\text{CH}_2\text{Cl}_2\cdot 0.5\text{CH}_3\text{OH}$  (3) were analyzed by the X-ray diffraction method, and it was revealed that the complex cation has an octahedral coordination geometry with a *cis(Cl),cis(P),cis(N)* (OC-6-32) configuration (Figs. 3 and 4). As a result of the strong *trans* influence of the phosphane donor, the two  $\text{Rh-Cl}$  and the two  $\text{Rh-N}$  bond lengths are significantly different from each other. The  $\text{Rh1-Cl1}$  bond [ $2.3787$  (6) Å] is longer by  $0.045$  Å than  $\text{Rh1-Cl2}$  [ $2.3338$  (7) Å], while  $\text{Rh1-N1}$  [ $2.168$  (2) Å] is longer by  $0.10$  Å than  $\text{Rh1-N11}$  [ $2.065$  (2) Å]. This fact suggests that



**Figure 5**  
A view of the crystal packing of  $[\text{PdCl}_2(\text{Ph}_2\text{Pqn})]$ , (1), illustrating the  $\pi$ - $\pi$  stacking interactions between the complexes. Color code: Pd, purple; Cl, green; P, yellow; N, blue; C, black; and H, gray.



**Figure 6**  
A view of the crystal packing of  $[\text{PtCl}_2(\text{Ph}_2\text{Pqn})]\cdot\text{CH}_2\text{Cl}_2$ , (2), illustrating the  $\pi$ - $\pi$  stacking interactions between the complexes. Color code: Pt, purple; Cl, green; P, yellow; N, blue; C, black; and H, gray.

the *trans* influence of the phosphane donor is much effective for the  $\text{Rh-N}$ (quinoline) bond rather than the  $\text{Rh-Cl}$  bond. Two slightly deviated  $\text{Rh-P}$  bond lengths [ $\text{Rh1-P1}$   $2.2897$  (7) vs.  $\text{Rh1-P2}$   $2.2531$  (8) Å] seem to result from different steric congestion around the P donor atoms. The larger bond angle of  $\text{P1-Rh1-P2}$  [ $100.55$  (3)°] than the ideal right angle is also suggestive of steric interaction between the two phosphane groups. However, the molecular structure of the complex cation (Fig. 3) also suggests an intramolecular  $\pi$ - $\pi$  stacking interaction between the C27-C32 and C39-C44 phenyl rings. The centroid-centroid distance between these rings is  $3.717$  (2) Å. An other intramolecular  $\pi$ - $\pi$  stacking interaction is also found between the N11/C12-C14/C20/C19 ring of the quinolyl substituent and the C21-C26 phenyl ring, the centroid-centroid distance being  $3.458$  (2) Å. These interactions could stabilize the *cis(Cl),cis(P),cis(N)* configuration of the  $\text{Rh}^{\text{III}}$  complex cation  $[\text{RhCl}_2(\text{Ph}_2\text{Pqn})_2]^+$ .

The crystal structures of the related complexes with (2-aminoethyl)diphenylphosphane,  $[\text{RhCl}_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{-NH}_2)_2]^+$ , were reported to have the *trans(Cl),cis(P)* (OC-6-13) or *cis(Cl),trans(P)* (OC-6-33) configuration (Fig. 4) (Galsbøl *et al.*, 1986). If the *trans(Cl),cis(P)* configuration were assumed for the present  $\text{Ph}_2\text{Pqn}$  complex, the molecule would have severe steric hindrance between the *ortho*-H atoms of the mutually *cis*-positioned quinolyl groups, as observed in the crystal structures of *cis(P)*- $[\text{Pd}(\text{Ph}_2\text{Pqn})_2]\text{X}_2$  (Suzuki, 2004). The *trans(P)* configurations, i.e., *trans(Cl),trans(P)* (OC-6-12) and *cis(Cl),trans(P)* (OC-6-33), would be unfavorable due to the mutually *trans* disposition of the phosphane groups having a strong *trans* influence. The last configuration, *cis(Cl),trans(N)* (OC-6-22), cannot form an intramolecular stacking interaction between the aryl groups of the phosphanes. Therefore, the observed *cis(Cl),cis(P),cis(N)* geome-

**Table 1**  
Experimental details.

	(1)	(2)	(3)
Crystal data			
Chemical formula	[PdCl <sub>2</sub> (C <sub>21</sub> H <sub>16</sub> NP)]	[PtCl <sub>2</sub> (C <sub>21</sub> H <sub>16</sub> NP)]·CH <sub>2</sub> Cl <sub>2</sub>	[RhCl <sub>2</sub> (C <sub>21</sub> H <sub>16</sub> NP) <sub>2</sub> ](PF <sub>6</sub> ) <sub>2</sub> ·0.5CH <sub>2</sub> Cl <sub>2</sub> ·0.5CH <sub>4</sub> O
<i>M<sub>r</sub></i>	490.62	664.23	1003.90
Crystal system, space group	Monoclinic, <i>P</i> <sub>2</sub> / <i>n</i>	Monoclinic, <i>P</i> <sub>2</sub> / <i>c</i>	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	200	200	200
<i>a</i> , <i>b</i> , <i>c</i> (Å)	9.0293 (5), 15.2154 (8), 13.7936 (6)	13.9280 (5), 9.2371 (3), 17.8941 (6)	9.841 (5), 13.825 (6), 16.167 (8)
$\alpha$ , $\beta$ , $\gamma$ (°)	90, 91.8197 (13), 90	90, 102.8447 (10), 90	87.307 (19), 81.80 (2), 70.819 (18)
<i>V</i> (Å <sup>3</sup> )	1894.07 (16)	2244.53 (13)	2056.2 (17)
<i>Z</i>	4	4	2
Radiation type	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$	Mo <i>K</i> $\alpha$
$\mu$ (mm <sup>-1</sup> )	1.35	6.81	0.79
Crystal size (mm)	0.18 × 0.15 × 0.12	0.25 × 0.24 × 0.05	0.20 × 0.20 × 0.15
Data collection			
Diffractometer	Rigaku R-AXIS RAPID	Rigaku R-AXIS RAPID	Rigaku R-AXIS RAPID
Absorption correction	Numerical ( <i>NUMABS</i> ; Rigaku, 1999)	Numerical ( <i>NUMABS</i> ; Rigaku, 1999)	Multi-scan ( <i>ABSCOR</i> ; Rigaku, 1995)
<i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub>	0.693, 0.850	0.281, 0.727	0.848, 0.882
No. of measured, independent and observed [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )] reflections	18103, 4283, 3913	20995, 5081, 4665	20385, 9338, 7245
<i>R</i> <sub>int</sub>	0.034	0.028	0.046
( <i>sin</i> $\theta$ / $\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )	0.649	0.648	0.649
Refinement			
<i>R</i> [ <i>F</i> <sup>2</sup> > 2 $\sigma$ ( <i>F</i> <sup>2</sup> )], <i>wR</i> [ <i>F</i> <sup>2</sup> ], <i>S</i>	0.028, 0.074, 0.92	0.017, 0.037, 0.98	0.044, 0.120, 1.06
No. of reflections	4283	5081	9338
No. of parameters	235	262	605
No. of restraints	0	0	20
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
$\Delta\rho_{\max}$ , $\Delta\rho_{\min}$ (e Å <sup>-3</sup> )	0.84, -0.50	0.58, -0.60	0.68, -1.03

Computer programs: *PROCESS-AUTO* (Rigaku, 1998), *CrystalStructure* (Rigaku, 2010), *DIRDIF99-PATY* (Beurskens *et al.*, 1999), *SHELXS2013* (Sheldrick, 2008), *SHELXL2013* (Sheldrick, 2015) and *ORTEP-3 for Windows* (Farrugia, 2012).

trical isomer could be the most favorable from the steric and electronic points of views.

### 3. Supramolecular features

In the crystal structure of (1), there is an intermolecular  $\pi$ - $\pi$  stacking interaction between the quinolyl planes, forming an inversion dimer (Fig. 5). The centroid-centroid distance between the N1/C2-C4/C10/C9 ring and the C5<sup>i</sup>-C10<sup>i</sup> ring of the neighbouring molecule [symmetry code: (i) 1 - *x*, 1 - *y*, 2 - *z*] is 3.633 (2) Å.

The Pt<sup>II</sup> complex in (2) also forms an inversion dimer unit by an intermolecular  $\pi$ - $\pi$  stacking interaction between the quinolyl rings of neighbouring molecules (Fig. 6). The centroid-centroid distance between the N1/C2-C4/C10/C9 ring and the C5<sup>ii</sup>-C10<sup>ii</sup> ring of the neighbouring molecule [symmetry code: (ii) 1 - *x*, -*y*, 1 - *z*] is 3.644 (2) Å.

No remarkable intermolecular stacking or hydrogen-bonding interactions are observed in the crystal structure of (3).

### 4. Database survey

The crystal structure of Ph<sub>2</sub>Pqn was reported previously (Nag *et al.*, 2010). Several metal complexes containing Ph<sub>2</sub>Pqn have also reported by us and others, *e.g.*, [Ni(Ph<sub>2</sub>Pqn)<sub>2</sub>](BF<sub>4</sub>)<sub>*n*</sub> (*n* =

1 or 2; Hashimoto *et al.*, 2010), [Pd(Ph<sub>2</sub>Pqn)<sub>2</sub>]*X*<sub>2</sub> (*X* = Cl, Br, or BF<sub>4</sub>; Suzuki, 2004), [Ru(bpy)<sub>2</sub>(Ph<sub>2</sub>Pqn)](PF<sub>6</sub>)<sub>2</sub> (bpy = 2,2-bipyridine; Suzuki *et al.*, 2002), [Cp\*Ir(N<sub>3</sub>)(Ph<sub>2</sub>Pqn)] (Cp\* = pentamethylcyclopentadienyl; Suzuki *et al.*, 2009), [Cu(Ph<sub>2</sub>Pqn)<sub>2</sub>]*BF*<sub>4</sub> (Suzuki *et al.*, 2011), [NiCl(C<sub>10</sub>H<sub>7</sub>)(Ph<sub>2</sub>Pqn)] (C<sub>10</sub>H<sub>7</sub> = 1-naphthyl; Sun *et al.*, 2002), [Cu(Ph<sub>2</sub>Pqn)<sub>2</sub>]*PF*<sub>6</sub> and [Zn*X*<sub>2</sub>(Ph<sub>2</sub>Pqn)] (*X* = Cl, Br, or I; Tsukuda *et al.*, 2009), [Cu(Ph<sub>2</sub>Pqn)]{(Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>)<sub>2</sub>O}]*BF*<sub>4</sub> (Qin *et al.*, 2009), [AuCl(Ph<sub>2</sub>Pqn)] (Monkowiak *et al.*, 2009), [PdCl(C<sub>3</sub>H<sub>5</sub>)(Ph<sub>2</sub>Pqn)], [Pd(C<sub>3</sub>H<sub>5</sub>)(Ph<sub>2</sub>Pqn)]*ClO*<sub>4</sub>, [Pd(Ph<sub>2</sub>Pqn)(MeOOC≡CCOOMe)] and [Pd(Ph<sub>2</sub>Pqn)]{MeOOC-(Me)C≡CCOOMe}] (C<sub>3</sub>H<sub>5</sub> = allyl; Canovese *et al.*, 2010). In addition, the crystal structure of [PdCl<sub>2</sub>(Ph<sub>2</sub>Pqn)]·CH<sub>2</sub>Cl<sub>2</sub> has been deposited (Bastanov *et al.*, 2009).

### 5. Synthesis and crystallization

The ligand, Ph<sub>2</sub>Pqn, was prepared according to a literature method (Feltham & Metzger, 1971; Aguirre *et al.*, 2007). The dichloridopalladium(II) and platinum(II) complexes, [PdCl<sub>2</sub>(Ph<sub>2</sub>Pqn)] and [PtCl<sub>2</sub>(Ph<sub>2</sub>Pqn)], were prepared by the method reported previously by Hudali *et al.* (1979). The palladium(II) complex was recrystallized from hot acetonitrile to afford yellow block-shaped crystals of [PdCl<sub>2</sub>(Ph<sub>2</sub>Pqn)], (1). Analysis calculated for C<sub>21</sub>H<sub>16</sub>Cl<sub>2</sub>NPPd: C 51.4, H 3.29, N 2.85%. Found: C 51.2, H 3.25, N 2.87%.

The colorless platelet crystals of the platinum(II) complex,  $[\text{PtCl}_2(\text{Ph}_2\text{Pqn})]\cdot\text{CH}_2\text{Cl}_2$ , (2), were obtained by recrystallization from dichloromethane. Analysis calculated for  $\text{C}_{21}\text{H}_{16}\text{Cl}_2\text{NPPt}$ : C 43.5, H 2.78, N 2.42%. Found (after drying completely): C 42.8, H 2.75, N 2.44%.

The  $\text{PF}_6$  salt of the dichloridorhodium(III) complex,  $[\text{RhCl}_2(\text{Ph}_2\text{Pqn})_2]\text{PF}_6$ , was precipitated from a methanol solution of  $\text{RhCl}_3(\text{Ph}_2\text{Pqn})_2(\text{H}_2\text{O})$ , which was prepared by a reaction of  $\text{RhCl}_3\cdot 3\text{H}_2\text{O}$  and two equivalent amounts of  $\text{Ph}_2\text{Pqn}$  in boiling water, by addition of a saturated methanol solution of  $\text{NH}_4\text{PF}_6$ . The crude product was recrystallized from a mixture of dichloromethane and methanol, affording pale-yellow prismatic crystals of  $[\text{RhCl}_2(\text{Ph}_2\text{Pqn})_2]\text{PF}_6\cdot 0.5\text{CH}_2\text{Cl}_2\cdot 0.5\text{CH}_3\text{OH}$  (3). These crystals were efflorescent when they were picked up from the mother liquor. Analysis calculated for  $\text{C}_{42}\text{H}_{32}\text{Cl}_2\text{F}_6\text{N}_2\text{P}_3\text{Rh}\cdot 2\text{H}_2\text{O}$ : C 51.4, H 3.70, N 2.85%. Found (after drying completely): C 51.6, H 3.55, N 2.85%.

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1. All H atoms were refined using a riding model, with  $\text{O}-\text{H} = 0.84 \text{ \AA}$  and  $\text{C}-\text{H} = 0.95$  (aromatic), 0.99 (methylene) or 0.98 (methyl)  $\text{ \AA}$ , and with  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5U_{\text{eq}}(\text{C}, \text{O})$ . During the refinement for (3), each F atom of the  $\text{PF}_6^-$  anion was found to have a large displacement ellipsoid elongated in the direction perpendicular to the  $\text{P}-\text{F}$  bond, which was attributable to rotational disorder of the anion over two positions. The occupancies of each F atom refined to 0.613 (14) and 0.387 (14). In addition, since the crystal structure of (3) contains a void accessible for a solvent molecule, disordered  $\text{CH}_2\text{Cl}_2$  and  $\text{CH}_3\text{OH}$  molecules with equal probabilities of 0.5 were assumed. In the refinement, the  $\text{P}-\text{F}$ ,  $\text{C}-\text{Cl}$  and  $\text{C}-\text{O}$  bond lengths and the  $\text{Cl}-\text{C}-\text{Cl}$  bond angle were restrained to be 1.55 (1), 1.75 (1), 1.42 (2)  $\text{ \AA}$  and  $112.0 (2)^\circ$ , respectively. Rigid bond restraints were also applied for the disordered  $\text{CH}_2\text{Cl}_2$  and  $\text{CH}_3\text{OH}$  molecules.

## Acknowledgements

This work was partly supported by a Grant-in-Aid for Scientific Research No. 25410070 from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

## References

- Aguirre, P. A., Lagos, C. A., Moya, S. A., Zúñiga, C., Vera-Oyarce, C., Sola, F., Peris, G. & Bayón, J. C. (2007). *Dalton Trans.* pp. 5419–5426.
- Bastanov, A. S., Bryant, D. E., Kilner, M. & Neletin, I. S. (2009). Private communication (refcode OBAWIZ). CCDC, Cambridge, England.
- Beurskens, P. T., Beurskens, G., de Gelder, R., García-Granda, S., Israel, R., Gould, R. O. & Smits, J. M. M. (1999). *The DIRDIF99 Program System*. Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
- Canovese, L., Visentin, F., Santo, C., Chessa, G. & Bertolasi, V. (2010). *Organometallics*, **29**, 3027–3038.
- Espinete, P. & Soulantica, K. (1999). *Coord. Chem. Rev.* **193–195**, 499–556.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Feltham, R. D. & Metzger, H. G. (1971). *J. Organomet. Chem.* **33**, 347–355.
- Galsbøl, F., Kojima, M., Ishii, T., Ohba, S., Saito, Y. & Fujita, J. (1986). *Bull. Chem. Soc. Jpn*, **59**, 1701–1707.
- Hashimoto, A., Yamaguchi, H., Suzuki, T., Kashiwabara, K., Kojima, M. & Takagi, H. D. (2010). *Eur. J. Inorg. Chem.* pp. 39–47.
- Hudali, H. A., Kingston, J. V. & Tayim, H. A. (1979). *Inorg. Chem.* **18**, 1391–1394.
- Issleib, K. & Hörnig, K. (1972). *Z. Anorg. Allg. Chem.* **389**, 263–268.
- Monkowitz, U., Zabel, M., Fleck, M. & Yersin, H. (2009). *Z. Naturforsch. Teil B*, **64**, 1513–1524.
- Nag, S., Cibian, M. & Hanan, G. S. (2010). *Acta Cryst. E* **66**, o2847.
- Qin, L., Zhang, Q., Sun, W., Wang, J., Lu, C., Cheng, Y. & Wang, L. (2009). *Dalton Trans.* pp. 9388–9391.
- Rigaku (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1999). *NUMABS*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2010). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
- Salem, G. & Wild, S. B. (1992). *Inorg. Chem.* **31**, 581–586.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Sun, W.-H., Li, Z., Hu, H., Wu, B., Yang, H., Zhu, N., Leng, X. & Wang, H. (2002). *New J. Chem.* **26**, 1474–1478.
- Suzuki, T. (2004). *Bull. Chem. Soc. Jpn*, **77**, 1869–1876.
- Suzuki, T., Kotera, M., Takayama, A. & Kojima, M. (2009). *Polyhedron*, **28**, 2287–2293.
- Suzuki, T., Kuchiyama, T., Kishi, S., Kaizaki, S. & Kato, M. (2002). *Bull. Chem. Soc. Jpn*, **75**, 2433–2439.
- Suzuki, T., Yamaguchi, H., Hashimoto, A., Nozaki, K., Doi, M., Inazumi, N., Ikeda, N., Kawata, S., Kojima, M. & Takagi, H. D. (2011). *Inorg. Chem.* **50**, 3981–3987.
- Tsukuda, T., Nishigata, C., Arai, K. & Tsubomura, T. (2009). *Polyhedron*, **28**, 7–12.
- Wehman, P., van Donge, H. M. A., Hagos, A., Kamer, P. C. J. & van Leeuwen, P. W. N. M. (1997). *J. Organomet. Chem.* **535**, 183–193.

## supporting information

*Acta Cryst.* (2015). E71, 447-451 [https://doi.org/10.1107/S2056989015006076]

## Crystal structures of dichloridopalladium(II), -platinum(II) and -rhodium(III) complexes containing 8-(diphenylphosphanyl)quinoline

Takayoshi Suzuki, Hiroshi Yamaguchi, Masayuki Fujiki, Akira Hashimoto and Hideo D. Takagi

### Computing details

For all compounds, data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO* (Rigaku, 1998); data reduction: *CrystalStructure* (Rigaku, 2010). Program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008) for (1), (2); *DIRDIF99-PATY* (Beurskens *et al.*, 1999) for (3). For all compounds, program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2015).

### (1) Dichlorido(8-diphenylphosphanylquinoline)palladium(II)

#### Crystal data

[PdCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)]

$M_r = 490.62$

Monoclinic,  $P2_1/n$

$a = 9.0293$  (5) Å

$b = 15.2154$  (8) Å

$c = 13.7936$  (6) Å

$\beta = 91.8197$  (13)°

$V = 1894.07$  (16) Å<sup>3</sup>

$Z = 4$

$F(000) = 976$

$D_x = 1.721$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71075$  Å

Cell parameters from 10456 reflections

$\theta = 3.0$ – $27.5$ °

$\mu = 1.35$  mm<sup>-1</sup>

$T = 200$  K

Block, yellow

$0.18 \times 0.15 \times 0.12$  mm

#### Data collection

Rigaku R-AXIS RAPID

diffractometer

Radiation source: fine-focus sealed tube

Detector resolution: 10.00 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: numerical

(*NUMABS*; Rigaku, 1999)

$T_{\min} = 0.693$ ,  $T_{\max} = 0.850$

18103 measured reflections

4283 independent reflections

3913 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\max} = 27.5$ °,  $\theta_{\min} = 3.1$ °

$h = -11 \rightarrow 9$

$k = -19 \rightarrow 19$

$l = -17 \rightarrow 17$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.074$

$S = 0.92$

4283 reflections

235 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0416P)^2 + 2.3772P]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.50 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.42495 (2)	0.29307 (2)	0.78114 (2)	0.02590 (7)
Cl1	0.57514 (7)	0.16783 (4)	0.81289 (5)	0.04011 (15)
Cl2	0.39715 (8)	0.25626 (5)	0.62073 (5)	0.04049 (16)
P1	0.28188 (6)	0.40803 (4)	0.75258 (4)	0.02469 (13)
N1	0.4378 (2)	0.33998 (13)	0.92166 (15)	0.0296 (4)
C2	0.5290 (3)	0.30655 (17)	0.9896 (2)	0.0359 (6)
H2	0.5960	0.2618	0.9714	0.043*
C3	0.5315 (3)	0.33376 (19)	1.0863 (2)	0.0414 (6)
H3	0.5995	0.3081	1.1320	0.050*
C4	0.4363 (3)	0.39706 (19)	1.11443 (19)	0.0400 (6)
H4	0.4334	0.4139	1.1807	0.048*
C5	0.2453 (3)	0.50817 (18)	1.06576 (19)	0.0396 (6)
H5	0.2397	0.5283	1.1307	0.047*
C6	0.1606 (3)	0.54738 (18)	0.9943 (2)	0.0391 (6)
H6	0.0978	0.5950	1.0101	0.047*
C7	0.1650 (3)	0.51819 (17)	0.89779 (18)	0.0333 (5)
H7	0.1053	0.5460	0.8489	0.040*
C8	0.2560 (3)	0.44917 (15)	0.87408 (17)	0.0271 (5)
C9	0.3462 (3)	0.40817 (15)	0.94699 (17)	0.0280 (5)
C10	0.3411 (3)	0.43799 (17)	1.04423 (18)	0.0346 (5)
C11	0.0989 (3)	0.39004 (15)	0.70011 (17)	0.0279 (5)
C12	0.0674 (3)	0.40697 (17)	0.60209 (19)	0.0335 (5)
H12	0.1406	0.4325	0.5629	0.040*
C13	-0.0709 (3)	0.38636 (19)	0.5624 (2)	0.0402 (6)
H13	-0.0932	0.3987	0.4960	0.048*
C14	-0.1766 (3)	0.3481 (2)	0.6186 (2)	0.0440 (7)
H14	-0.2709	0.3336	0.5905	0.053*
C15	-0.1463 (3)	0.3307 (2)	0.7151 (2)	0.0432 (7)
H15	-0.2198	0.3042	0.7533	0.052*
C16	-0.0090 (3)	0.35166 (17)	0.75691 (19)	0.0341 (5)
H16	0.0114	0.3400	0.8237	0.041*
C17	0.3685 (3)	0.49474 (15)	0.68551 (16)	0.0260 (5)
C18	0.5132 (3)	0.48296 (17)	0.65615 (18)	0.0325 (5)
H18	0.5608	0.4277	0.6651	0.039*
C19	0.5880 (3)	0.55202 (19)	0.61371 (19)	0.0391 (6)
H19	0.6871	0.5441	0.5943	0.047*

C20	0.5192 (3)	0.63163 (19)	0.59974 (19)	0.0405 (6)
H20	0.5711	0.6788	0.5711	0.049*
C21	0.3749 (3)	0.64331 (18)	0.6273 (2)	0.0393 (6)
H21	0.3273	0.6983	0.6163	0.047*
C22	0.2985 (3)	0.57554 (17)	0.67083 (18)	0.0333 (5)
H22	0.1996	0.5841	0.6904	0.040*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pd1	0.02371 (11)	0.02217 (11)	0.03178 (12)	−0.00047 (6)	0.00035 (7)	0.00150 (6)
Cl1	0.0335 (3)	0.0306 (3)	0.0562 (4)	0.0070 (3)	0.0014 (3)	0.0060 (3)
Cl2	0.0487 (4)	0.0367 (3)	0.0361 (3)	0.0004 (3)	0.0012 (3)	−0.0069 (2)
P1	0.0232 (3)	0.0233 (3)	0.0276 (3)	−0.0006 (2)	0.0002 (2)	0.0007 (2)
N1	0.0268 (10)	0.0295 (11)	0.0323 (11)	−0.0045 (8)	−0.0015 (8)	0.0046 (8)
C2	0.0336 (14)	0.0307 (13)	0.0427 (15)	−0.0045 (10)	−0.0076 (11)	0.0079 (10)
C3	0.0452 (16)	0.0406 (15)	0.0376 (15)	−0.0086 (13)	−0.0108 (11)	0.0079 (11)
C4	0.0478 (17)	0.0442 (16)	0.0276 (13)	−0.0135 (13)	−0.0061 (11)	0.0016 (11)
C5	0.0490 (17)	0.0379 (14)	0.0322 (14)	−0.0090 (12)	0.0081 (11)	−0.0077 (10)
C6	0.0426 (16)	0.0317 (14)	0.0436 (16)	−0.0012 (11)	0.0109 (11)	−0.0056 (11)
C7	0.0337 (13)	0.0300 (13)	0.0366 (14)	−0.0012 (10)	0.0057 (10)	−0.0008 (10)
C8	0.0260 (12)	0.0269 (12)	0.0286 (12)	−0.0045 (9)	0.0020 (9)	0.0000 (9)
C9	0.0273 (12)	0.0252 (11)	0.0314 (12)	−0.0074 (9)	0.0006 (9)	0.0010 (9)
C10	0.0397 (15)	0.0321 (13)	0.0320 (13)	−0.0117 (11)	0.0001 (10)	0.0006 (10)
C11	0.0247 (12)	0.0237 (11)	0.0354 (13)	0.0009 (9)	−0.0006 (9)	−0.0019 (9)
C12	0.0334 (13)	0.0317 (13)	0.0351 (14)	−0.0009 (10)	−0.0032 (10)	0.0010 (10)
C13	0.0360 (15)	0.0385 (15)	0.0451 (16)	0.0083 (11)	−0.0136 (11)	−0.0063 (11)
C14	0.0280 (14)	0.0427 (16)	0.0607 (19)	0.0062 (12)	−0.0074 (12)	−0.0177 (13)
C15	0.0275 (14)	0.0441 (16)	0.0583 (19)	−0.0037 (12)	0.0083 (12)	−0.0111 (13)
C16	0.0296 (13)	0.0351 (13)	0.0375 (14)	−0.0036 (10)	0.0023 (10)	−0.0022 (10)
C17	0.0257 (12)	0.0273 (11)	0.0250 (11)	−0.0040 (9)	−0.0015 (8)	0.0006 (8)
C18	0.0302 (13)	0.0298 (13)	0.0375 (14)	0.0002 (10)	0.0030 (10)	−0.0015 (10)
C19	0.0343 (14)	0.0434 (15)	0.0400 (15)	−0.0089 (12)	0.0096 (11)	−0.0019 (11)
C20	0.0496 (17)	0.0358 (14)	0.0362 (14)	−0.0142 (12)	0.0028 (11)	0.0044 (11)
C21	0.0473 (16)	0.0287 (13)	0.0415 (15)	−0.0018 (11)	−0.0053 (11)	0.0064 (10)
C22	0.0295 (13)	0.0328 (13)	0.0374 (14)	0.0000 (10)	−0.0013 (10)	0.0033 (10)

*Geometric parameters (Å, °)*

Pd1—N1	2.065 (2)	C9—C10	1.418 (3)
Pd1—P1	2.2026 (6)	C11—C12	1.397 (4)
Pd1—Cl2	2.2885 (7)	C11—C16	1.397 (3)
Pd1—Cl1	2.3716 (6)	C12—C13	1.384 (4)
P1—C11	1.804 (2)	C12—H12	0.9500
P1—C17	1.804 (2)	C13—C14	1.377 (4)
P1—C8	1.811 (2)	C13—H13	0.9500
N1—C2	1.329 (3)	C14—C15	1.377 (4)
N1—C9	1.379 (3)	C14—H14	0.9500



C2—C3	1.396 (4)	C15—C16	1.387 (4)
C2—H2	0.9500	C15—H15	0.9500
C3—C4	1.356 (4)	C16—H16	0.9500
C3—H3	0.9500	C17—C22	1.394 (3)
C4—C10	1.418 (4)	C17—C18	1.392 (3)
C4—H4	0.9500	C18—C19	1.388 (4)
C5—C6	1.365 (4)	C18—H18	0.9500
C5—C10	1.412 (4)	C19—C20	1.372 (4)
C5—H5	0.9500	C19—H19	0.9500
C6—C7	1.405 (4)	C20—C21	1.381 (4)
C6—H6	0.9500	C20—H20	0.9500
C7—C8	1.379 (3)	C21—C22	1.388 (4)
C7—H7	0.9500	C21—H21	0.9500
C8—C9	1.418 (3)	C22—H22	0.9500
N1—Pd1—P1	84.75 (6)	C5—C10—C9	118.6 (2)
N1—Pd1—Cl2	173.28 (6)	C5—C10—C4	123.4 (2)
P1—Pd1—Cl2	88.61 (2)	C9—C10—C4	117.9 (3)
N1—Pd1—Cl1	95.14 (6)	C12—C11—C16	119.7 (2)
P1—Pd1—Cl1	178.94 (2)	C12—C11—P1	121.08 (19)
Cl2—Pd1—Cl1	91.51 (3)	C16—C11—P1	119.00 (19)
C11—P1—C17	108.18 (11)	C13—C12—C11	119.7 (3)
C11—P1—C8	106.28 (11)	C13—C12—H12	120.2
C17—P1—C8	107.01 (11)	C11—C12—H12	120.2
C11—P1—Pd1	118.41 (8)	C12—C13—C14	120.4 (3)
C17—P1—Pd1	114.27 (8)	C12—C13—H13	119.8
C8—P1—Pd1	101.60 (8)	C14—C13—H13	119.8
C2—N1—C9	118.2 (2)	C15—C14—C13	120.4 (3)
C2—N1—Pd1	122.99 (19)	C15—C14—H14	119.8
C9—N1—Pd1	118.79 (16)	C13—C14—H14	119.8
N1—C2—C3	123.4 (3)	C14—C15—C16	120.3 (3)
N1—C2—H2	118.3	C14—C15—H15	119.8
C3—C2—H2	118.3	C16—C15—H15	119.8
C4—C3—C2	119.5 (3)	C15—C16—C11	119.5 (3)
C4—C3—H3	120.2	C15—C16—H16	120.2
C2—C3—H3	120.2	C11—C16—H16	120.2
C3—C4—C10	119.5 (3)	C22—C17—C18	119.8 (2)
C3—C4—H4	120.3	C22—C17—P1	121.15 (18)
C10—C4—H4	120.3	C18—C17—P1	118.78 (19)
C6—C5—C10	120.8 (2)	C19—C18—C17	119.9 (2)
C6—C5—H5	119.6	C19—C18—H18	120.0
C10—C5—H5	119.6	C17—C18—H18	120.0
C5—C6—C7	120.9 (3)	C20—C19—C18	120.2 (3)
C5—C6—H6	119.5	C20—C19—H19	119.9
C7—C6—H6	119.5	C18—C19—H19	119.9
C8—C7—C6	120.0 (3)	C19—C20—C21	120.1 (2)
C8—C7—H7	120.0	C19—C20—H20	119.9
C6—C7—H7	120.0	C21—C20—H20	119.9

C7—C8—C9	120.0 (2)	C20—C21—C22	120.6 (3)
C7—C8—P1	125.31 (19)	C20—C21—H21	119.7
C9—C8—P1	114.59 (18)	C22—C21—H21	119.7
N1—C9—C8	119.1 (2)	C21—C22—C17	119.3 (2)
N1—C9—C10	121.2 (2)	C21—C22—H22	120.4
C8—C9—C10	119.7 (2)	C17—C22—H22	120.4
C9—N1—C2—C3	3.4 (4)	C3—C4—C10—C9	2.3 (4)
Pd1—N1—C2—C3	-175.5 (2)	C17—P1—C11—C12	28.7 (2)
N1—C2—C3—C4	0.5 (4)	C8—P1—C11—C12	143.3 (2)
C2—C3—C4—C10	-3.3 (4)	Pd1—P1—C11—C12	-103.4 (2)
C10—C5—C6—C7	0.9 (4)	C17—P1—C11—C16	-156.7 (2)
C5—C6—C7—C8	-0.2 (4)	C8—P1—C11—C16	-42.1 (2)
C6—C7—C8—C9	-0.6 (4)	Pd1—P1—C11—C16	71.2 (2)
C6—C7—C8—P1	-176.4 (2)	C16—C11—C12—C13	0.6 (4)
C11—P1—C8—C7	-50.7 (2)	P1—C11—C12—C13	175.2 (2)
C17—P1—C8—C7	64.7 (2)	C11—C12—C13—C14	-1.0 (4)
Pd1—P1—C8—C7	-175.1 (2)	C12—C13—C14—C15	0.6 (4)
C11—P1—C8—C9	133.35 (18)	C13—C14—C15—C16	0.1 (4)
C17—P1—C8—C9	-111.23 (18)	C14—C15—C16—C11	-0.5 (4)
Pd1—P1—C8—C9	8.88 (18)	C12—C11—C16—C15	0.1 (4)
C2—N1—C9—C8	175.3 (2)	P1—C11—C16—C15	-174.6 (2)
Pd1—N1—C9—C8	-5.7 (3)	C11—P1—C17—C22	52.8 (2)
C2—N1—C9—C10	-4.5 (3)	C8—P1—C17—C22	-61.3 (2)
Pd1—N1—C9—C10	174.54 (17)	Pd1—P1—C17—C22	-172.94 (17)
C7—C8—C9—N1	-179.2 (2)	C11—P1—C17—C18	-133.20 (19)
P1—C8—C9—N1	-3.0 (3)	C8—P1—C17—C18	112.7 (2)
C7—C8—C9—C10	0.6 (3)	Pd1—P1—C17—C18	1.0 (2)
P1—C8—C9—C10	176.80 (18)	C22—C17—C18—C19	1.0 (4)
C6—C5—C10—C9	-0.9 (4)	P1—C17—C18—C19	-173.1 (2)
C6—C5—C10—C4	177.3 (3)	C17—C18—C19—C20	-0.6 (4)
N1—C9—C10—C5	179.9 (2)	C18—C19—C20—C21	-0.4 (4)
C8—C9—C10—C5	0.1 (3)	C19—C20—C21—C22	1.2 (4)
N1—C9—C10—C4	1.7 (3)	C20—C21—C22—C17	-0.8 (4)
C8—C9—C10—C4	-178.1 (2)	C18—C17—C22—C21	-0.2 (4)
C3—C4—C10—C5	-175.9 (3)	P1—C17—C22—C21	173.7 (2)

## (2) Dichlorido(8-diphenylphosphanylquinoline)platinum(II) dichloromethane monosolvate

*Crystal data*[PtCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)]·CH<sub>2</sub>Cl<sub>2</sub>*M<sub>r</sub>* = 664.23Monoclinic, *P*2<sub>1</sub>/*c**a* = 13.9280 (5) Å*b* = 9.2371 (3) Å*c* = 17.8941 (6) Å $\beta$  = 102.8447 (10)°*V* = 2244.53 (13) Å<sup>3</sup>*Z* = 4*F*(000) = 1272*D<sub>x</sub>* = 1.966 Mg m<sup>-3</sup>Mo *K*α radiation,  $\lambda$  = 0.71075 Å

Cell parameters from 12710 reflections

 $\theta$  = 3.0–27.4° $\mu$  = 6.81 mm<sup>-1</sup>*T* = 200 K

Platelet, colorless

0.25 × 0.24 × 0.05 mm

*Data collection*

Rigaku R-AXIS RAPID  
diffractometer

Radiation source: fine-focus sealed tube

Detector resolution: 10.00 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: numerical  
(NUMABS; Rigaku, 1999)

$T_{\min} = 0.281$ ,  $T_{\max} = 0.727$

20995 measured reflections

5081 independent reflections

4665 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 27.4^\circ$ ,  $\theta_{\min} = 3.0^\circ$

$h = -18 \rightarrow 18$

$k = -11 \rightarrow 10$

$l = -23 \rightarrow 23$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.017$

$wR(F^2) = 0.037$

$S = 0.98$

5081 reflections

262 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0073P)^2 + 3.0468P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$

$\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.60 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pt1	0.34435 (2)	0.08493 (2)	0.69614 (2)	0.02160 (3)
Cl1	0.40053 (5)	-0.06049 (7)	0.80665 (3)	0.03408 (14)
Cl2	0.20025 (5)	0.12939 (8)	0.73477 (4)	0.03670 (15)
Cl3	0.19749 (8)	0.61854 (9)	0.43864 (4)	0.0575 (2)
Cl4	0.04531 (7)	0.67431 (12)	0.30251 (5)	0.0619 (2)
P1	0.29474 (5)	0.22285 (6)	0.59493 (3)	0.02219 (12)
N1	0.46929 (15)	0.0549 (2)	0.65523 (11)	0.0237 (4)
C1	0.1712 (2)	0.6794 (3)	0.34355 (16)	0.0393 (6)
H1A	0.1953	0.7800	0.3420	0.047*
H1B	0.2065	0.6182	0.3131	0.047*
C2	0.5401 (2)	-0.0380 (3)	0.68572 (14)	0.0313 (6)
H2	0.5313	-0.0960	0.7275	0.038*
C3	0.6271 (2)	-0.0547 (3)	0.65944 (16)	0.0367 (6)
H3	0.6758	-0.1220	0.6834	0.044*
C4	0.6411 (2)	0.0261 (3)	0.59951 (15)	0.0342 (6)
H4	0.7001	0.0164	0.5816	0.041*
C5	0.5755 (2)	0.2104 (3)	0.49983 (15)	0.0352 (6)
H5	0.6324	0.2022	0.4791	0.042*
C6	0.5028 (2)	0.3044 (3)	0.46766 (15)	0.0374 (6)
H6	0.5101	0.3626	0.4255	0.045*

C7	0.4172 (2)	0.3161 (3)	0.49634 (14)	0.0319 (6)
H7	0.3670	0.3821	0.4733	0.038*
C8	0.40500 (18)	0.2329 (3)	0.55748 (13)	0.0241 (5)
C9	0.48110 (18)	0.1374 (3)	0.59276 (13)	0.0239 (5)
C10	0.5677 (2)	0.1246 (3)	0.56378 (14)	0.0283 (5)
C11	0.2562 (2)	0.4050 (3)	0.61167 (13)	0.0274 (5)
C12	0.1573 (2)	0.4345 (3)	0.60854 (17)	0.0403 (7)
H12	0.1089	0.3618	0.5922	0.048*
C13	0.1296 (3)	0.5704 (3)	0.6294 (2)	0.0524 (9)
H13	0.0622	0.5909	0.6272	0.063*
C14	0.2001 (3)	0.6758 (3)	0.65334 (17)	0.0481 (8)
H14	0.1808	0.7685	0.6678	0.058*
C15	0.2971 (3)	0.6479 (3)	0.65643 (16)	0.0440 (8)
H15	0.3449	0.7213	0.6728	0.053*
C16	0.3265 (2)	0.5121 (3)	0.63574 (13)	0.0318 (6)
H16	0.3941	0.4929	0.6381	0.038*
C17	0.20016 (18)	0.1489 (3)	0.51834 (13)	0.0249 (5)
C18	0.1620 (2)	0.0112 (3)	0.52486 (15)	0.0305 (6)
H18	0.1829	-0.0425	0.5709	0.037*
C19	0.0931 (2)	-0.0475 (3)	0.46383 (17)	0.0386 (6)
H19	0.0670	-0.1413	0.4685	0.046*
C20	0.0623 (2)	0.0296 (3)	0.39660 (16)	0.0420 (7)
H20	0.0150	-0.0107	0.3553	0.050*
C21	0.1009 (2)	0.1660 (4)	0.38972 (16)	0.0450 (7)
H21	0.0807	0.2186	0.3433	0.054*
C22	0.1687 (2)	0.2259 (3)	0.45004 (15)	0.0383 (6)
H22	0.1941	0.3200	0.4451	0.046*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pt1	0.02203 (5)	0.02250 (5)	0.02056 (5)	-0.00063 (4)	0.00538 (3)	-0.00020 (3)
Cl1	0.0310 (4)	0.0435 (4)	0.0267 (3)	0.0012 (3)	0.0042 (2)	0.0108 (3)
Cl2	0.0304 (4)	0.0427 (4)	0.0423 (3)	0.0051 (3)	0.0193 (3)	0.0086 (3)
Cl3	0.0775 (7)	0.0463 (4)	0.0421 (4)	0.0134 (4)	-0.0009 (4)	0.0046 (3)
Cl4	0.0372 (5)	0.0920 (7)	0.0503 (4)	0.0008 (4)	-0.0034 (4)	-0.0148 (4)
P1	0.0220 (3)	0.0218 (3)	0.0232 (3)	0.0003 (2)	0.0059 (2)	0.0005 (2)
N1	0.0226 (11)	0.0256 (10)	0.0228 (9)	0.0005 (8)	0.0051 (8)	-0.0043 (8)
C1	0.0342 (17)	0.0445 (16)	0.0389 (14)	-0.0001 (13)	0.0073 (12)	-0.0018 (13)
C2	0.0304 (15)	0.0352 (14)	0.0272 (12)	0.0058 (11)	0.0044 (11)	-0.0013 (11)
C3	0.0298 (15)	0.0410 (15)	0.0380 (14)	0.0096 (12)	0.0046 (12)	-0.0075 (12)
C4	0.0260 (15)	0.0414 (15)	0.0365 (14)	-0.0005 (12)	0.0094 (11)	-0.0142 (12)
C5	0.0317 (16)	0.0429 (16)	0.0355 (13)	-0.0102 (12)	0.0173 (12)	-0.0097 (12)
C6	0.0428 (18)	0.0401 (15)	0.0333 (13)	-0.0084 (13)	0.0171 (12)	0.0016 (12)
C7	0.0351 (16)	0.0312 (13)	0.0299 (12)	-0.0027 (11)	0.0080 (11)	0.0010 (11)
C8	0.0228 (13)	0.0256 (12)	0.0245 (11)	-0.0032 (10)	0.0063 (9)	-0.0048 (9)
C9	0.0257 (13)	0.0250 (11)	0.0212 (11)	-0.0051 (10)	0.0057 (9)	-0.0074 (9)
C10	0.0255 (14)	0.0306 (13)	0.0297 (12)	-0.0058 (10)	0.0078 (10)	-0.0104 (10)

C11	0.0363 (15)	0.0233 (12)	0.0242 (11)	0.0019 (11)	0.0104 (10)	0.0017 (9)
C12	0.0383 (18)	0.0309 (14)	0.0558 (18)	0.0018 (12)	0.0191 (14)	-0.0023 (13)
C13	0.060 (2)	0.0383 (17)	0.068 (2)	0.0172 (16)	0.0349 (19)	0.0037 (15)
C14	0.085 (3)	0.0241 (14)	0.0436 (16)	0.0096 (15)	0.0329 (17)	0.0021 (12)
C15	0.076 (3)	0.0254 (13)	0.0333 (14)	-0.0086 (15)	0.0173 (15)	-0.0041 (11)
C16	0.0416 (17)	0.0291 (13)	0.0251 (12)	-0.0041 (12)	0.0086 (11)	0.0010 (10)
C17	0.0236 (13)	0.0237 (11)	0.0273 (11)	0.0027 (10)	0.0052 (10)	-0.0020 (9)
C18	0.0283 (15)	0.0304 (13)	0.0329 (13)	-0.0022 (11)	0.0072 (11)	-0.0004 (11)
C19	0.0331 (16)	0.0346 (14)	0.0483 (16)	-0.0072 (12)	0.0091 (13)	-0.0118 (13)
C20	0.0305 (16)	0.0529 (18)	0.0391 (15)	-0.0009 (14)	0.0002 (12)	-0.0158 (14)
C21	0.0432 (19)	0.0534 (19)	0.0322 (14)	0.0019 (15)	-0.0052 (13)	0.0027 (13)
C22	0.0399 (17)	0.0322 (14)	0.0375 (14)	-0.0011 (12)	-0.0026 (12)	0.0057 (12)

*Geometric parameters (Å, °)*

Pt1—N1	2.051 (2)	C7—H7	0.9500
Pt1—P1	2.1963 (6)	C8—C9	1.416 (3)
Pt1—C12	2.3002 (7)	C9—C10	1.420 (3)
Pt1—C11	2.3747 (6)	C11—C16	1.392 (4)
C13—C1	1.752 (3)	C11—C12	1.393 (4)
C14—C1	1.744 (3)	C12—C13	1.388 (4)
P1—C17	1.810 (3)	C12—H12	0.9500
P1—C8	1.809 (2)	C13—C14	1.381 (5)
P1—C11	1.811 (2)	C13—H13	0.9500
N1—C2	1.329 (3)	C14—C15	1.366 (5)
N1—C9	1.392 (3)	C14—H14	0.9500
C1—H1A	0.9900	C15—C16	1.394 (4)
C1—H1B	0.9900	C15—H15	0.9500
C2—C3	1.402 (4)	C16—H16	0.9500
C2—H2	0.9500	C17—C18	1.393 (4)
C3—C4	1.356 (4)	C17—C22	1.398 (3)
C3—H3	0.9500	C18—C19	1.393 (4)
C4—C10	1.411 (4)	C18—H18	0.9500
C4—H4	0.9500	C19—C20	1.381 (4)
C5—C6	1.361 (4)	C19—H19	0.9500
C5—C10	1.416 (4)	C20—C21	1.386 (4)
C5—H5	0.9500	C20—H20	0.9500
C6—C7	1.404 (4)	C21—C22	1.382 (4)
C6—H6	0.9500	C21—H21	0.9500
C7—C8	1.378 (3)	C22—H22	0.9500
N1—Pt1—P1	85.44 (6)	N1—C9—C8	119.3 (2)
N1—Pt1—C12	175.93 (6)	N1—C9—C10	120.6 (2)
P1—Pt1—C12	90.50 (2)	C8—C9—C10	120.1 (2)
N1—Pt1—C11	94.13 (6)	C4—C10—C9	118.5 (2)
P1—Pt1—C11	178.80 (2)	C4—C10—C5	123.3 (2)
C12—Pt1—C11	89.93 (2)	C9—C10—C5	118.2 (2)
C17—P1—C8	105.88 (11)	C16—C11—C12	119.6 (2)

C17—P1—C11	106.45 (12)	C16—C11—P1	119.9 (2)
C8—P1—C11	108.71 (12)	C12—C11—P1	120.1 (2)
C17—P1—Pt1	116.61 (8)	C13—C12—C11	119.8 (3)
C8—P1—Pt1	101.42 (8)	C13—C12—H12	120.1
C11—P1—Pt1	116.93 (8)	C11—C12—H12	120.1
C2—N1—C9	118.2 (2)	C14—C13—C12	120.1 (3)
C2—N1—Pt1	123.48 (17)	C14—C13—H13	120.0
C9—N1—Pt1	118.27 (16)	C12—C13—H13	120.0
C14—C1—C13	111.96 (17)	C15—C14—C13	120.5 (3)
C14—C1—H1A	109.2	C15—C14—H14	119.7
C13—C1—H1A	109.2	C13—C14—H14	119.7
C14—C1—H1B	109.2	C14—C15—C16	120.3 (3)
C13—C1—H1B	109.2	C14—C15—H15	119.8
H1A—C1—H1B	107.9	C16—C15—H15	119.8
N1—C2—C3	123.4 (2)	C11—C16—C15	119.6 (3)
N1—C2—H2	118.3	C11—C16—H16	120.2
C3—C2—H2	118.3	C15—C16—H16	120.2
C4—C3—C2	119.5 (3)	C18—C17—C22	119.1 (2)
C4—C3—H3	120.3	C18—C17—P1	120.54 (19)
C2—C3—H3	120.3	C22—C17—P1	120.2 (2)
C3—C4—C10	119.7 (3)	C17—C18—C19	119.9 (2)
C3—C4—H4	120.1	C17—C18—H18	120.0
C10—C4—H4	120.1	C19—C18—H18	120.0
C6—C5—C10	121.0 (3)	C20—C19—C18	120.5 (3)
C6—C5—H5	119.5	C20—C19—H19	119.7
C10—C5—H5	119.5	C18—C19—H19	119.7
C5—C6—C7	120.5 (3)	C19—C20—C21	119.6 (3)
C5—C6—H6	119.8	C19—C20—H20	120.2
C7—C6—H6	119.8	C21—C20—H20	120.2
C8—C7—C6	120.8 (3)	C22—C21—C20	120.4 (3)
C8—C7—H7	119.6	C22—C21—H21	119.8
C6—C7—H7	119.6	C20—C21—H21	119.8
C7—C8—C9	119.3 (2)	C21—C22—C17	120.4 (3)
C7—C8—P1	126.0 (2)	C21—C22—H22	119.8
C9—C8—P1	114.57 (17)	C17—C22—H22	119.8
C9—N1—C2—C3	1.6 (4)	C6—C5—C10—C9	1.0 (4)
Pt1—N1—C2—C3	-177.6 (2)	C17—P1—C11—C16	-147.98 (19)
N1—C2—C3—C4	-0.5 (4)	C8—P1—C11—C16	-34.3 (2)
C2—C3—C4—C10	-0.7 (4)	Pt1—P1—C11—C16	79.7 (2)
C10—C5—C6—C7	-1.4 (4)	C17—P1—C11—C12	38.9 (2)
C5—C6—C7—C8	0.0 (4)	C8—P1—C11—C12	152.6 (2)
C6—C7—C8—C9	1.6 (4)	Pt1—P1—C11—C12	-93.4 (2)
C6—C7—C8—P1	-173.7 (2)	C16—C11—C12—C13	0.0 (4)
C17—P1—C8—C7	62.9 (2)	P1—C11—C12—C13	173.1 (2)
C11—P1—C8—C7	-51.1 (2)	C11—C12—C13—C14	-0.1 (5)
Pt1—P1—C8—C7	-174.9 (2)	C12—C13—C14—C15	0.3 (5)
C17—P1—C8—C9	-112.53 (18)	C13—C14—C15—C16	-0.3 (4)

C11—P1—C8—C9	133.44 (18)	C12—C11—C16—C15	0.0 (4)
Pt1—P1—C8—C9	9.65 (18)	P1—C11—C16—C15	-173.17 (19)
C2—N1—C9—C8	177.6 (2)	C14—C15—C16—C11	0.2 (4)
Pt1—N1—C9—C8	-3.1 (3)	C8—P1—C17—C18	112.0 (2)
C2—N1—C9—C10	-1.5 (3)	C11—P1—C17—C18	-132.4 (2)
Pt1—N1—C9—C10	177.82 (17)	Pt1—P1—C17—C18	0.1 (2)
C7—C8—C9—N1	179.1 (2)	C8—P1—C17—C22	-64.2 (2)
P1—C8—C9—N1	-5.1 (3)	C11—P1—C17—C22	51.3 (2)
C7—C8—C9—C10	-1.9 (3)	Pt1—P1—C17—C22	-176.13 (19)
P1—C8—C9—C10	173.94 (17)	C22—C17—C18—C19	-0.3 (4)
C3—C4—C10—C9	0.8 (4)	P1—C17—C18—C19	-176.5 (2)
C3—C4—C10—C5	-178.5 (2)	C17—C18—C19—C20	0.2 (4)
N1—C9—C10—C4	0.3 (3)	C18—C19—C20—C21	0.4 (5)
C8—C9—C10—C4	-178.7 (2)	C19—C20—C21—C22	-1.0 (5)
N1—C9—C10—C5	179.6 (2)	C20—C21—C22—C17	0.9 (5)
C8—C9—C10—C5	0.6 (3)	C18—C17—C22—C21	-0.2 (4)
C6—C5—C10—C4	-179.7 (3)	P1—C17—C22—C21	176.0 (2)

(3) *cis*-Dichloridobis(8-diphenylphosphanylquinoline)rhodium(III) hexafluoridophosphate  
dichloromethane/methanol hemisolvate

*Crystal data*

[RhCl<sub>2</sub>(C<sub>21</sub>H<sub>16</sub>NP)<sub>2</sub>](PF<sub>6</sub>)·0.5CH<sub>2</sub>Cl<sub>2</sub>·0.5CH<sub>4</sub>O

*M<sub>r</sub>* = 1003.90

Triclinic, *P*1̄

*a* = 9.841 (5) Å

*b* = 13.825 (6) Å

*c* = 16.167 (8) Å

*α* = 87.307 (19)°

*β* = 81.80 (2)°

*γ* = 70.819 (18)°

*V* = 2056.2 (17) Å<sup>3</sup>

*Z* = 2

*F*(000) = 1012

*D<sub>x</sub>* = 1.621 Mg m<sup>-3</sup>

Mo *Kα* radiation, *λ* = 0.71075 Å

Cell parameters from 15480 reflections

*θ* = 3.1–27.5°

*μ* = 0.79 mm<sup>-1</sup>

*T* = 200 K

Prism, colorless

0.20 × 0.20 × 0.15 mm

*Data collection*

Rigaku R-Axis RAPID

diffractometer

Radiation source: fine-focus sealed tube

Detector resolution: 10.00 pixels mm<sup>-1</sup>

*ω* scans

Absorption correction: multi-scan

(*ABSCOR*.; Rigaku, 1995)

*T<sub>min</sub>* = 0.848, *T<sub>max</sub>* = 0.882

20385 measured reflections

9338 independent reflections

7245 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.046

*θ<sub>max</sub>* = 27.5°, *θ<sub>min</sub>* = 3.1°

*h* = -12→12

*k* = -17→17

*l* = -20→20

*Refinement*

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.044

*wR*(*F*<sup>2</sup>) = 0.120

*S* = 1.06

9338 reflections

605 parameters

20 restraints

Primary atom site location: heavy-atom method

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

*w* = 1/[σ<sup>2</sup>(*F<sub>o</sub>*<sup>2</sup>) + (0.064*P*)<sup>2</sup>]

where *P* = (*F<sub>o</sub>*<sup>2</sup> + 2*F<sub>c</sub>*<sup>2</sup>)/3

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.68 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.03 \text{ e } \text{\AA}^{-3}$$

### Special details

**Experimental.** The  $^{31}\text{P}$  NMR spectrum of 3 in  $\text{CD}_3\text{CN}$  (400 MHz, 22 °C) showed two doublet of doublets resonances at  $\delta$  38.66 ( $J_{\text{Rh-P}} = 110$ ,  $J_{\text{P-P}} = 21$  Hz) and 40.47 ( $J_{\text{Rh-P}} = 113$  Hz).

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Rh1	0.81791 (2)	0.88958 (2)	0.68292 (2)	0.02967 (9)	
Cl1	0.92494 (8)	0.90055 (6)	0.54278 (4)	0.03774 (17)	
Cl2	1.04776 (8)	0.82096 (6)	0.72496 (4)	0.03852 (17)	
P1	0.71063 (8)	0.90351 (6)	0.81912 (4)	0.03163 (17)	
P2	0.80756 (8)	0.73608 (6)	0.64925 (4)	0.03480 (18)	
P3	0.55681 (15)	0.62928 (12)	0.29883 (9)	0.0909 (4)	
F1A	0.6674 (12)	0.5251 (7)	0.3142 (9)	0.163 (6)	0.613 (14)
F2A	0.4337 (9)	0.5838 (8)	0.3072 (9)	0.132 (4)	0.613 (14)
F3A	0.601 (2)	0.6117 (9)	0.2058 (4)	0.182 (6)	0.613 (14)
F4A	0.559 (3)	0.633 (2)	0.3936 (6)	0.322 (12)	0.613 (14)
F5A	0.6781 (9)	0.6811 (7)	0.2764 (13)	0.174 (5)	0.613 (14)
F6A	0.4534 (11)	0.7384 (6)	0.2966 (15)	0.211 (8)	0.613 (14)
F1B	0.530 (4)	0.5418 (19)	0.356 (2)	0.300 (16)	0.387 (14)
F2B	0.451 (2)	0.6177 (18)	0.240 (2)	0.208 (10)	0.387 (14)
F3B	0.6695 (18)	0.5392 (19)	0.249 (2)	0.212 (13)	0.387 (14)
F4B	0.449 (2)	0.6981 (12)	0.3668 (10)	0.147 (7)	0.387 (14)
F5B	0.6769 (14)	0.6458 (17)	0.3408 (15)	0.163 (9)	0.387 (14)
F6B	0.522 (3)	0.7243 (11)	0.2443 (9)	0.232 (15)	0.387 (14)
N1	0.8319 (3)	1.03784 (18)	0.71011 (14)	0.0333 (5)	
N11	0.6185 (3)	0.9470 (2)	0.64031 (13)	0.0382 (6)	
Cl3A	0.3228 (12)	0.6504 (8)	0.8186 (7)	0.322 (5)	0.5
Cl4A	0.3702 (13)	0.5700 (9)	0.9744 (6)	0.407 (8)	0.5
C1A	0.289 (2)	0.5586 (14)	0.8897 (10)	0.307 (11)	0.5
H1A	0.1829	0.5731	0.9058	0.369*	0.5
H1B	0.3314	0.4890	0.8655	0.369*	0.5
O1B	0.3204 (14)	0.7010 (12)	0.8793 (10)	0.213 (8)	0.5
H1F	0.3593	0.7057	0.8302	0.256*	0.5
C1B	0.326 (2)	0.5987 (17)	0.8956 (10)	0.227 (11)	0.5
H1C	0.2739	0.5942	0.9514	0.341*	0.5
H1E	0.4273	0.5547	0.8932	0.341*	0.5
H1D	0.2801	0.5761	0.8537	0.341*	0.5
C2	0.8789 (4)	1.0945 (2)	0.6523 (2)	0.0478 (8)	
H2	0.8939	1.0739	0.5956	0.057*	
C3	0.9078 (4)	1.1828 (3)	0.6702 (2)	0.0567 (9)	
H3	0.9431	1.2203	0.6264	0.068*	



C4	0.8852 (4)	1.2148 (3)	0.7497 (2)	0.0571 (10)
H4	0.9020	1.2760	0.7624	0.069*
C5	0.8172 (5)	1.1806 (3)	0.8995 (2)	0.0696 (12)
H5	0.8330	1.2406	0.9159	0.083*
C6	0.7766 (5)	1.1198 (4)	0.9585 (2)	0.0743 (13)
H6	0.7667	1.1368	1.0157	0.089*
C7	0.7488 (4)	1.0322 (3)	0.93688 (19)	0.0542 (9)
H7	0.7205	0.9901	0.9794	0.065*
C8	0.7622 (3)	1.0064 (2)	0.85439 (17)	0.0377 (7)
C9	0.8102 (3)	1.0674 (2)	0.79179 (18)	0.0369 (6)
C10	0.8366 (4)	1.1571 (3)	0.8139 (2)	0.0484 (8)
C12	0.5505 (4)	1.0465 (3)	0.63529 (19)	0.0525 (9)
H12	0.5979	1.0922	0.6490	0.063*
C13	0.4129 (4)	1.0875 (4)	0.6109 (2)	0.0713 (13)
H13	0.3684	1.1596	0.6081	0.086*
C14	0.3440 (4)	1.0250 (4)	0.5916 (2)	0.0761 (15)
H14	0.2496	1.0525	0.5755	0.091*
C15	0.3453 (5)	0.8472 (6)	0.5772 (3)	0.0870 (18)
H15	0.2503	0.8714	0.5618	0.104*
C16	0.4156 (6)	0.7439 (6)	0.5816 (3)	0.098 (2)
H16	0.3687	0.6971	0.5697	0.118*
C17	0.5564 (4)	0.7070 (4)	0.6036 (3)	0.0723 (12)
H17	0.6048	0.6353	0.6055	0.087*
C18	0.6254 (4)	0.7737 (3)	0.62245 (19)	0.0495 (9)
C19	0.5529 (3)	0.8803 (3)	0.61891 (17)	0.0467 (8)
C20	0.4116 (4)	0.9172 (4)	0.59511 (19)	0.0615 (11)
C21	0.5141 (3)	0.9536 (3)	0.82721 (17)	0.0432 (8)
C22	0.4385 (4)	1.0554 (3)	0.8417 (2)	0.0644 (11)
H22	0.4881	1.1015	0.8512	0.077*
C23	0.2882 (6)	1.0909 (5)	0.8423 (3)	0.107 (2)
H23	0.2347	1.1613	0.8522	0.129*
C24	0.2165 (6)	1.0214 (8)	0.8282 (3)	0.125 (3)
H24	0.1141	1.0447	0.8291	0.151*
C25	0.2927 (6)	0.9224 (7)	0.8134 (3)	0.104 (2)
H25	0.2442	0.8760	0.8030	0.125*
C26	0.4391 (4)	0.8879 (4)	0.8133 (2)	0.0639 (11)
H26	0.4911	0.8174	0.8035	0.077*
C27	0.7478 (4)	0.8059 (2)	0.89991 (17)	0.0428 (7)
C28	0.6348 (5)	0.7924 (3)	0.9570 (2)	0.0645 (11)
H28	0.5373	0.8325	0.9520	0.077*
C29	0.6633 (6)	0.7216 (4)	1.0204 (3)	0.0891 (16)
H29	0.5859	0.7139	1.0596	0.107*
C30	0.8036 (7)	0.6625 (4)	1.0267 (3)	0.1008 (19)
H30	0.8233	0.6127	1.0699	0.121*
C31	0.9143 (6)	0.6745 (4)	0.9719 (3)	0.102 (2)
H31	1.0112	0.6331	0.9769	0.122*
C32	0.8874 (5)	0.7475 (4)	0.9078 (2)	0.0761 (14)
H32	0.9659	0.7562	0.8701	0.091*

C33	0.9307 (3)	0.6735 (2)	0.55799 (18)	0.0373 (7)
C34	1.0796 (3)	0.6486 (2)	0.55612 (19)	0.0437 (7)
H34	1.1176	0.6639	0.6028	0.052*
C35	1.1731 (4)	0.6015 (3)	0.4863 (2)	0.0505 (8)
H35	1.2750	0.5846	0.4852	0.061*
C36	1.1187 (4)	0.5794 (2)	0.4188 (2)	0.0487 (8)
H36	1.1828	0.5464	0.3713	0.058*
C37	0.9725 (4)	0.6047 (2)	0.41995 (19)	0.0481 (8)
H37	0.9356	0.5894	0.3728	0.058*
C38	0.8767 (4)	0.6523 (2)	0.48863 (18)	0.0427 (7)
H38	0.7750	0.6703	0.4883	0.051*
C39	0.8269 (4)	0.6356 (2)	0.72758 (18)	0.0407 (7)
C40	0.7092 (5)	0.6125 (3)	0.7699 (3)	0.0689 (12)
H40	0.6136	0.6507	0.7595	0.083*
C41	0.7312 (6)	0.5334 (3)	0.8277 (3)	0.0844 (15)
H41	0.6498	0.5179	0.8565	0.101*
C42	0.8658 (6)	0.4779 (3)	0.8439 (2)	0.0714 (12)
H42	0.8781	0.4249	0.8845	0.086*
C43	0.9853 (6)	0.4982 (3)	0.8016 (2)	0.0724 (12)
H43	1.0802	0.4584	0.8121	0.087*
C44	0.9657 (4)	0.5771 (3)	0.7437 (2)	0.0597 (10)
H44	1.0479	0.5915	0.7147	0.072*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Rh1	0.03129 (14)	0.03778 (14)	0.02306 (12)	-0.01406 (10)	-0.00624 (8)	-0.00226 (9)
Cl1	0.0437 (4)	0.0465 (4)	0.0252 (3)	-0.0185 (3)	-0.0016 (3)	-0.0013 (3)
Cl2	0.0332 (4)	0.0482 (4)	0.0393 (4)	-0.0168 (3)	-0.0119 (3)	-0.0035 (3)
P1	0.0335 (4)	0.0417 (4)	0.0227 (3)	-0.0152 (3)	-0.0058 (3)	-0.0020 (3)
P2	0.0358 (4)	0.0452 (4)	0.0305 (4)	-0.0218 (4)	-0.0038 (3)	-0.0078 (3)
P3	0.0619 (8)	0.1049 (11)	0.0944 (10)	-0.0099 (8)	-0.0086 (7)	-0.0199 (9)
F1A	0.099 (7)	0.145 (8)	0.206 (11)	0.007 (6)	-0.030 (8)	0.076 (9)
F2A	0.072 (4)	0.155 (8)	0.176 (9)	-0.058 (5)	0.023 (5)	-0.025 (7)
F3A	0.281 (16)	0.160 (11)	0.067 (4)	-0.040 (10)	0.018 (6)	0.022 (5)
F4A	0.30 (2)	0.51 (4)	0.108 (8)	-0.04 (3)	-0.068 (11)	-0.103 (13)
F5A	0.106 (7)	0.117 (6)	0.290 (16)	-0.033 (5)	-0.008 (8)	0.011 (9)
F6A	0.088 (6)	0.112 (7)	0.42 (2)	0.032 (5)	-0.092 (10)	-0.112 (12)
F1B	0.28 (4)	0.20 (2)	0.42 (4)	-0.11 (2)	-0.03 (4)	0.16 (3)
F2B	0.164 (18)	0.21 (2)	0.26 (2)	-0.038 (13)	-0.078 (18)	-0.12 (2)
F3B	0.115 (14)	0.20 (2)	0.30 (3)	-0.040 (14)	0.08 (2)	-0.14 (2)
F4B	0.140 (13)	0.143 (12)	0.121 (11)	-0.015 (9)	0.044 (9)	-0.052 (8)
F5B	0.079 (9)	0.22 (2)	0.184 (17)	-0.009 (10)	-0.078 (11)	-0.073 (14)
F6B	0.43 (4)	0.145 (17)	0.110 (10)	-0.11 (2)	-0.001 (15)	0.079 (12)
N1	0.0396 (14)	0.0320 (12)	0.0291 (11)	-0.0126 (11)	-0.0045 (10)	-0.0033 (10)
N11	0.0332 (14)	0.0595 (16)	0.0202 (11)	-0.0122 (12)	-0.0057 (9)	0.0011 (11)
Cl3A	0.302 (11)	0.251 (10)	0.445 (14)	-0.111 (8)	-0.131 (12)	0.083 (10)
Cl4A	0.480 (19)	0.444 (17)	0.221 (8)	-0.130 (12)	0.146 (10)	0.022 (9)

---

C1A	0.35 (3)	0.218 (19)	0.33 (2)	-0.15 (2)	0.233 (17)	-0.174 (15)
O1B	0.118 (9)	0.287 (16)	0.257 (16)	-0.110 (10)	0.061 (10)	-0.168 (13)
C1B	0.26 (2)	0.48 (3)	0.114 (11)	-0.33 (2)	-0.086 (12)	0.119 (16)
C2	0.067 (2)	0.0415 (17)	0.0344 (15)	-0.0181 (16)	-0.0036 (15)	0.0000 (14)
C3	0.080 (3)	0.0396 (18)	0.052 (2)	-0.0268 (18)	0.0016 (18)	0.0005 (16)
C4	0.075 (3)	0.0420 (18)	0.060 (2)	-0.0295 (19)	-0.0005 (19)	-0.0092 (17)
C5	0.100 (3)	0.072 (3)	0.053 (2)	-0.054 (3)	0.008 (2)	-0.027 (2)
C6	0.111 (4)	0.093 (3)	0.0392 (19)	-0.064 (3)	0.006 (2)	-0.026 (2)
C7	0.070 (2)	0.071 (2)	0.0324 (15)	-0.040 (2)	0.0014 (15)	-0.0128 (16)
C8	0.0358 (16)	0.0489 (17)	0.0311 (14)	-0.0173 (14)	-0.0017 (12)	-0.0085 (13)
C9	0.0331 (16)	0.0410 (16)	0.0358 (15)	-0.0096 (13)	-0.0041 (12)	-0.0090 (13)
C10	0.056 (2)	0.0469 (19)	0.0454 (18)	-0.0220 (17)	0.0004 (15)	-0.0139 (15)
C12	0.047 (2)	0.067 (2)	0.0326 (16)	-0.0030 (17)	-0.0095 (14)	0.0074 (16)
C13	0.046 (2)	0.100 (3)	0.044 (2)	0.009 (2)	-0.0103 (17)	0.015 (2)
C14	0.034 (2)	0.142 (5)	0.0343 (18)	-0.003 (3)	-0.0126 (15)	0.010 (2)
C15	0.038 (2)	0.180 (6)	0.051 (2)	-0.044 (3)	-0.0051 (18)	-0.029 (3)
C16	0.064 (3)	0.184 (6)	0.077 (3)	-0.079 (4)	-0.001 (2)	-0.049 (4)
C17	0.054 (2)	0.106 (3)	0.075 (3)	-0.048 (2)	0.000 (2)	-0.036 (2)
C18	0.0394 (19)	0.082 (3)	0.0365 (16)	-0.0320 (18)	-0.0013 (13)	-0.0200 (16)
C19	0.0312 (17)	0.087 (3)	0.0233 (13)	-0.0201 (17)	-0.0037 (11)	-0.0091 (15)
C20	0.0347 (19)	0.123 (4)	0.0258 (15)	-0.023 (2)	-0.0073 (13)	-0.0020 (19)
C21	0.0369 (17)	0.071 (2)	0.0236 (13)	-0.0205 (16)	-0.0054 (12)	0.0023 (14)
C22	0.046 (2)	0.084 (3)	0.051 (2)	-0.005 (2)	-0.0067 (17)	0.006 (2)
C23	0.058 (3)	0.150 (6)	0.069 (3)	0.024 (3)	-0.003 (2)	0.012 (3)
C24	0.034 (3)	0.271 (10)	0.056 (3)	-0.029 (4)	-0.011 (2)	0.009 (4)
C25	0.050 (3)	0.241 (8)	0.044 (2)	-0.075 (4)	-0.003 (2)	-0.022 (3)
C26	0.051 (2)	0.120 (4)	0.0341 (16)	-0.047 (2)	-0.0011 (15)	-0.0148 (19)
C27	0.059 (2)	0.0453 (17)	0.0256 (13)	-0.0170 (16)	-0.0089 (13)	-0.0009 (13)
C28	0.075 (3)	0.074 (3)	0.0450 (19)	-0.028 (2)	-0.0082 (18)	0.0157 (19)
C29	0.124 (5)	0.091 (4)	0.048 (2)	-0.037 (3)	-0.001 (3)	0.025 (2)
C30	0.149 (5)	0.081 (3)	0.047 (2)	-0.008 (3)	-0.005 (3)	0.021 (2)
C31	0.099 (4)	0.110 (4)	0.051 (2)	0.026 (3)	-0.012 (3)	0.017 (3)
C32	0.068 (3)	0.101 (3)	0.0350 (18)	0.003 (2)	-0.0040 (18)	0.014 (2)
C33	0.0431 (18)	0.0379 (15)	0.0359 (15)	-0.0207 (14)	-0.0012 (13)	-0.0082 (13)
C34	0.0431 (19)	0.0511 (19)	0.0405 (16)	-0.0209 (15)	0.0000 (13)	-0.0132 (14)
C35	0.050 (2)	0.0480 (19)	0.054 (2)	-0.0198 (16)	0.0054 (16)	-0.0127 (16)
C36	0.068 (2)	0.0366 (16)	0.0392 (16)	-0.0194 (16)	0.0109 (16)	-0.0116 (14)
C37	0.072 (3)	0.0428 (17)	0.0344 (15)	-0.0260 (17)	-0.0039 (15)	-0.0079 (14)
C38	0.055 (2)	0.0463 (17)	0.0341 (15)	-0.0263 (16)	-0.0058 (14)	-0.0066 (13)
C39	0.054 (2)	0.0420 (17)	0.0341 (15)	-0.0283 (15)	0.0012 (13)	-0.0080 (13)
C40	0.066 (3)	0.052 (2)	0.083 (3)	-0.023 (2)	0.014 (2)	0.008 (2)
C41	0.093 (4)	0.070 (3)	0.087 (3)	-0.040 (3)	0.027 (3)	0.011 (3)
C42	0.118 (4)	0.055 (2)	0.049 (2)	-0.042 (3)	-0.004 (2)	0.0045 (19)
C43	0.099 (4)	0.081 (3)	0.050 (2)	-0.042 (3)	-0.025 (2)	0.017 (2)
C44	0.067 (3)	0.083 (3)	0.0443 (19)	-0.042 (2)	-0.0161 (17)	0.0071 (19)

---

*Geometric parameters (Å, °)*

Rh1—N11	2.065 (2)	C14—C20	1.420 (7)
Rh1—N1	2.168 (2)	C14—H14	0.9500
Rh1—P2	2.2531 (8)	C15—C16	1.370 (8)
Rh1—P1	2.2897 (7)	C15—C20	1.392 (7)
Rh1—C12	2.3338 (7)	C15—H15	0.9500
Rh1—C11	2.3787 (6)	C16—C17	1.402 (7)
P1—C8	1.800 (3)	C16—H16	0.9500
P1—C21	1.815 (3)	C17—C18	1.378 (5)
P1—C27	1.818 (3)	C17—H17	0.9500
P2—C18	1.806 (3)	C18—C19	1.413 (5)
P2—C39	1.816 (3)	C19—C20	1.418 (5)
P2—C33	1.821 (3)	C21—C22	1.372 (5)
P3—F3A	1.513 (6)	C21—C26	1.389 (5)
P3—F6B	1.518 (8)	C22—C23	1.396 (6)
P3—F6A	1.519 (7)	C22—H22	0.9500
P3—F2A	1.526 (6)	C23—C24	1.409 (10)
P3—F5B	1.526 (8)	C23—H23	0.9500
P3—F1A	1.528 (6)	C24—C25	1.341 (9)
P3—F4B	1.529 (8)	C24—H24	0.9500
P3—F3B	1.538 (8)	C25—C26	1.360 (6)
P3—F4A	1.539 (7)	C25—H25	0.9500
P3—F2B	1.552 (8)	C26—H26	0.9500
P3—F1B	1.558 (9)	C27—C32	1.367 (5)
P3—F5A	1.576 (7)	C27—C28	1.397 (5)
N1—C2	1.322 (4)	C28—C29	1.376 (5)
N1—C9	1.367 (4)	C28—H28	0.9500
N11—C12	1.321 (4)	C29—C30	1.370 (7)
N11—C19	1.368 (4)	C29—H29	0.9500
C13A—C1A	1.754 (10)	C30—C31	1.353 (7)
C14A—C1A	1.721 (10)	C30—H30	0.9500
C1A—H1A	0.9900	C31—C32	1.402 (5)
C1A—H1B	0.9900	C31—H31	0.9500
O1B—C1B	1.412 (16)	C32—H32	0.9500
O1B—H1F	0.8400	C33—C34	1.387 (4)
C1B—H1C	0.9800	C33—C38	1.388 (4)
C1B—H1E	0.9800	C34—C35	1.387 (4)
C1B—H1D	0.9800	C34—H34	0.9500
C2—C3	1.392 (5)	C35—C36	1.370 (5)
C2—H2	0.9500	C35—H35	0.9500
C3—C4	1.344 (5)	C36—C37	1.361 (5)
C3—H3	0.9500	C36—H36	0.9500
C4—C10	1.405 (5)	C37—C38	1.385 (4)
C4—H4	0.9500	C37—H37	0.9500
C5—C6	1.346 (5)	C38—H38	0.9500
C5—C10	1.407 (5)	C39—C40	1.381 (5)
C5—H5	0.9500	C39—C44	1.395 (5)

C6—C7	1.396 (5)	C40—C41	1.385 (6)
C6—H6	0.9500	C40—H40	0.9500
C7—C8	1.373 (4)	C41—C42	1.351 (6)
C7—H7	0.9500	C41—H41	0.9500
C8—C9	1.416 (4)	C42—C43	1.377 (6)
C9—C10	1.418 (4)	C42—H42	0.9500
C12—C13	1.392 (5)	C43—C44	1.387 (5)
C12—H12	0.9500	C43—H43	0.9500
C13—C14	1.332 (7)	C44—H44	0.9500
C13—H13	0.9500		
N11—Rh1—N1	95.37 (10)	C4—C10—C9	118.3 (3)
N11—Rh1—P2	84.53 (8)	C5—C10—C9	117.6 (3)
N1—Rh1—P2	177.65 (6)	N11—C12—C13	123.2 (4)
N11—Rh1—P1	91.56 (6)	N11—C12—H12	118.4
N1—Rh1—P1	81.81 (6)	C13—C12—H12	118.4
P2—Rh1—P1	100.55 (3)	C14—C13—C12	119.5 (4)
N11—Rh1—Cl2	177.29 (7)	C14—C13—H13	120.2
N1—Rh1—Cl2	86.06 (7)	C12—C13—H13	120.2
P2—Rh1—Cl2	93.94 (3)	C13—C14—C20	120.1 (4)
P1—Rh1—Cl2	90.92 (3)	C13—C14—H14	120.0
N11—Rh1—Cl1	87.29 (6)	C20—C14—H14	120.0
N1—Rh1—Cl1	90.19 (6)	C16—C15—C20	120.8 (4)
P2—Rh1—Cl1	87.45 (3)	C16—C15—H15	119.6
P1—Rh1—Cl1	171.78 (3)	C20—C15—H15	119.6
Cl2—Rh1—Cl1	90.41 (3)	C15—C16—C17	120.3 (4)
C8—P1—C21	104.72 (15)	C15—C16—H16	119.8
C8—P1—C27	105.37 (14)	C17—C16—H16	119.8
C21—P1—C27	104.93 (15)	C18—C17—C16	120.7 (5)
C8—P1—Rh1	100.60 (9)	C18—C17—H17	119.6
C21—P1—Rh1	111.85 (9)	C16—C17—H17	119.6
C27—P1—Rh1	127.10 (11)	C17—C18—C19	119.2 (4)
C18—P2—C39	108.44 (16)	C17—C18—P2	125.0 (3)
C18—P2—C33	107.38 (14)	C19—C18—P2	115.7 (2)
C39—P2—C33	104.33 (14)	N11—C19—C18	119.6 (3)
C18—P2—Rh1	100.25 (12)	N11—C19—C20	120.5 (4)
C39—P2—Rh1	119.73 (10)	C18—C19—C20	119.8 (3)
C33—P2—Rh1	116.00 (9)	C15—C20—C19	119.1 (5)
F3A—P3—F6A	98.7 (10)	C15—C20—C14	123.3 (4)
F3A—P3—F2A	96.5 (6)	C19—C20—C14	117.6 (4)
F6A—P3—F2A	92.8 (7)	C22—C21—C26	119.1 (4)
F6B—P3—F5B	99.3 (12)	C22—C21—P1	122.0 (3)
F3A—P3—F1A	89.1 (5)	C26—C21—P1	118.8 (3)
F6A—P3—F1A	170.5 (9)	C21—C22—C23	119.6 (5)
F2A—P3—F1A	91.7 (7)	C21—C22—H22	120.2
F6B—P3—F4B	85.3 (8)	C23—C22—H22	120.2
F5B—P3—F4B	87.0 (10)	C22—C23—C24	119.4 (6)
F6B—P3—F3B	109.7 (17)	C22—C23—H23	120.3

F5B—P3—F3B	89.9 (12)	C24—C23—H23	120.3
F4B—P3—F3B	165.0 (17)	C25—C24—C23	120.0 (5)
F3A—P3—F4A	163.5 (12)	C25—C24—H24	120.0
F6A—P3—F4A	93.0 (9)	C23—C24—H24	120.0
F2A—P3—F4A	94.5 (11)	C24—C25—C26	120.5 (6)
F1A—P3—F4A	78.3 (11)	C24—C25—H25	119.8
F6B—P3—F2B	73.3 (16)	C26—C25—H25	119.8
F5B—P3—F2B	169.0 (18)	C25—C26—C21	121.4 (5)
F4B—P3—F2B	100.3 (14)	C25—C26—H26	119.3
F3B—P3—F2B	85.1 (12)	C21—C26—H26	119.3
F6B—P3—F1B	158.1 (19)	C32—C27—C28	119.0 (3)
F5B—P3—F1B	98.6 (18)	C32—C27—P1	120.4 (3)
F4B—P3—F1B	83.2 (15)	C28—C27—P1	120.6 (3)
F3B—P3—F1B	82.8 (16)	C29—C28—C27	120.7 (4)
F2B—P3—F1B	90.6 (15)	C29—C28—H28	119.7
F3A—P3—F5A	76.1 (9)	C27—C28—H28	119.7
F6A—P3—F5A	84.0 (6)	C30—C29—C28	119.7 (4)
F2A—P3—F5A	171.3 (9)	C30—C29—H29	120.1
F1A—P3—F5A	92.7 (7)	C28—C29—H29	120.1
F4A—P3—F5A	93.7 (10)	C31—C30—C29	120.3 (4)
C2—N1—C9	118.3 (3)	C31—C30—H30	119.8
C2—N1—Rh1	122.6 (2)	C29—C30—H30	119.8
C9—N1—Rh1	118.45 (18)	C30—C31—C32	120.7 (5)
C12—N11—C19	119.0 (3)	C30—C31—H31	119.7
C12—N11—Rh1	121.9 (2)	C32—C31—H31	119.7
C19—N11—Rh1	119.2 (2)	C27—C32—C31	119.6 (4)
C14A—C1A—C13A	104.2 (8)	C27—C32—H32	120.2
C14A—C1A—H1A	110.9	C31—C32—H32	120.2
C13A—C1A—H1A	110.9	C34—C33—C38	119.2 (3)
C14A—C1A—H1B	110.9	C34—C33—P2	120.3 (2)
C13A—C1A—H1B	110.9	C38—C33—P2	120.5 (2)
H1A—C1A—H1B	108.9	C33—C34—C35	120.2 (3)
C1B—O1B—H1F	109.5	C33—C34—H34	119.9
O1B—C1B—H1C	109.5	C35—C34—H34	119.9
O1B—C1B—H1E	109.5	C36—C35—C34	120.1 (3)
H1C—C1B—H1E	109.5	C36—C35—H35	120.0
O1B—C1B—H1D	109.5	C34—C35—H35	120.0
H1C—C1B—H1D	109.5	C37—C36—C35	119.9 (3)
H1E—C1B—H1D	109.5	C37—C36—H36	120.0
N1—C2—C3	123.5 (3)	C35—C36—H36	120.0
N1—C2—H2	118.2	C36—C37—C38	121.2 (3)
C3—C2—H2	118.2	C36—C37—H37	119.4
C4—C3—C2	119.5 (3)	C38—C37—H37	119.4
C4—C3—H3	120.2	C37—C38—C33	119.4 (3)
C2—C3—H3	120.2	C37—C38—H38	120.3
C3—C4—C10	119.5 (3)	C33—C38—H38	120.3
C3—C4—H4	120.2	C40—C39—C44	118.6 (3)
C10—C4—H4	120.2	C40—C39—P2	122.4 (3)

C6—C5—C10	121.5 (3)	C44—C39—P2	119.0 (2)
C6—C5—H5	119.3	C39—C40—C41	119.7 (4)
C10—C5—H5	119.3	C39—C40—H40	120.2
C5—C6—C7	121.1 (3)	C41—C40—H40	120.2
C5—C6—H6	119.5	C42—C41—C40	121.5 (4)
C7—C6—H6	119.5	C42—C41—H41	119.3
C8—C7—C6	120.3 (3)	C40—C41—H41	119.3
C8—C7—H7	119.9	C41—C42—C43	120.1 (4)
C6—C7—H7	119.9	C41—C42—H42	120.0
C7—C8—C9	119.2 (3)	C43—C42—H42	120.0
C7—C8—P1	124.0 (2)	C42—C43—C44	119.4 (4)
C9—C8—P1	116.7 (2)	C42—C43—H43	120.3
N1—C9—C8	118.9 (3)	C44—C43—H43	120.3
N1—C9—C10	120.8 (3)	C43—C44—C39	120.8 (4)
C8—C9—C10	120.3 (3)	C43—C44—H44	119.6
C4—C10—C5	124.1 (3)	C39—C44—H44	119.6
C9—N1—C2—C3	0.1 (5)	C13—C14—C20—C15	178.9 (3)
Rh1—N1—C2—C3	-170.8 (3)	C13—C14—C20—C19	-0.5 (5)
N1—C2—C3—C4	-1.2 (6)	C8—P1—C21—C22	11.2 (3)
C2—C3—C4—C10	1.7 (6)	C27—P1—C21—C22	121.9 (3)
C10—C5—C6—C7	-1.6 (8)	Rh1—P1—C21—C22	-96.9 (3)
C5—C6—C7—C8	-0.1 (7)	C8—P1—C21—C26	-172.8 (2)
C6—C7—C8—C9	2.6 (6)	C27—P1—C21—C26	-62.0 (3)
C6—C7—C8—P1	-173.1 (3)	Rh1—P1—C21—C26	79.2 (2)
C21—P1—C8—C7	77.0 (3)	C26—C21—C22—C23	0.1 (5)
C27—P1—C8—C7	-33.4 (3)	P1—C21—C22—C23	176.2 (3)
Rh1—P1—C8—C7	-166.8 (3)	C21—C22—C23—C24	0.1 (6)
C21—P1—C8—C9	-98.7 (2)	C22—C23—C24—C25	-0.7 (8)
C27—P1—C8—C9	150.9 (2)	C23—C24—C25—C26	1.1 (8)
Rh1—P1—C8—C9	17.4 (3)	C24—C25—C26—C21	-0.8 (7)
C2—N1—C9—C8	-179.0 (3)	C22—C21—C26—C25	0.2 (5)
Rh1—N1—C9—C8	-7.6 (4)	P1—C21—C26—C25	-175.9 (3)
C2—N1—C9—C10	0.4 (5)	C8—P1—C27—C32	-73.9 (3)
Rh1—N1—C9—C10	171.7 (2)	C21—P1—C27—C32	175.8 (3)
C7—C8—C9—N1	176.0 (3)	Rh1—P1—C27—C32	42.6 (4)
P1—C8—C9—N1	-8.1 (4)	C8—P1—C27—C28	103.4 (3)
C7—C8—C9—C10	-3.4 (5)	C21—P1—C27—C28	-6.9 (3)
P1—C8—C9—C10	172.6 (3)	Rh1—P1—C27—C28	-140.1 (3)
C3—C4—C10—C5	176.4 (4)	C32—C27—C28—C29	0.2 (6)
C3—C4—C10—C9	-1.2 (6)	P1—C27—C28—C29	-177.2 (4)
C6—C5—C10—C4	-176.9 (4)	C27—C28—C29—C30	-1.2 (7)
C6—C5—C10—C9	0.7 (7)	C28—C29—C30—C31	1.1 (9)
N1—C9—C10—C4	0.2 (5)	C29—C30—C31—C32	0.0 (9)
C8—C9—C10—C4	179.5 (3)	C28—C27—C32—C31	0.9 (7)
N1—C9—C10—C5	-177.6 (3)	P1—C27—C32—C31	178.3 (4)
C8—C9—C10—C5	1.7 (5)	C30—C31—C32—C27	-1.0 (8)
C19—N11—C12—C13	1.5 (4)	C18—P2—C33—C34	171.2 (3)

---

Rh1—N11—C12—C13	-176.6 (2)	C39—P2—C33—C34	-73.8 (3)
N11—C12—C13—C14	0.2 (5)	Rh1—P2—C33—C34	60.1 (3)
C12—C13—C14—C20	-0.7 (5)	C18—P2—C33—C38	-7.1 (3)
C20—C15—C16—C17	-0.6 (7)	C39—P2—C33—C38	107.8 (3)
C15—C16—C17—C18	1.1 (7)	Rh1—P2—C33—C38	-118.2 (2)
C16—C17—C18—C19	-0.4 (5)	C38—C33—C34—C35	-1.2 (5)
C16—C17—C18—P2	-178.5 (3)	P2—C33—C34—C35	-179.6 (2)
C39—P2—C18—C17	-47.8 (3)	C33—C34—C35—C36	0.0 (5)
C33—P2—C18—C17	64.4 (3)	C34—C35—C36—C37	0.8 (5)
Rh1—P2—C18—C17	-174.1 (3)	C35—C36—C37—C38	-0.4 (5)
C39—P2—C18—C19	134.0 (2)	C36—C37—C38—C33	-0.8 (5)
C33—P2—C18—C19	-113.8 (2)	C34—C33—C38—C37	1.6 (4)
Rh1—P2—C18—C19	7.8 (2)	P2—C33—C38—C37	180.0 (2)
C12—N11—C19—C18	178.8 (3)	C18—P2—C39—C40	-13.0 (3)
Rh1—N11—C19—C18	-3.0 (3)	C33—P2—C39—C40	-127.2 (3)
C12—N11—C19—C20	-2.7 (4)	Rh1—P2—C39—C40	100.9 (3)
Rh1—N11—C19—C20	175.5 (2)	C18—P2—C39—C44	164.7 (3)
C17—C18—C19—N11	177.8 (3)	C33—P2—C39—C44	50.5 (3)
P2—C18—C19—N11	-4.0 (4)	Rh1—P2—C39—C44	-81.3 (3)
C17—C18—C19—C20	-0.8 (5)	C44—C39—C40—C41	0.8 (6)
P2—C18—C19—C20	177.5 (2)	P2—C39—C40—C41	178.6 (3)
C16—C15—C20—C19	-0.6 (6)	C39—C40—C41—C42	0.1 (7)
C16—C15—C20—C14	-180.0 (4)	C40—C41—C42—C43	-1.2 (7)
N11—C19—C20—C15	-177.2 (3)	C41—C42—C43—C44	1.3 (7)
C18—C19—C20—C15	1.3 (4)	C42—C43—C44—C39	-0.3 (6)
N11—C19—C20—C14	2.2 (4)	C40—C39—C44—C43	-0.7 (6)
C18—C19—C20—C14	-179.3 (3)	P2—C39—C44—C43	-178.5 (3)

---