



# Crystal structure of bis[4-(dimethylamino)pyridine- $\kappa N^1$ ]bis(methanol- $\kappa O$ )-bis(thiocyanato- $\kappa N$ )manganese(II)

Stefan Suckert,\* Inke Jess and Christian Näther

Institut für Anorganische Chemie, Christian-Albrechts-Universität Kiel, Max-Eyth-Strasse 2, 24118 Kiel, Germany. \*Correspondence e-mail: ssuckert@ac.uni-kiel.de

Received 11 March 2015; accepted 13 April 2015

Edited by J. Simpson, University of Otago, New Zealand


The whole molecule of the title compound,  $[\text{Mn}(\text{NCS})_2(\text{CH}_3\text{OH})_2(\text{C}_5\text{H}_6\text{N}_2)_2]$ , is generated by inversion symmetry. The  $\text{Mn}^{\text{II}}$  ion, which is located on an inversion center, is coordinated by two 4-(dimethylamino)pyridine ligands, two methanol ligands and two terminally  $N$ -bonded thiocyanate anions, forming a slightly distorted octahedron. In the crystal, molecules are linked by  $\text{O}-\text{H}\cdots\text{S}$  hydrogen bonds, forming chains extending along the  $a$ -axis direction.

**Keywords:** crystal structure; discrete complex; octahedral coordination; hydrogen bonding.

**CCDC reference:** 1059105

## 1. Related literature

For the structure of another discrete complex with 4-(dimethylamino)pyridine and thiocyanate ligands, see: Chen *et al.* (2007). For general background to this work, see: Näther *et al.* (2013).



## 2. Experimental

### 2.1. Crystal data

$[\text{Mn}(\text{NCS})_2(\text{CH}_3\text{OH})_2(\text{C}_5\text{H}_6\text{N}_2)_2]$	$\gamma = 79.809 (11)^\circ$
$M_r = 479.52$	$V = 584.72 (10) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 7.0771 (7) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.1586 (8) \text{ \AA}$	$\mu = 0.77 \text{ mm}^{-1}$
$c = 10.6491 (10) \text{ \AA}$	$T = 180 \text{ K}$
$\alpha = 76.381 (11)^\circ$	$0.16 \times 0.10 \times 0.04 \text{ mm}$
$\beta = 81.672 (11)^\circ$	

### 2.2. Data collection

Stoe IPDS-1 diffractometer	4585 measured reflections
Absorption correction: numerical ( <i>X-SHAPE</i> and <i>X-RED32</i> ; Stoe & Cie, 2008)	2459 independent reflections
$T_{\text{min}} = 0.903$ , $T_{\text{max}} = 0.959$	1885 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.039$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	133 parameters
$wR(F^2) = 0.100$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
2459 reflections	$\Delta\rho_{\text{min}} = -0.59 \text{ e \AA}^{-3}$

**Table 1**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O31}-\text{H1O}\cdots\text{S1}^i$	0.85	2.42	3.2409 (18)	161

Symmetry code: (i)  $x - 1, y, z$ .

Data collection: *X-AREA* (Stoe & Cie, 2008); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008) and *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

## Acknowledgements

We gratefully acknowledge financial support by the DFG (project number NA 720/5-1) and the State of Schleswig-Holstein. We thank Professor Dr Wolfgang Bensch for access to his experimental facilities.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SJ5446).

## References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Chen, F.-J., Liu, G.-Q. & Zeng, Z.-Z. (2007). *Anal. Sci.* **23**, x253–x254.
- Näther, C., Wöhlert, S., Boeckmann, J., Wriedt, M. & Jess, I. (2013). *Z. Anorg. Allg. Chem.* **639**, 2696–2714.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Stoe & Cie (2008). *X-AREA*, *X-RED32* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2015). E71, m126 [https://doi.org/10.1107/S2056989015007318]

## Crystal structure of bis[4-(dimethylamino)pyridine- $\kappa N^1$ ]bis(methanol- $\kappa O$ )bis-(thiocyanato- $\kappa N$ )manganese(II)

Stefan Suckert, Inke Jess and Christian Näther

### S1. Synthesis and crystallization

MnSO<sub>4</sub>·H<sub>2</sub>O was purchased from Merck and 4-(dimethylamino)pyridine and Ba(NCS)<sub>2</sub>·3 H<sub>2</sub>O were purchased from Alfa Aesar. Mn(NCS)<sub>2</sub> was synthesized by stirring 17.97 g (58.44 mmol) Ba(NCS)<sub>2</sub>·3 H<sub>2</sub>O and 9.88 g (58.44 mmol) MnSO<sub>4</sub>·H<sub>2</sub>O in 300 mL H<sub>2</sub>O at room temperature for three hours. The white precipitate of BaSO<sub>4</sub> was filtered off and the solvent removed with a rotary evaporator. The homogeneity of the product was investigated by X-ray powder diffraction and elemental analysis. The title compound was prepared by the reaction of (0.18 mmol) 30.8 mg Mn(NCS)<sub>2</sub> and (0.3 mmol) 36.7 mg 4-(dimethylamino)pyridine in 1.0 mL methanol at room temperature. After a few days colorless plate shaped crystals of the title compound were obtained.

### S2. Refinement

The C—H H atoms were positioned with idealized geometry and were refined isotropically with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  (1.5 for methyl H atoms) using a riding model with C—H = 0.95 Å for aromatic and C—H = 0.98 Å for methyl H atoms. The O—H H atom was located in a difference map, its bond length set to ideal values of 0.85 Å and refined with  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$  using a riding model.




Figure 1

Structure of the title complex with atom labelling. Displacement ellipsoids are drawn at the 50% probability level. Symmetry code:  $i = -x+1, -y, -z+1$ .




Figure 2

Crystal structure of the title compound viewed perpendicular to the crystallographic  $a,c$  plane.

**Bis[4-(dimethylamino)pyridine- $\kappa N^1$ ]bis(methanol- $\kappa O$ )bis(thiocyanato- $\kappa N$ )manganese(II)]**

*Crystal data*

[Mn(NCS)<sub>2</sub>(CH<sub>4</sub>O)<sub>2</sub>(C<sub>3</sub>H<sub>6</sub>N<sub>2</sub>)<sub>2</sub>]

$M_r = 479.52$

Triclinic,  $P\bar{1}$

$a = 7.0771$  (7) Å

$b = 8.1586$  (8) Å

$c = 10.6491$  (10) Å

$\alpha = 76.381$  (11)°

$\beta = 81.672$  (11)°

$\gamma = 79.809$  (11)°

$V = 584.72$  (10) Å<sup>3</sup>

$Z = 1$

$F(000) = 251$

$D_x = 1.362$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4585 reflections

$\theta = 2.6$ – $27.0$ °

$\mu = 0.77$  mm<sup>-1</sup>

$T = 180$  K

Plate, colorless

$0.16 \times 0.10 \times 0.04$  mm

*Data collection*

Stoe IPDS-1  
diffractometer

Radiation source: fine-focus sealed tube  
phi scans

Absorption correction: numerical  
(*X-SHAPE* and *X-RED32*; Stoe & Cie, 2008)

$T_{\min} = 0.903$ ,  $T_{\max} = 0.959$

4585 measured reflections

2459 independent reflections

1885 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

$\theta_{\max} = 27.0$ °,  $\theta_{\min} = 2.6$ °

$h = -9 \rightarrow 9$

$k = -10 \rightarrow 10$

$l = -13 \rightarrow 13$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.100$

$S = 1.04$

2459 reflections

133 parameters

0 restraints

Hydrogen site location: mixed

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0598P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.30$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.59$  e Å<sup>-3</sup>

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.5000	0.0000	0.5000	0.02111 (15)
N1	0.6294 (3)	0.0926 (3)	0.63997 (19)	0.0312 (5)
C1	0.7395 (3)	0.1199 (3)	0.7004 (2)	0.0236 (5)
S1	0.89299 (9)	0.15954 (10)	0.78630 (6)	0.0394 (2)
O31	0.2205 (2)	0.1737 (2)	0.53900 (16)	0.0339 (4)
H1O	0.1462	0.1444	0.6079	0.051*
C31	0.0945 (4)	0.2643 (4)	0.4439 (3)	0.0486 (7)
H31A	-0.0136	0.3337	0.4847	0.073*
H31B	0.0448	0.1828	0.4074	0.073*
H31C	0.1659	0.3386	0.3743	0.073*
N11	0.5945 (3)	0.2090 (2)	0.33954 (17)	0.0222 (4)
N12	0.7263 (3)	0.6192 (2)	0.03652 (18)	0.0274 (4)
C11	0.5795 (3)	0.3682 (3)	0.3576 (2)	0.0264 (5)
H11	0.5369	0.3866	0.4427	0.032*
C12	0.6210 (3)	0.5063 (3)	0.2624 (2)	0.0252 (5)
H12	0.6057	0.6157	0.2825	0.030*
C13	0.6867 (3)	0.4861 (3)	0.1345 (2)	0.0210 (4)
C14	0.7077 (3)	0.3182 (3)	0.1162 (2)	0.0230 (4)
H14	0.7554	0.2942	0.0334	0.028*
C15	0.6594 (3)	0.1895 (3)	0.2177 (2)	0.0235 (5)
H15	0.6727	0.0783	0.2011	0.028*
C16	0.6858 (4)	0.7919 (3)	0.0569 (3)	0.0353 (6)
H16A	0.7210	0.8713	-0.0243	0.053*
H16B	0.7612	0.8015	0.1244	0.053*
H16C	0.5479	0.8196	0.0844	0.053*
C17	0.7938 (3)	0.5957 (3)	-0.0944 (2)	0.0309 (5)
H17A	0.8140	0.7059	-0.1514	0.046*
H17B	0.6972	0.5482	-0.1267	0.046*
H17C	0.9158	0.5171	-0.0934	0.046*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Mn1	0.0216 (2)	0.0237 (3)	0.0185 (2)	-0.00575 (18)	0.00127 (17)	-0.00564 (18)
N1	0.0330 (10)	0.0364 (12)	0.0282 (10)	-0.0112 (9)	-0.0016 (8)	-0.0117 (9)
C1	0.0254 (11)	0.0242 (11)	0.0201 (10)	-0.0035 (8)	0.0040 (8)	-0.0067 (9)
S1	0.0261 (3)	0.0703 (5)	0.0288 (3)	-0.0110 (3)	0.0005 (2)	-0.0240 (3)
O31	0.0250 (8)	0.0444 (11)	0.0258 (8)	0.0030 (7)	0.0033 (6)	-0.0046 (8)
C31	0.0373 (15)	0.062 (2)	0.0343 (14)	0.0113 (13)	-0.0022 (11)	-0.0017 (13)

N11	0.0272 (9)	0.0210 (9)	0.0188 (9)	-0.0037 (7)	0.0014 (7)	-0.0070 (7)
N12	0.0300 (10)	0.0241 (10)	0.0248 (10)	-0.0039 (8)	0.0015 (8)	-0.0017 (8)
C11	0.0325 (12)	0.0263 (12)	0.0210 (11)	-0.0043 (9)	0.0028 (8)	-0.0100 (9)
C12	0.0306 (11)	0.0201 (11)	0.0267 (11)	-0.0032 (9)	-0.0011 (9)	-0.0100 (9)
C13	0.0143 (9)	0.0245 (11)	0.0238 (10)	-0.0031 (8)	-0.0019 (7)	-0.0040 (9)
C14	0.0205 (10)	0.0270 (11)	0.0210 (10)	-0.0009 (8)	0.0022 (8)	-0.0090 (9)
C15	0.0222 (10)	0.0242 (11)	0.0258 (11)	-0.0021 (8)	0.0013 (8)	-0.0119 (9)
C16	0.0405 (14)	0.0234 (12)	0.0399 (14)	-0.0064 (10)	-0.0011 (11)	-0.0034 (11)
C17	0.0274 (11)	0.0402 (14)	0.0217 (11)	-0.0070 (10)	-0.0001 (9)	0.0001 (10)

*Geometric parameters (Å, °)*

Mn1—N1 <sup>i</sup>	2.192 (2)	N12—C16	1.448 (3)
Mn1—N1	2.192 (2)	N12—C17	1.452 (3)
Mn1—N11	2.2302 (17)	C11—C12	1.370 (3)
Mn1—N11 <sup>i</sup>	2.2302 (17)	C11—H11	0.9500
Mn1—O31	2.2676 (17)	C12—C13	1.412 (3)
Mn1—O31 <sup>i</sup>	2.2676 (17)	C12—H12	0.9500
N1—C1	1.160 (3)	C13—C14	1.408 (3)
C1—S1	1.634 (2)	C14—C15	1.368 (3)
O31—C31	1.429 (3)	C14—H14	0.9500
O31—H1O	0.8500	C15—H15	0.9500
C31—H31A	0.9800	C16—H16A	0.9800
C31—H31B	0.9800	C16—H16B	0.9800
C31—H31C	0.9800	C16—H16C	0.9800
N11—C11	1.341 (3)	C17—H17A	0.9800
N11—C15	1.349 (3)	C17—H17B	0.9800
N12—C13	1.355 (3)	C17—H17C	0.9800
N1 <sup>i</sup> —Mn1—N1	180.0	C13—N12—C17	121.3 (2)
N1 <sup>i</sup> —Mn1—N11	89.19 (7)	C16—N12—C17	117.81 (19)
N1—Mn1—N11	90.81 (7)	N11—C11—C12	124.8 (2)
N1 <sup>i</sup> —Mn1—N11 <sup>i</sup>	90.81 (7)	N11—C11—H11	117.6
N1—Mn1—N11 <sup>i</sup>	89.19 (7)	C12—C11—H11	117.6
N11—Mn1—N11 <sup>i</sup>	180.0	C11—C12—C13	120.0 (2)
N1 <sup>i</sup> —Mn1—O31	90.50 (7)	C11—C12—H12	120.0
N1—Mn1—O31	89.50 (7)	C13—C12—H12	120.0
N11—Mn1—O31	89.07 (6)	N12—C13—C14	122.6 (2)
N11 <sup>i</sup> —Mn1—O31	90.93 (6)	N12—C13—C12	122.2 (2)
N1 <sup>i</sup> —Mn1—O31 <sup>i</sup>	89.50 (7)	C14—C13—C12	115.22 (19)
N1—Mn1—O31 <sup>i</sup>	90.50 (7)	C15—C14—C13	120.1 (2)
N11—Mn1—O31 <sup>i</sup>	90.93 (6)	C15—C14—H14	120.0
N11 <sup>i</sup> —Mn1—O31 <sup>i</sup>	89.07 (6)	C13—C14—H14	120.0
O31—Mn1—O31 <sup>i</sup>	180.0	N11—C15—C14	124.7 (2)
C1—N1—Mn1	162.79 (19)	N11—C15—H15	117.6
N1—C1—S1	179.4 (2)	C14—C15—H15	117.6
C31—O31—Mn1	125.44 (16)	N12—C16—H16A	109.5
C31—O31—H1O	105.1	N12—C16—H16B	109.5

Mn1—O31—H1O	119.1	H16A—C16—H16B	109.5
O31—C31—H31A	109.5	N12—C16—H16C	109.5
O31—C31—H31B	109.5	H16A—C16—H16C	109.5
H31A—C31—H31B	109.5	H16B—C16—H16C	109.5
O31—C31—H31C	109.5	N12—C17—H17A	109.5
H31A—C31—H31C	109.5	N12—C17—H17B	109.5
H31B—C31—H31C	109.5	H17A—C17—H17B	109.5
C11—N11—C15	115.11 (18)	N12—C17—H17C	109.5
C11—N11—Mn1	120.90 (14)	H17A—C17—H17C	109.5
C15—N11—Mn1	123.89 (15)	H17B—C17—H17C	109.5
C13—N12—C16	120.6 (2)		

Symmetry code: (i)  $-x+1, -y, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O31—H1O $\cdots$ S1 <sup>ii</sup>	0.85	2.42	3.2409 (18)	161

Symmetry code: (ii)  $x-1, y, z$ .