

Crystal structure of *trans*-(1,8-dibutyl-1,3,6,8,10,13-hexaazacyclotetradecane- $\kappa^4N^3,N^6,N^{10},N^{13}$)bis(thiocyanato- κN)nickel(II) from synchrotron data

Dae-Woong Kim,^a Jong Jin Kim,^b Jong Won Shin,^c Jin Hong Kim^c and Dohyun Moon^{c*}

Received 28 May 2015
Accepted 8 June 2015

Edited by M. Gdaniec, Adam Mickiewicz University, Poland

^aDepartment of Chemistry and Green-Nano Materials Research Center, Kyungpook National University, Daegu, 702-701, Republic of Korea, ^bDepartment of Applied Chemistry, College of Engineering, Kyungpook National University, Daegu, 702-701, Republic of Korea, and ^cBeamline Department, Pohang Accelerator Laboratory 80, Jigokro-127-beongil, Nam-Gu Pohang, Gyeongbuk 790-784, South Korea. *Correspondence e-mail: dmoon@postech.ac.kr


Keywords: crystal structure; azamacrocyclic ligand; Jahn–Teller distortion; sodium thiocyanate; hydrogen bonding; synchrotron data


CCDC reference: 1405450
Supporting information: this article has supporting information at journals.iucr.org/e

The crystal structure of the title compound, $[\text{Ni}(\text{NCS})_2(\text{C}_{16}\text{H}_{38}\text{N}_6)]$, has been determined from synchrotron data. The asymmetric unit consists of two halves of the complex molecules which have their Ni^{II} atoms located on inversion centres. The Ni^{II} ions show a tetragonally distorted octahedral coordination geometry, with four secondary amine N atoms of the azamacrocyclic ligand in the equatorial plane and two N atoms of the thiocyanate anions in the axial positions. The average equatorial Ni–N bond length [2.070 (5) Å] is shorter than the average axial Ni–N bond length [2.107 (18) Å]. Only half of the macrocyclic ligand N–H groups are involved in hydrogen bonding. The complex molecules are connected *via* intermolecular N–H...S hydrogen bonds into two symmetry-independent one-dimensional polymeric structures extending along the *b*-axis direction. One of the *n*-butyl substituents of the macrocycle exhibits conformational disorder with a refined occupancy ratio of 0.630:0.370.

1. Chemical context

Coordination compounds, including those formed by macrocyclic ligands, have attracted wide interest of material sciences, because of their potential applications (Lehn, 1995; Zhou *et al.*, 2012). In particular, Ni^{II} macrocyclic complexes having vacant sites in the axial positions have been used for the synthesis of new supramolecular materials with interesting properties, including chiral recognition (Ryoo *et al.*, 2010) and gas storage (Suh *et al.*, 2012). For example, Ni^{II} complexes with alkyl-substituted tetra-azamacrocyclic ligands and anionic tetrazole derivatives, metal cyanide and azide (Shen *et al.*, 2012; Kim *et al.*, 2015) have been studied as magnetic materials and substrates for crystal engineering. The thiocyanate ion is a versatile anionic ligand which can easily bind to a transition metal ion as a terminal or bridging ligand through the nitrogen and/or the sulfur atoms, thus allowing the assembly of multi-dimensional compounds or heterometallic complexes (Safarifar & Morsali, 2012; Wang & Wang, 2015). Here, we report the synthesis and crystal structure of an Ni^{II} complex with an azamacrocyclic ligand and two thiocyanate anions, *trans*-(1,8-dibutyl-1,3,6,8,10,13-hexaazacyclotetradecane- $\kappa^4N^3,N^6,N^{10},N^{13}$)bis(thiocyanato- κN)nickel(II) (I).





2. Structural commentary

The title compound (**I**) contains two crystallographically independent complex molecules that are centrosymmetric. Each Ni^{II} ion lies on an inversion centre and is coordinated by four secondary amine N atoms of the azamacrocyclic ligand in a square-planar fashion in the equatorial plane, and by two N atoms from the thiocyanate anions at the axial positions, resulting in a tetragonally distorted octahedral geometry, as shown in Fig. 1. The average equatorial bond lengths, Ni1A–N_{eq} and Ni1B–N_{eq}, are 2.070 (8) and 2.070 (3) Å, respectively. The axial bond lengths, Ni1A–N_{ax} and Ni1B–N_{ax} are 2.119 (1) and 2.093 (1) Å, respectively. The axial bonds are longer than the equatorial bonds, which can be attributed either to a large Jahn–Teller distortion effect of the Ni^{II} ion and/or to a ring contraction of the azamacrocyclic ligand (Halcrow, 2013; Kim *et al.*, 2015). The average N–C and C–S bond lengths of the thiocyanate ligands are 1.157 (1) and

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N1A–H1A···S1A ⁱ	1.00	2.73	3.5154 (17)	136
N2B–H2B···S1B ⁱⁱ	1.00	2.66	3.4556 (17)	137

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, -y + 1, -z$.

1.627 (11) Å, respectively. The former is very similar to a C≡N triple-bond length, while the latter is slightly shorter than reported C–S single-bond lengths (Bradforth *et al.*, 1993; Shin *et al.*, 2010). The six-membered chelate rings involving C2A, C3A and C2B, C3B atoms adopt a *chair* conformation, whereas the five-membered chelate rings involving C1A, C4A and C1B, C4B assume a *gauche* conformation (Min & Suh, 2001; Kim *et al.*, 2015).

3. Supramolecular features

The S atoms of the thiocyanate groups form intermolecular N–H···S hydrogen bonds with adjacent secondary amine groups of the azamacrocyclic ligand, giving rise to two symmetry-independent one-dimensional polymeric chains propagating along the *b*-axis direction (Fig. 2 and Table 1).

4. Database survey

A search of the Cambridge Structural Database (Version 5.36, Feb 2015 with two updates; Groom & Allen, 2014) indicated one complex of Ni^{II} with the same azamacrocyclic ligand having an anionic tetrazole derivative at the axial positions (Kim *et al.*, 2015).




Figure 1

View of the molecular structure of the title compound, showing the atom-labelling scheme, with displacement ellipsoids drawn at the 30% probability level. H atoms have been omitted for clarity. The minor position of the *n*-butyl substituent in the *A* molecule is not shown.

Table 2
Experimental details.

Crystal data	
Chemical formula	[Ni(NCS) ₂ (C ₁₆ H ₃₈ N ₆)]
<i>M_r</i>	489.39
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	180
<i>a</i> , <i>b</i> , <i>c</i> (Å)	8.6610 (17), 12.027 (2), 12.560 (3)
α , β , γ (°)	94.66 (3), 97.99 (3), 110.04 (3)
<i>V</i> (Å ³)	1205.4 (5)
<i>Z</i>	2
Radiation type	Synchrotron, $\lambda = 0.630$ Å
μ (mm ⁻¹)	0.72
Crystal size (mm)	0.25 × 0.15 × 0.13
Data collection	
Diffractometer	ADSC Q210 CCD area detector
Absorption correction	Empirical (using intensity measurements) (<i>HKL3000sm SCALEPACK</i> ; Otwinowski & Minor, 1997)
<i>T_{min}</i> , <i>T_{max}</i>	0.841, 0.916
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	12812, 6583, 6243
<i>R_{int}</i>	0.014
(sin θ/λ) _{max} (Å ⁻¹)	0.696
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.042, 0.111, 1.06
No. of reflections	6583
No. of parameters	287
No. of restraints	11
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	1.58, -1.11

Computer programs: *PAL ADSC Quantum-210 ADX* (Arvai & Nielsen, 1983), *HKL3000sm* (Otwinowski & Minor, 1997), *SHELXT2014* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *DIAMOND* (Putz & Brandenburg, 2007) and *publCIF* (Westrip, 2010).

5. Synthesis and crystallization

The title compound (**I**) was prepared as follows. The starting complex, [Ni(C₁₆H₃₈N₆)](ClO₄)₂, was prepared by a slightly modified method reported by Jung *et al.* (1989). To a MeCN solution (10 mL) of [Ni(C₁₆H₃₈N₆)](ClO₄)₂ (0.15 g, 0.26 mmol) was slowly added a MeCN solution (5 mL) containing sodium thiocyanate (0.042 g, 0.52 mmol) at room temperature. A pale-pink precipitate was formed, which was filtered off, washed with MeCN, and diethyl ether, and dried in air. Single crystals of the title compound were obtained by layering of the MeCN solution of sodium thiocyanate on the MeCN solution of [Ni(C₁₆H₃₈N₆)](ClO₄)₂ for several days. Yield: 0.062 g (49%). FT-IR (KBr, cm⁻¹): 3304, 3243, 2929, 2867, 2069, 1468, 1386, 1273, 1204, 1070, 925.

Safety note: Although we have experienced no problem with the compounds reported in this study, perchlorate salts of metal complexes are often explosive and should be handled with great caution.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were placed in geometrically idealized positions and constrained to ride on




Figure 2
View of the crystal packing, with N–H...S hydrogen bonds drawn as red dashed lines. H atoms have been omitted for clarity.

their parent atoms, with C–H distances of 0.98–0.99 Å and an N–H distance of 1.0 Å with *U*_{iso}(H) values of 1.2 or 1.5*U*_{eq} of the parent atoms. The C7A and C8A atoms of the macrocyclic ligand were refined as disordered over two sets of sites (C71A, C72A and C81A, C82A) with refined occupancies of 0.630 and 0.370, respectively. The bond lengths and angles of the disordered part were restrained to ensure proper geometry using DFIX and DANG instructions of *SHELXL2014* (Sheldrick, 2015b).

Acknowledgements

This work was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2014R1A1A2058815) and supported by the Institute for Basic Science (IBS, IBS-R007-D1-2015 – a01). The X-ray crystallography BL2D-SMC beamline at PLS-II was supported in part by MSIP and POSTECH.

References

- Arvai, A. J. & Nielsen, C. (1983). *ADSC Quantum-210 ADX*. Area Detector System Corporation, Poway, CA, USA.
- Bradforth, S. E., Kim, E. H., Arnold, D. W. & Neumark, D. M. (1993). *J. Chem. Phys.* **98**, 800–810.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Halcrow, M. A. (2013). *Chem. Soc. Rev.* **42**, 1784–1795.
- Jung, S. K., Kang, S. G. & Suh, M. P. (1989). *Bull. Korean Chem. Soc.* **10**, 362–366.
- Kim, D.-W., Shin, J. W., Kim, J. H. & Moon, D. (2015). *Acta Cryst.* **E71**, 173–175.
- Lehn, J.-M. (1995). In *Supramolecular Chemistry; Concepts and Perspectives*. Weinheim: VCH.
- Min, K. S. & Suh, M. P. (2001). *Chem. Eur. J.* **7**, 303–313.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. Academic Press, New York.
- Putz, H. & Brandenburg, K. (2007). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.

- Ryoo, J. J., Shin, J. W., Dho, H. S. & Min, K. S. (2010). *Inorg. Chem.* **49**, 7232–7234.
- Safarifard, V. & Morsali, A. (2012). *CrystEngComm*, **14**, 5130–5132.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Shen, X., Zhou, H., Zhang, Q., Xu, Y. & Zhou, H. (2012). *Eur. J. Inorg. Chem.* pp. 5050–5057.
- Shin, J. W., Rowthu, S. R., Ryoo, J. J. & Min, K. S. (2010). *Acta Cryst.* **E66**, m919–m920.
- Suh, M. P., Park, H. J., Prasad, T. K. & Lim, D.-W. (2012). *Chem. Rev.* **112**, 782–835.
- Wang, H.-T. & Wang, X.-L. (2015). *Acta Cryst.* **C71**, 318–321.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhou, Z., Shen, M., Cao, C., Liu, Q. & Yan, Z. (2012). *Chem. Eur. J.* **18**, 7675–7679.

supporting information

Acta Cryst. (2015). E71, 779-782 [doi:10.1107/S205698901501110X]

Crystal structure of *trans*-(1,8-dibutyl-1,3,6,8,10,13-hexaazacyclotetradecane- $\kappa^4N^3,N^6,N^{10},N^{13}$)bis(thiocyanato- κN)nickel(II) from synchrotron data

Dae-Woong Kim, Jong Jin Kim, Jong Won Shin, Jin Hong Kim and Dohyun Moon

Computing details

Data collection: *PAL ADSC Quantum-210 ADX* (Arvai & Nielsen, 1983); cell refinement: *HKL3000sm* (Otwinowski & Minor, 1997); data reduction: *HKL3000sm* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXT2014* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *DIAMOND* (Putz & Brandenburg, 2007); software used to prepare material for publication: *publCIF* (Westrip, 2010).

trans-(1,8-Dibutyl-1,3,6,8,10,13-hexaazacyclotetradecane- $\kappa^4N^3,N^6,N^{10},N^{13}$)bis(thiocyanato- κN)nickel(II)

Crystal data

[Ni(NCS)₂(C₁₆H₃₈N₆)]

$M_r = 489.39$

Triclinic, $P\bar{1}$

$a = 8.6610$ (17) Å

$b = 12.027$ (2) Å

$c = 12.560$ (3) Å

$\alpha = 94.66$ (3)°

$\beta = 97.99$ (3)°

$\gamma = 110.04$ (3)°

$V = 1205.4$ (5) Å³

$Z = 2$

$F(000) = 524$

$D_x = 1.348$ Mg m⁻³

Synchrotron radiation, $\lambda = 0.630$ Å

Cell parameters from 49914 reflections

$\theta = 0.4$ –33.6°

$\mu = 0.72$ mm⁻¹

$T = 180$ K

Block, pale pink

$0.25 \times 0.15 \times 0.13$ mm

Data collection

ADSC Q210 CCD area-detector
diffractometer

Radiation source: PLSII 2D bending magnet

ω scan

Absorption correction: empirical (using
intensity measurements)

(*HKL3000sm SCALEPACK*; Otwinowski &
Minor, 1997)

$T_{\min} = 0.841$, $T_{\max} = 0.916$

12812 measured reflections

6583 independent reflections

6243 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.014$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -12 \rightarrow 12$

$k = -16 \rightarrow 16$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.111$

$S = 1.06$

6583 reflections

287 parameters

11 restraints

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0541P)^2 + 0.7946P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$

$$\Delta\rho_{\max} = 1.58 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.11 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ni1A	0.0000	0.0000	0.5000	0.02295 (8)	
S1A	-0.39772 (6)	0.09973 (6)	0.68745 (4)	0.05126 (15)	
N1A	0.20720 (17)	0.11708 (13)	0.60541 (11)	0.0301 (3)	
H1A	0.3083	0.1069	0.5828	0.036*	
N2A	0.02319 (18)	0.10693 (13)	0.37714 (12)	0.0321 (3)	
H2A	0.1105	0.0955	0.3377	0.039*	
N3A	0.2318 (2)	0.27979 (15)	0.49597 (17)	0.0460 (4)	
N4A	-0.16012 (18)	0.07346 (14)	0.56843 (13)	0.0343 (3)	
C1A	0.1944 (2)	0.07739 (18)	0.71320 (14)	0.0373 (4)	
H1A1	0.1128	0.1036	0.7459	0.045*	
H1A2	0.3042	0.1129	0.7621	0.045*	
C2A	0.2242 (3)	0.24393 (17)	0.60232 (18)	0.0426 (4)	
H2A1	0.1280	0.2567	0.6285	0.051*	
H2A2	0.3269	0.2957	0.6528	0.051*	
C3A	0.0780 (3)	0.23584 (17)	0.41904 (19)	0.0431 (4)	
H3A1	0.0916	0.2821	0.3571	0.052*	
H3A2	-0.0106	0.2499	0.4539	0.052*	
C4A	-0.1381 (2)	0.05792 (19)	0.30124 (15)	0.0387 (4)	
H4A1	-0.1251	0.0872	0.2304	0.046*	
H4A2	-0.2223	0.0840	0.3307	0.046*	
C5A	0.3770 (3)	0.2784 (3)	0.4483 (3)	0.0632 (7)	
H5A1	0.3705	0.3095	0.3778	0.076*	
H5A2	0.3693	0.1945	0.4332	0.076*	
C6A	0.5445 (3)	0.3493 (3)	0.5164 (4)	0.0870 (11)	
H6A1	0.5622	0.3131	0.5829	0.104*	
H6A2	0.5534	0.4327	0.5378	0.104*	
C71A	0.6816 (5)	0.3457 (6)	0.4395 (4)	0.084 (2)	0.63
H71A	0.6781	0.2628	0.4233	0.100*	0.63
H71B	0.6563	0.3747	0.3701	0.100*	0.63
C81A	0.8493 (5)	0.4237 (4)	0.4995 (4)	0.0654 (11)	0.63
H81A	0.8483	0.5035	0.5213	0.098*	0.63
H81B	0.9324	0.4296	0.4525	0.098*	0.63
H81C	0.8780	0.3895	0.5642	0.098*	0.63
C72A	0.7095 (11)	0.3319 (9)	0.5052 (9)	0.077 (2)	0.37
H72A	0.8005	0.3782	0.5662	0.093*	0.37
H72B	0.6971	0.2465	0.4991	0.093*	0.37

C82A	0.7346 (11)	0.3790 (12)	0.4057 (7)	0.090 (3)	0.37
H82A	0.6498	0.3254	0.3461	0.135*	0.37
H82B	0.8460	0.3857	0.3921	0.135*	0.37
H82C	0.7255	0.4582	0.4107	0.135*	0.37
C9A	-0.2607 (2)	0.08368 (15)	0.61628 (13)	0.0304 (3)	
Ni2B	1.0000	0.5000	0.0000	0.02474 (8)	
S1B	0.48562 (7)	0.51503 (7)	-0.18976 (6)	0.0654 (2)	
N1B	0.90438 (18)	0.33335 (13)	-0.09357 (12)	0.0319 (3)	
H1B	0.7910	0.3226	-0.1338	0.038*	
N2B	0.86585 (17)	0.44340 (13)	0.12179 (11)	0.0293 (3)	
H2B	0.7493	0.4404	0.0979	0.035*	
N3B	0.7849 (2)	0.23013 (14)	0.05324 (14)	0.0368 (3)	
N4B	0.79629 (19)	0.53663 (16)	-0.07763 (14)	0.0384 (3)	
C1B	1.0161 (2)	0.34041 (17)	-0.17379 (16)	0.0395 (4)	
H1B1	0.9614	0.2749	-0.2350	0.047*	
H1B2	1.1210	0.3322	-0.1394	0.047*	
C2B	0.8871 (3)	0.23310 (16)	-0.02905 (18)	0.0401 (4)	
H2B1	0.8377	0.1568	-0.0789	0.048*	
H2B2	0.9997	0.2392	0.0065	0.048*	
C3B	0.8572 (2)	0.32257 (17)	0.14559 (15)	0.0363 (3)	
H3B1	0.9715	0.3258	0.1735	0.044*	
H3B2	0.7904	0.3008	0.2037	0.044*	
C4B	0.9457 (2)	0.53919 (17)	0.21515 (14)	0.0366 (4)	
H4B1	1.0503	0.5321	0.2515	0.044*	
H4B2	0.8700	0.5320	0.2685	0.044*	
C5B	0.6079 (2)	0.20371 (17)	0.00919 (15)	0.0366 (4)	
H5B1	0.5718	0.1390	-0.0530	0.044*	
H5B2	0.5956	0.2756	-0.0184	0.044*	
C6B	0.4936 (2)	0.16605 (17)	0.09165 (15)	0.0382 (4)	
H6B1	0.5143	0.0997	0.1253	0.046*	
H6B2	0.5206	0.2340	0.1498	0.046*	
C7B	0.3107 (3)	0.12633 (19)	0.04111 (16)	0.0413 (4)	
H7B1	0.2794	0.0511	-0.0089	0.050*	
H7B2	0.2932	0.1877	-0.0020	0.050*	
C8B	0.1979 (3)	0.1066 (2)	0.12585 (18)	0.0456 (4)	
H8B1	0.2212	0.0508	0.1725	0.068*	
H8B2	0.0808	0.0734	0.0895	0.068*	
H8B3	0.2190	0.1831	0.1701	0.068*	
C9B	0.6659 (2)	0.52684 (14)	-0.12328 (13)	0.0294 (3)	

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1A	0.02036 (13)	0.02643 (14)	0.02328 (13)	0.00930 (10)	0.00718 (9)	0.00057 (9)
S1A	0.0351 (2)	0.0888 (4)	0.0415 (3)	0.0343 (3)	0.01563 (19)	0.0052 (3)
N1A	0.0239 (6)	0.0347 (7)	0.0292 (6)	0.0085 (5)	0.0064 (5)	-0.0031 (5)
N2A	0.0294 (6)	0.0369 (7)	0.0333 (7)	0.0130 (5)	0.0111 (5)	0.0087 (5)
N3A	0.0402 (8)	0.0320 (7)	0.0611 (11)	0.0053 (6)	0.0126 (8)	0.0082 (7)

N4A	0.0292 (6)	0.0389 (7)	0.0376 (7)	0.0161 (6)	0.0098 (5)	-0.0021 (6)
C1A	0.0309 (8)	0.0535 (10)	0.0258 (7)	0.0150 (7)	0.0046 (6)	-0.0020 (7)
C2A	0.0391 (9)	0.0315 (8)	0.0492 (10)	0.0061 (7)	0.0071 (8)	-0.0078 (7)
C3A	0.0437 (10)	0.0344 (9)	0.0555 (11)	0.0155 (8)	0.0138 (8)	0.0144 (8)
C4A	0.0354 (8)	0.0551 (11)	0.0305 (8)	0.0204 (8)	0.0076 (6)	0.0118 (7)
C5A	0.0370 (11)	0.0618 (15)	0.0824 (18)	-0.0004 (10)	0.0211 (11)	0.0257 (13)
C6A	0.0414 (13)	0.0560 (16)	0.147 (3)	-0.0010 (11)	0.0047 (17)	0.0253 (19)
C71A	0.042 (2)	0.129 (4)	0.055 (2)	-0.011 (2)	0.0025 (17)	0.067 (3)
C81A	0.051 (2)	0.062 (2)	0.076 (3)	0.0168 (18)	0.0036 (19)	-0.003 (2)
C72A	0.062 (5)	0.074 (5)	0.098 (7)	0.026 (4)	0.016 (5)	0.019 (5)
C82A	0.083 (7)	0.129 (10)	0.054 (5)	0.049 (7)	-0.015 (5)	-0.011 (6)
C9A	0.0253 (7)	0.0374 (8)	0.0300 (7)	0.0144 (6)	0.0043 (5)	-0.0002 (6)
Ni2B	0.02097 (13)	0.02802 (14)	0.02616 (14)	0.01142 (10)	0.00207 (9)	0.00079 (10)
S1B	0.0371 (3)	0.0927 (5)	0.0605 (4)	0.0332 (3)	-0.0185 (2)	-0.0201 (3)
N1B	0.0280 (6)	0.0312 (6)	0.0350 (7)	0.0090 (5)	0.0090 (5)	-0.0017 (5)
N2B	0.0241 (6)	0.0333 (6)	0.0283 (6)	0.0093 (5)	0.0025 (5)	0.0009 (5)
N3B	0.0358 (7)	0.0314 (7)	0.0425 (8)	0.0102 (6)	0.0093 (6)	0.0065 (6)
N4B	0.0285 (7)	0.0482 (9)	0.0418 (8)	0.0189 (6)	0.0019 (6)	0.0083 (7)
C1B	0.0345 (8)	0.0373 (9)	0.0424 (9)	0.0079 (7)	0.0145 (7)	-0.0086 (7)
C2B	0.0419 (9)	0.0297 (8)	0.0524 (11)	0.0152 (7)	0.0153 (8)	0.0037 (7)
C3B	0.0341 (8)	0.0393 (9)	0.0347 (8)	0.0119 (7)	0.0039 (6)	0.0102 (7)
C4B	0.0299 (8)	0.0446 (9)	0.0289 (7)	0.0070 (7)	0.0060 (6)	-0.0040 (7)
C5B	0.0354 (8)	0.0332 (8)	0.0363 (8)	0.0054 (6)	0.0084 (7)	0.0042 (6)
C6B	0.0386 (9)	0.0371 (8)	0.0339 (8)	0.0067 (7)	0.0089 (7)	0.0046 (7)
C7B	0.0397 (9)	0.0445 (10)	0.0339 (8)	0.0069 (8)	0.0105 (7)	0.0029 (7)
C8B	0.0434 (10)	0.0489 (11)	0.0440 (10)	0.0125 (8)	0.0158 (8)	0.0083 (8)
C9B	0.0283 (7)	0.0303 (7)	0.0309 (7)	0.0131 (6)	0.0055 (6)	0.0003 (6)

Geometric parameters (Å, °)

Ni1A—N1A ⁱ	2.0640 (17)	C82A—H82A	0.9800
Ni1A—N1A	2.0640 (17)	C82A—H82B	0.9800
Ni1A—N2A ⁱ	2.0754 (15)	C82A—H82C	0.9800
Ni1A—N2A	2.0754 (15)	Ni2B—N2B ⁱⁱ	2.0675 (15)
Ni1A—N4A ⁱ	2.1190 (15)	Ni2B—N2B	2.0675 (15)
Ni1A—N4A	2.1190 (15)	Ni2B—N1B ⁱⁱ	2.0719 (16)
S1A—C9A	1.6339 (17)	Ni2B—N1B	2.0719 (16)
N1A—C1A	1.478 (2)	Ni2B—N4B ⁱⁱ	2.0933 (16)
N1A—C2A	1.486 (2)	Ni2B—N4B	2.0933 (16)
N1A—H1A	1.0000	S1B—C9B	1.6190 (18)
N2A—C4A	1.477 (2)	N1B—C1B	1.479 (2)
N2A—C3A	1.483 (3)	N1B—C2B	1.484 (2)
N2A—H2A	1.0000	N1B—H1B	1.0000
N3A—C3A	1.436 (3)	N2B—C4B	1.480 (2)
N3A—C2A	1.440 (3)	N2B—C3B	1.486 (2)
N3A—C5A	1.470 (3)	N2B—H2B	1.0000
N4A—C9A	1.158 (2)	N3B—C3B	1.444 (3)
C1A—C4A ⁱ	1.517 (3)	N3B—C2B	1.446 (2)

C1A—H1A1	0.9900	N3B—C5B	1.469 (3)
C1A—H1A2	0.9900	N4B—C9B	1.156 (2)
C2A—H2A1	0.9900	C1B—C4B ⁱⁱ	1.523 (3)
C2A—H2A2	0.9900	C1B—H1B1	0.9900
C3A—H3A1	0.9900	C1B—H1B2	0.9900
C3A—H3A2	0.9900	C2B—H2B1	0.9900
C4A—C1A ⁱ	1.517 (3)	C2B—H2B2	0.9900
C4A—H4A1	0.9900	C3B—H3B1	0.9900
C4A—H4A2	0.9900	C3B—H3B2	0.9900
C5A—C6A	1.501 (4)	C4B—C1B ⁱⁱ	1.522 (3)
C5A—H5A1	0.9900	C4B—H4B1	0.9900
C5A—H5A2	0.9900	C4B—H4B2	0.9900
C6A—C72A	1.537 (9)	C5B—C6B	1.520 (3)
C6A—C71A	1.641 (6)	C5B—H5B1	0.9900
C6A—H6A1	0.9900	C5B—H5B2	0.9900
C6A—H6A2	0.9900	C6B—C7B	1.514 (3)
C71A—C81A	1.485 (5)	C6B—H6B1	0.9900
C71A—H71A	0.9900	C6B—H6B2	0.9900
C71A—H71B	0.9900	C7B—C8B	1.522 (3)
C81A—H81A	0.9800	C7B—H7B1	0.9900
C81A—H81B	0.9800	C7B—H7B2	0.9900
C81A—H81C	0.9800	C8B—H8B1	0.9800
C72A—C82A	1.427 (12)	C8B—H8B2	0.9800
C72A—H72A	0.9900	C8B—H8B3	0.9800
C72A—H72B	0.9900		
N1A ⁱ —Ni1A—N1A	180.00 (7)	C72A—C82A—H82B	109.5
N1A ⁱ —Ni1A—N2A ⁱ	95.00 (7)	H82A—C82A—H82B	109.5
N1A—Ni1A—N2A ⁱ	85.00 (6)	C72A—C82A—H82C	109.5
N1A ⁱ —Ni1A—N2A	85.00 (6)	H82A—C82A—H82C	109.5
N1A—Ni1A—N2A	95.00 (6)	H82B—C82A—H82C	109.5
N2A ⁱ —Ni1A—N2A	180.00 (8)	N4A—C9A—S1A	178.09 (16)
N1A ⁱ —Ni1A—N4A ⁱ	91.75 (6)	N2B ⁱⁱ —Ni2B—N2B	180.0
N1A—Ni1A—N4A ⁱ	88.25 (6)	N2B ⁱⁱ —Ni2B—N1B ⁱⁱ	93.91 (6)
N2A ⁱ —Ni1A—N4A ⁱ	92.85 (6)	N2B—Ni2B—N1B ⁱⁱ	86.09 (6)
N2A—Ni1A—N4A ⁱ	87.15 (6)	N2B ⁱⁱ —Ni2B—N1B	86.09 (6)
N1A ⁱ —Ni1A—N4A	88.25 (6)	N2B—Ni2B—N1B	93.91 (6)
N1A—Ni1A—N4A	91.75 (6)	N1B ⁱⁱ —Ni2B—N1B	180.0
N2A ⁱ —Ni1A—N4A	87.15 (6)	N2B ⁱⁱ —Ni2B—N4B ⁱⁱ	88.26 (6)
N2A—Ni1A—N4A	92.85 (6)	N2B—Ni2B—N4B ⁱⁱ	91.74 (6)
N4A ⁱ —Ni1A—N4A	180.0	N1B ⁱⁱ —Ni2B—N4B ⁱⁱ	88.42 (7)
C1A—N1A—C2A	114.56 (15)	N1B—Ni2B—N4B ⁱⁱ	91.58 (7)
C1A—N1A—Ni1A	106.14 (11)	N2B ⁱⁱ —Ni2B—N4B	91.74 (6)
C2A—N1A—Ni1A	112.51 (12)	N2B—Ni2B—N4B	88.26 (6)
C1A—N1A—H1A	107.8	N1B ⁱⁱ —Ni2B—N4B	91.58 (7)
C2A—N1A—H1A	107.8	N1B—Ni2B—N4B	88.42 (7)
Ni1A—N1A—H1A	107.8	N4B ⁱⁱ —Ni2B—N4B	180.0
C4A—N2A—C3A	115.13 (15)	C1B—N1B—C2B	114.04 (15)

C4A—N2A—Ni1A	105.93 (11)	C1B—N1B—Ni2B	104.88 (11)
C3A—N2A—Ni1A	112.70 (12)	C2B—N1B—Ni2B	113.56 (11)
C4A—N2A—H2A	107.6	C1B—N1B—H1B	108.0
C3A—N2A—H2A	107.6	C2B—N1B—H1B	108.0
Ni1A—N2A—H2A	107.6	Ni2B—N1B—H1B	108.0
C3A—N3A—C2A	116.46 (17)	C4B—N2B—C3B	114.37 (14)
C3A—N3A—C5A	113.3 (2)	C4B—N2B—Ni2B	104.95 (10)
C2A—N3A—C5A	116.6 (2)	C3B—N2B—Ni2B	112.98 (11)
C9A—N4A—Ni1A	161.18 (15)	C4B—N2B—H2B	108.1
N1A—C1A—C4A ⁱ	108.32 (14)	C3B—N2B—H2B	108.1
N1A—C1A—H1A1	110.0	Ni2B—N2B—H2B	108.1
C4A ⁱ —C1A—H1A1	110.0	C3B—N3B—C2B	115.91 (15)
N1A—C1A—H1A2	110.0	C3B—N3B—C5B	115.92 (16)
C4A ⁱ —C1A—H1A2	110.0	C2B—N3B—C5B	113.83 (16)
H1A1—C1A—H1A2	108.4	C9B—N4B—Ni2B	163.23 (16)
N3A—C2A—N1A	113.64 (16)	N1B—C1B—C4B ⁱⁱ	108.49 (15)
N3A—C2A—H2A1	108.8	N1B—C1B—H1B1	110.0
N1A—C2A—H2A1	108.8	C4B ⁱⁱ —C1B—H1B1	110.0
N3A—C2A—H2A2	108.8	N1B—C1B—H1B2	110.0
N1A—C2A—H2A2	108.8	C4B ⁱⁱ —C1B—H1B2	110.0
H2A1—C2A—H2A2	107.7	H1B1—C1B—H1B2	108.4
N3A—C3A—N2A	113.85 (16)	N3B—C2B—N1B	113.94 (15)
N3A—C3A—H3A1	108.8	N3B—C2B—H2B1	108.8
N2A—C3A—H3A1	108.8	N1B—C2B—H2B1	108.8
N3A—C3A—H3A2	108.8	N3B—C2B—H2B2	108.8
N2A—C3A—H3A2	108.8	N1B—C2B—H2B2	108.8
H3A1—C3A—H3A2	107.7	H2B1—C2B—H2B2	107.7
N2A—C4A—C1A ⁱ	108.23 (15)	N3B—C3B—N2B	114.19 (14)
N2A—C4A—H4A1	110.1	N3B—C3B—H3B1	108.7
C1A ⁱ —C4A—H4A1	110.1	N2B—C3B—H3B1	108.7
N2A—C4A—H4A2	110.1	N3B—C3B—H3B2	108.7
C1A ⁱ —C4A—H4A2	110.1	N2B—C3B—H3B2	108.7
H4A1—C4A—H4A2	108.4	H3B1—C3B—H3B2	107.6
N3A—C5A—C6A	115.5 (3)	N2B—C4B—C1B ⁱⁱ	108.66 (15)
N3A—C5A—H5A1	108.4	N2B—C4B—H4B1	110.0
C6A—C5A—H5A1	108.4	C1B ⁱⁱ —C4B—H4B1	110.0
N3A—C5A—H5A2	108.4	N2B—C4B—H4B2	110.0
C6A—C5A—H5A2	108.4	C1B ⁱⁱ —C4B—H4B2	110.0
H5A1—C5A—H5A2	107.5	H4B1—C4B—H4B2	108.3
C5A—C6A—C72A	125.6 (5)	N3B—C5B—C6B	113.59 (16)
C5A—C6A—C71A	105.5 (3)	N3B—C5B—H5B1	108.8
C5A—C6A—H6A1	110.6	C6B—C5B—H5B1	108.8
C71A—C6A—H6A1	110.6	N3B—C5B—H5B2	108.8
C5A—C6A—H6A2	110.6	C6B—C5B—H5B2	108.8
C71A—C6A—H6A2	110.6	H5B1—C5B—H5B2	107.7
H6A1—C6A—H6A2	108.8	C7B—C6B—C5B	112.38 (16)
C81A—C71A—C6A	107.8 (4)	C7B—C6B—H6B1	109.1
C81A—C71A—H71A	110.2	C5B—C6B—H6B1	109.1

C6A—C71A—H71A	110.2	C7B—C6B—H6B2	109.1
C81A—C71A—H71B	110.2	C5B—C6B—H6B2	109.1
C6A—C71A—H71B	110.2	H6B1—C6B—H6B2	107.9
H71A—C71A—H71B	108.5	C6B—C7B—C8B	112.29 (17)
C71A—C81A—H81A	109.5	C6B—C7B—H7B1	109.1
C71A—C81A—H81B	109.5	C8B—C7B—H7B1	109.1
H81A—C81A—H81B	109.5	C6B—C7B—H7B2	109.1
C71A—C81A—H81C	109.5	C8B—C7B—H7B2	109.1
H81A—C81A—H81C	109.5	H7B1—C7B—H7B2	107.9
H81B—C81A—H81C	109.5	C7B—C8B—H8B1	109.5
C82A—C72A—C6A	99.0 (7)	C7B—C8B—H8B2	109.5
C82A—C72A—H72A	112.0	H8B1—C8B—H8B2	109.5
C6A—C72A—H72A	112.0	C7B—C8B—H8B3	109.5
C82A—C72A—H72B	112.0	H8B1—C8B—H8B3	109.5
C6A—C72A—H72B	112.0	H8B2—C8B—H8B3	109.5
H72A—C72A—H72B	109.6	N4B—C9B—S1B	178.44 (17)
C72A—C82A—H82A	109.5		
C2A—N1A—C1A—C4A ⁱ	167.05 (14)	C5A—C6A—C72A—C82A	71.8 (8)
Ni1A—N1A—C1A—C4A ⁱ	42.27 (15)	C2B—N1B—C1B—C4B ⁱⁱ	-167.20 (15)
C3A—N3A—C2A—N1A	73.7 (2)	Ni2B—N1B—C1B—C4B ⁱⁱ	-42.36 (16)
C5A—N3A—C2A—N1A	-64.4 (2)	C3B—N3B—C2B—N1B	-71.3 (2)
C1A—N1A—C2A—N3A	-178.39 (15)	C5B—N3B—C2B—N1B	66.9 (2)
Ni1A—N1A—C2A—N3A	-57.03 (18)	C1B—N1B—C2B—N3B	176.89 (15)
C2A—N3A—C3A—N2A	-73.1 (2)	Ni2B—N1B—C2B—N3B	56.80 (19)
C5A—N3A—C3A—N2A	66.4 (2)	C2B—N3B—C3B—N2B	72.1 (2)
C4A—N2A—C3A—N3A	177.61 (16)	C5B—N3B—C3B—N2B	-65.2 (2)
Ni1A—N2A—C3A—N3A	55.96 (19)	C4B—N2B—C3B—N3B	-177.77 (14)
C3A—N2A—C4A—C1A ⁱ	-167.20 (15)	Ni2B—N2B—C3B—N3B	-57.79 (17)
Ni1A—N2A—C4A—C1A ⁱ	-41.95 (15)	C3B—N2B—C4B—C1B ⁱⁱ	166.62 (14)
C3A—N3A—C5A—C6A	166.0 (2)	Ni2B—N2B—C4B—C1B ⁱⁱ	42.25 (15)
C2A—N3A—C5A—C6A	-54.7 (3)	C3B—N3B—C5B—C6B	-58.2 (2)
N3A—C5A—C6A—C72A	159.1 (5)	C2B—N3B—C5B—C6B	163.62 (16)
N3A—C5A—C6A—C71A	-173.6 (3)	N3B—C5B—C6B—C7B	-173.67 (16)
C5A—C6A—C71A—C81A	175.0 (4)	C5B—C6B—C7B—C8B	-171.39 (18)

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+2, -y+1, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1A—H1A \cdots S1A ⁱⁱⁱ	1.00	2.73	3.5154 (17)	136
N2B—H2B \cdots S1B ^{iv}	1.00	2.66	3.4556 (17)	137

Symmetry codes: (iii) $x+1, y, z$; (iv) $-x+1, -y+1, -z$.