



Crystal structure of benzobicyclon

Gihaeng Kang, Jineun Kim,* Hansu Lim and Tae Ho Kim*

Department of Chemistry and Research Institute of Natural Sciences, Gyeongsang National University, Jinju 52828, Republic of Korea. *Correspondence e-mail: thkim@gnu.ac.kr, jekim@gnu.ac.kr

Received 27 November 2015; accepted 2 December 2015

Edited by J. Simpson, University of Otago, New Zealand


In the title compound, $C_{22}H_{19}ClO_4S_2$ [systematic name: 3-(2-chloro-4-mesylbenzoyl)-4-(phenylsulfanyl)bicyclo[3.2.1]oct-2-en-2-one], which is an unclassified herbicide, the dihedral angle between the plane of the phenyl and chlorobenzene rings is $19.9(2)^\circ$. In the crystal, $C-H \cdots O$ hydrogen bonds link adjacent molecules, generating two-dimensional networks extending parallel to (011).

Keywords: crystal structure; benzobicyclon; bicyclo[3.2.1]oct-2-en-4-one; herbicide.

CCDC reference: 1440215

1. Related literature

For information on the herbicidal properties of the title compound, see: Im *et al.* (2015). For related crystal structures, see: Brown *et al.* (2007); Hou *et al.* (2010).



2. Experimental

2.1. Crystal data

$C_{22}H_{19}ClO_4S_2$
 $M_r = 446.94$

Monoclinic, $P2_1/c$
 $a = 15.5111(8) \text{ \AA}$

$b = 10.7218(6) \text{ \AA}$
 $c = 12.0169(6) \text{ \AA}$
 $\beta = 91.449(3)^\circ$
 $V = 1997.85(18) \text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.43 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
 $0.15 \times 0.11 \times 0.06 \text{ mm}$

2.2. Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2014)
 $T_{\min} = 0.687$, $T_{\max} = 0.746$

15850 measured reflections
3504 independent reflections
2584 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.142$
 $S = 1.07$
3504 reflections

263 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.97 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C1-H1B \cdots O4^i$	0.98	2.44	3.322 (4)	150
$C1-H1C \cdots O4^{ii}$	0.98	2.52	3.477 (5)	165
$C3-H3 \cdots O2^{iii}$	0.95	2.53	3.422 (4)	157

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x, -y + 1, -z + 2$.

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINT* (Bruker, 2014); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Acknowledgements

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2015R1D1A4A01020317).

Supporting information for this paper is available from the IUCr electronic archives (Reference: SJ5485).

References

- Brandenburg, K. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Brown, P. M., Käppel, N., Murphy, P. J., Coles, S. J. & Hursthouse, M. B. (2007). *Tetrahedron*, **63**, 1100–1106.
Bruker (2014). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Hou, Y.-J., Chu, W.-Y., Sui, J. & Sun, Z.-Z. (2010). *Z. Kristallogr.* **225**, 465–466.
Im, S. J., Abd El-Aty, A. M., Lee, Y.-J., Rahman, M. M., Kim, S.-W., Choi, J.-H. & Shim, J.-H. (2015). *Food Chem.* **168**, 404–409.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.

supporting information

Acta Cryst. (2015). E71, o1035 [https://doi.org/10.1107/S2056989015023221]

Crystal structure of benzobicyclon

Gihaeng Kang, Jineun Kim, Hansu Lim and Tae Ho Kim

S1. Comment

Benzobicyclon is a newly developed compound with potent herbicidal activity against *Scirpus juncooides*, one of the most troublesome weeds in paddy fields. (Im *et al.*, 2015). However, until now its crystal structure has not been reported. In the title compound (Fig. 1), the dihedral angle between the plane of the phenyl and chlorobenzene rings is 19.9 (2)°. All bond lengths and bond angles are normal and comparable to those observed in the crystal structures of similar compounds (Brown *et al.*, 2007; Hou *et al.*, 2010).

In the crystal structure (Fig. 2), intermolecular C1—H1B···O4 hydrogen bond link adjacent molecules, forming chains along the *c*-axis direction. These chains are linked by C1—H1C···O4 and C3—H3···O2 hydrogen bonds (Table 1), resulting in two-dimensional networks extending parallel to the (011) plane.

S2. Experimental

The title compound was purchased from the Dr. Ehrenstorfer GmbH Company. Slow evaporation of a solution in CH₂Cl₂ gave single crystals suitable for X-ray analysis.

S3. Refinement

All H-atoms were positioned geometrically and refined using a riding model with $d(\text{C—H}) = 1.00 \text{ \AA}$, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for Csp³—H, $d(\text{C—H}) = 0.99 \text{ \AA}$, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for CH₂ groups, $d(\text{C—H}) = 0.98 \text{ \AA}$, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for CH₃ groups, $d(\text{C—H}) = 0.95 \text{ \AA}$, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ for aromatic C—H.




Figure 1

The asymmetric unit of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are shown as small spheres of arbitrary radius.




Figure 2

Crystal packing viewed along the c axis. The intermolecular hydrogen bonds are shown as dashed lines.

3-(2-Chloro-4-mesitylbenzoyl)-4-(phenylsulfanyl)bicyclo[3.2.1]oct-3-en-2-one

Crystal data

$C_{22}H_{19}ClO_4S_2$

$M_r = 446.94$

Monoclinic, $P2_1/c$

$a = 15.5111(8) \text{ \AA}$

$b = 10.7218(6) \text{ \AA}$

$c = 12.0169(6) \text{ \AA}$

$\beta = 91.449(3)^\circ$

$V = 1997.85(18) \text{ \AA}^3$

$Z = 4$

$F(000) = 928$

$D_x = 1.486 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2645 reflections

$\theta = 2.6\text{--}24.7^\circ$

$\mu = 0.43 \text{ mm}^{-1}$

$T = 173 \text{ K}$

Block, colourless

$0.15 \times 0.11 \times 0.06 \text{ mm}$

Data collection

Bruker APEXII CCD

diffractometer

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2014)

$T_{\min} = 0.687$, $T_{\max} = 0.746$

15850 measured reflections

3504 independent reflections

2584 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.055$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -18 \rightarrow 18$

$k = -11 \rightarrow 12$

$l = -14 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.053$

$wR(F^2) = 0.142$

$S = 1.07$

3504 reflections

263 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0619P)^2 + 2.0045P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.97 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.22882 (6)	0.40049 (9)	0.83242 (9)	0.0464 (3)
S1	-0.08667 (6)	0.26508 (9)	0.95779 (9)	0.0361 (3)
S2	0.41445 (6)	0.12501 (8)	0.64532 (8)	0.0324 (3)
O1	-0.16416 (16)	0.2346 (3)	0.8963 (2)	0.0495 (8)
O2	-0.07793 (17)	0.3867 (2)	1.0065 (3)	0.0499 (8)
O3	0.24478 (16)	0.0592 (2)	0.6713 (2)	0.0429 (7)
O4	0.14394 (16)	0.3801 (3)	0.5647 (3)	0.0479 (8)
C1	-0.0691 (2)	0.1524 (4)	1.0616 (3)	0.0401 (10)
H1A	-0.1083	0.1674	1.1228	0.060*
H1B	-0.0093	0.1576	1.0895	0.060*
H1C	-0.0799	0.0692	1.0304	0.060*
C2	0.0015 (2)	0.2413 (3)	0.8691 (3)	0.0303 (8)
C3	0.0716 (2)	0.3211 (3)	0.8792 (3)	0.0339 (9)
H3	0.0715	0.3884	0.9305	0.041*
C4	0.1417 (2)	0.3002 (3)	0.8126 (3)	0.0309 (8)
C5	0.1442 (2)	0.2011 (3)	0.7389 (3)	0.0290 (8)
C6	0.0730 (2)	0.1218 (3)	0.7314 (3)	0.0335 (9)
H6	0.0736	0.0530	0.6816	0.040*
C7	0.0013 (2)	0.1420 (3)	0.7954 (3)	0.0338 (9)
H7	-0.0473	0.0883	0.7890	0.041*
C8	0.2214 (2)	0.1671 (3)	0.6715 (3)	0.0302 (8)
C9	0.2696 (2)	0.2642 (3)	0.6103 (3)	0.0336 (9)
C10	0.3565 (2)	0.2546 (3)	0.5982 (3)	0.0310 (8)
C11	0.4045 (2)	0.3595 (4)	0.5419 (3)	0.0382 (9)
H11	0.4656	0.3361	0.5274	0.046*
C12	0.3542 (4)	0.3935 (5)	0.4330 (3)	0.0668 (16)
H12A	0.3347	0.3191	0.3906	0.080*
H12B	0.3870	0.4499	0.3845	0.080*
C13	0.2767 (3)	0.4631 (4)	0.4939 (5)	0.0711 (17)
H13	0.2407	0.5144	0.4411	0.085*
C14	0.3207 (4)	0.5376 (5)	0.5754 (5)	0.0755 (17)
H14A	0.2838	0.5499	0.6405	0.091*
H14B	0.3341	0.6204	0.5438	0.091*
C15	0.3992 (3)	0.4753 (4)	0.6092 (4)	0.0622 (14)
H15A	0.3983	0.4548	0.6895	0.075*
H15B	0.4495	0.5295	0.5958	0.075*
C16	0.2224 (2)	0.3702 (4)	0.5594 (4)	0.0466 (11)
C17	0.5234 (2)	0.1795 (3)	0.6555 (3)	0.0299 (8)
C18	0.5496 (3)	0.2579 (3)	0.7411 (3)	0.0404 (10)

H18	0.5085	0.2892	0.7914	0.048*
C19	0.6355 (3)	0.2910 (4)	0.7538 (4)	0.0532 (12)
H19	0.6536	0.3460	0.8118	0.064*
C20	0.6948 (3)	0.2430 (4)	0.6807 (4)	0.0566 (13)
H20	0.7540	0.2643	0.6896	0.068*
C21	0.6690 (3)	0.1654 (4)	0.5962 (4)	0.0488 (12)
H21	0.7103	0.1336	0.5465	0.059*
C22	0.5831 (2)	0.1329 (3)	0.5825 (3)	0.0374 (9)
H22	0.5652	0.0791	0.5236	0.045*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0325 (5)	0.0437 (6)	0.0632 (7)	-0.0172 (4)	0.0084 (5)	-0.0109 (5)
S1	0.0220 (5)	0.0359 (5)	0.0509 (6)	-0.0007 (4)	0.0085 (4)	-0.0012 (4)
S2	0.0273 (5)	0.0273 (5)	0.0430 (6)	0.0021 (4)	0.0086 (4)	0.0007 (4)
O1	0.0215 (13)	0.0641 (19)	0.063 (2)	0.0002 (12)	-0.0001 (13)	-0.0002 (15)
O2	0.0421 (16)	0.0354 (15)	0.073 (2)	-0.0006 (12)	0.0196 (15)	-0.0119 (14)
O3	0.0372 (15)	0.0298 (14)	0.0624 (19)	-0.0013 (12)	0.0170 (14)	-0.0004 (13)
O4	0.0221 (14)	0.0543 (17)	0.067 (2)	0.0025 (12)	0.0014 (13)	0.0214 (15)
C1	0.030 (2)	0.046 (2)	0.045 (2)	-0.0030 (17)	0.0105 (18)	0.0022 (19)
C2	0.0229 (18)	0.0305 (18)	0.038 (2)	-0.0010 (14)	0.0032 (16)	0.0033 (16)
C3	0.0296 (19)	0.0289 (18)	0.043 (2)	-0.0007 (15)	0.0015 (17)	-0.0017 (17)
C4	0.0235 (17)	0.0294 (18)	0.040 (2)	-0.0053 (15)	0.0033 (16)	0.0032 (16)
C5	0.0254 (18)	0.0278 (18)	0.034 (2)	0.0000 (14)	0.0047 (16)	0.0064 (16)
C6	0.032 (2)	0.0277 (18)	0.041 (2)	-0.0078 (15)	0.0045 (17)	-0.0026 (17)
C7	0.0230 (18)	0.0325 (19)	0.046 (2)	-0.0066 (15)	0.0019 (17)	0.0022 (17)
C8	0.0258 (18)	0.0282 (19)	0.037 (2)	-0.0045 (15)	0.0044 (16)	0.0028 (16)
C9	0.0239 (18)	0.036 (2)	0.041 (2)	0.0005 (15)	0.0052 (16)	0.0093 (17)
C10	0.0284 (19)	0.0329 (19)	0.032 (2)	0.0003 (15)	0.0050 (16)	0.0034 (16)
C11	0.0225 (18)	0.044 (2)	0.049 (2)	0.0000 (16)	0.0069 (17)	0.0151 (19)
C12	0.107 (4)	0.068 (3)	0.026 (2)	-0.048 (3)	0.012 (3)	0.002 (2)
C13	0.026 (2)	0.044 (3)	0.141 (5)	0.0005 (19)	-0.023 (3)	0.043 (3)
C14	0.095 (4)	0.061 (3)	0.072 (4)	0.008 (3)	0.031 (3)	0.018 (3)
C15	0.094 (4)	0.047 (3)	0.046 (3)	-0.030 (3)	0.011 (3)	0.000 (2)
C16	0.029 (2)	0.048 (2)	0.063 (3)	0.0016 (18)	0.005 (2)	0.020 (2)
C17	0.0255 (18)	0.0270 (18)	0.037 (2)	0.0048 (14)	0.0025 (16)	0.0080 (16)
C18	0.040 (2)	0.037 (2)	0.044 (2)	0.0001 (17)	0.0027 (19)	0.0038 (19)
C19	0.049 (3)	0.055 (3)	0.055 (3)	-0.011 (2)	-0.012 (2)	0.009 (2)
C20	0.027 (2)	0.068 (3)	0.074 (3)	-0.009 (2)	-0.007 (2)	0.029 (3)
C21	0.033 (2)	0.056 (3)	0.058 (3)	0.016 (2)	0.013 (2)	0.028 (2)
C22	0.031 (2)	0.038 (2)	0.044 (2)	0.0097 (17)	0.0096 (18)	0.0064 (18)

Geometric parameters (Å, °)

C11—C4	1.739 (3)	C11—C15	1.485 (6)
S1—O1	1.432 (3)	C11—C12	1.551 (6)
S1—O2	1.434 (3)	C11—H11	1.0000

S1—C1	1.753 (4)	C12—C13	1.607 (7)
S1—C2	1.774 (3)	C12—H12A	0.9900
S2—C10	1.742 (4)	C12—H12B	0.9900
S2—C17	1.790 (4)	C13—C14	1.425 (8)
O3—C8	1.213 (4)	C13—C16	1.534 (6)
O4—C16	1.226 (4)	C13—H13	1.0000
C1—H1A	0.9800	C14—C15	1.437 (7)
C1—H1B	0.9800	C14—H14A	0.9900
C1—H1C	0.9800	C14—H14B	0.9900
C2—C7	1.385 (5)	C15—H15A	0.9900
C2—C3	1.386 (5)	C15—H15B	0.9900
C3—C4	1.384 (5)	C17—C18	1.382 (5)
C3—H3	0.9500	C17—C22	1.385 (5)
C4—C5	1.385 (5)	C18—C19	1.383 (6)
C5—C6	1.395 (5)	C18—H18	0.9500
C5—C8	1.508 (5)	C19—C20	1.387 (6)
C6—C7	1.384 (5)	C19—H19	0.9500
C6—H6	0.9500	C20—C21	1.365 (7)
C7—H7	0.9500	C20—H20	0.9500
C8—C9	1.487 (5)	C21—C22	1.383 (6)
C9—C10	1.363 (5)	C21—H21	0.9500
C9—C16	1.476 (5)	C22—H22	0.9500
C10—C11	1.517 (5)		
O1—S1—O2	119.16 (18)	C11—C12—C13	95.2 (3)
O1—S1—C1	108.84 (18)	C11—C12—H12A	112.7
O2—S1—C1	108.9 (2)	C13—C12—H12A	112.7
O1—S1—C2	107.89 (18)	C11—C12—H12B	112.7
O2—S1—C2	108.04 (16)	C13—C12—H12B	112.7
C1—S1—C2	102.72 (17)	H12A—C12—H12B	110.2
C10—S2—C17	103.94 (16)	C14—C13—C16	105.7 (5)
S1—C1—H1A	109.5	C14—C13—C12	102.8 (4)
S1—C1—H1B	109.5	C16—C13—C12	111.0 (3)
H1A—C1—H1B	109.5	C14—C13—H13	112.2
S1—C1—H1C	109.5	C16—C13—H13	112.2
H1A—C1—H1C	109.5	C12—C13—H13	112.2
H1B—C1—H1C	109.5	C13—C14—C15	108.8 (4)
C7—C2—C3	121.3 (3)	C13—C14—H14A	109.9
C7—C2—S1	120.4 (3)	C15—C14—H14A	109.9
C3—C2—S1	118.3 (3)	C13—C14—H14B	109.9
C4—C3—C2	118.4 (3)	C15—C14—H14B	109.9
C4—C3—H3	120.8	H14A—C14—H14B	108.3
C2—C3—H3	120.8	C14—C15—C11	107.1 (4)
C3—C4—C5	121.9 (3)	C14—C15—H15A	110.3
C3—C4—C11	116.1 (3)	C11—C15—H15A	110.3
C5—C4—C11	121.8 (3)	C14—C15—H15B	110.3
C4—C5—C6	118.3 (3)	C11—C15—H15B	110.3
C4—C5—C8	124.4 (3)	H15A—C15—H15B	108.6

C6—C5—C8	117.2 (3)	O4—C16—C9	121.8 (3)
C7—C6—C5	120.9 (3)	O4—C16—C13	121.9 (4)
C7—C6—H6	119.5	C9—C16—C13	116.1 (3)
C5—C6—H6	119.5	C18—C17—C22	120.2 (3)
C6—C7—C2	119.1 (3)	C18—C17—S2	120.6 (3)
C6—C7—H7	120.4	C22—C17—S2	119.0 (3)
C2—C7—H7	120.4	C17—C18—C19	120.1 (4)
O3—C8—C9	120.9 (3)	C17—C18—H18	119.9
O3—C8—C5	118.2 (3)	C19—C18—H18	119.9
C9—C8—C5	120.9 (3)	C18—C19—C20	119.2 (4)
C10—C9—C16	119.7 (3)	C18—C19—H19	120.4
C10—C9—C8	120.7 (3)	C20—C19—H19	120.4
C16—C9—C8	119.6 (3)	C21—C20—C19	120.8 (4)
C9—C10—C11	119.4 (3)	C21—C20—H20	119.6
C9—C10—S2	121.9 (3)	C19—C20—H20	119.6
C11—C10—S2	118.7 (3)	C20—C21—C22	120.3 (4)
C15—C11—C10	110.0 (3)	C20—C21—H21	119.9
C15—C11—C12	103.2 (4)	C22—C21—H21	119.9
C10—C11—C12	107.9 (3)	C21—C22—C17	119.5 (4)
C15—C11—H11	111.8	C21—C22—H22	120.3
C10—C11—H11	111.8	C17—C22—H22	120.3
C12—C11—H11	111.8		
O1—S1—C2—C7	-37.9 (4)	C17—S2—C10—C11	19.8 (3)
O2—S1—C2—C7	-167.9 (3)	C9—C10—C11—C15	64.1 (5)
C1—S1—C2—C7	77.0 (3)	S2—C10—C11—C15	-116.0 (4)
O1—S1—C2—C3	145.0 (3)	C9—C10—C11—C12	-47.8 (5)
O2—S1—C2—C3	14.9 (4)	S2—C10—C11—C12	132.1 (3)
C1—S1—C2—C3	-100.1 (3)	C15—C11—C12—C13	-42.9 (4)
C7—C2—C3—C4	0.8 (6)	C10—C11—C12—C13	73.5 (4)
S1—C2—C3—C4	177.9 (3)	C11—C12—C13—C14	44.6 (4)
C2—C3—C4—C5	-1.5 (6)	C11—C12—C13—C16	-68.0 (5)
C2—C3—C4—C11	-178.0 (3)	C16—C13—C14—C15	86.1 (5)
C3—C4—C5—C6	0.9 (6)	C12—C13—C14—C15	-30.4 (5)
C11—C4—C5—C6	177.3 (3)	C13—C14—C15—C11	2.1 (5)
C3—C4—C5—C8	-174.6 (3)	C10—C11—C15—C14	-86.7 (4)
C11—C4—C5—C8	1.8 (5)	C12—C11—C15—C14	28.3 (4)
C4—C5—C6—C7	0.4 (6)	C10—C9—C16—O4	177.3 (4)
C8—C5—C6—C7	176.2 (3)	C8—C9—C16—O4	-1.1 (7)
C5—C6—C7—C2	-1.0 (6)	C10—C9—C16—C13	0.7 (6)
C3—C2—C7—C6	0.5 (6)	C8—C9—C16—C13	-177.6 (4)
S1—C2—C7—C6	-176.6 (3)	C14—C13—C16—O4	106.6 (5)
C4—C5—C8—O3	131.3 (4)	C12—C13—C16—O4	-142.6 (5)
C6—C5—C8—O3	-44.2 (5)	C14—C13—C16—C9	-76.9 (5)
C4—C5—C8—C9	-46.5 (5)	C12—C13—C16—C9	33.9 (6)
C6—C5—C8—C9	138.0 (4)	C10—S2—C17—C18	74.3 (3)
O3—C8—C9—C10	-31.3 (6)	C10—S2—C17—C22	-111.6 (3)
C5—C8—C9—C10	146.4 (4)	C22—C17—C18—C19	0.5 (6)

O3—C8—C9—C16	147.0 (4)	S2—C17—C18—C19	174.5 (3)
C5—C8—C9—C16	-35.3 (5)	C17—C18—C19—C20	-1.0 (6)
C16—C9—C10—C11	5.7 (6)	C18—C19—C20—C21	1.0 (7)
C8—C9—C10—C11	-176.0 (3)	C19—C20—C21—C22	-0.4 (6)
C16—C9—C10—S2	-174.2 (3)	C20—C21—C22—C17	-0.1 (6)
C8—C9—C10—S2	4.1 (5)	C18—C17—C22—C21	0.1 (5)
C17—S2—C10—C9	-160.3 (3)	S2—C17—C22—C21	-174.0 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C1—H1B \cdots O4 ⁱ	0.98	2.44	3.322 (4)	150
C1—H1C \cdots O4 ⁱⁱ	0.98	2.52	3.477 (5)	165
C3—H3 \cdots O2 ⁱⁱⁱ	0.95	2.53	3.422 (4)	157

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x, y-1/2, -z+3/2$; (iii) $-x, -y+1, -z+2$.