



Crystal structure of *trans*-diammine(1,4,8,11-tetraazacyclotetradecane- κ^4N)chromium(III) tetrachloridozincate chloride monohydrate from synchrotron data

Dohyun Moon^a and Jong-Ha Choi^{b*}

Received 4 February 2016

Accepted 1 March 2016

^aPohang Accelerator Laboratory, POSTECH, Pohang 37673, Republic of Korea, and ^bDepartment of Chemistry, Andong National University, Andong 36729, Republic of Korea. *Correspondence e-mail: jhchoi@anu.ac.kr

Edited by M. Weil, Vienna University of Technology, Austria

Keywords: crystal structure; cyclam; ammine ligand; tetrachloridozincate chloride double salt; *trans*-III conformation; chromium(III) complex.


CCDC reference: 1456673

Supporting information: this article has supporting information at journals.iucr.org/e


The asymmetric unit of the title complex salt, $[\text{Cr}(\text{C}_{10}\text{H}_{24}\text{N}_4)(\text{NH}_3)_2][\text{ZnCl}_4]\text{Cl}\cdot\text{H}_2\text{O}$, is comprised of four halves of the Cr^{III} complex cations (the counterparts being generated by application of inversion symmetry), two tetrachloridozincate anions, two chloride anions and two water molecules. Each Cr^{III} ion is coordinated by the four N atoms of the cyclam (1,4,8,11-tetraazacyclotetradecane) ligand in the equatorial plane and by two N atoms of ammine ligands in axial positions, displaying an overall distorted octahedral coordination environment. The $\text{Cr}-\text{N}(\text{cyclam})$ bond lengths range from 2.0501 (15) to 2.0615 (15) Å, while the $\text{Cr}-\text{N}(\text{NH}_3)$ bond lengths range from 2.0976 (13) to 2.1062 (13) Å. The macrocyclic cyclam moieties adopt the *trans*-III conformation with six- and five-membered chelate rings in chair and *gauche* conformations. The $[\text{ZnCl}_4]^{2-}$ anions have a slightly distorted tetrahedral shape. In the crystal, the Cl^- anions link the complex cations, as well as the solvent water molecules, through $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen-bonding interactions. The supramolecular set-up also includes $\text{N}-\text{H}\cdots\text{Cl}$, $\text{C}-\text{H}\cdots\text{Cl}$, $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonding between $\text{N}-\text{H}$ or $\text{C}-\text{H}$ groups of cyclam, ammine $\text{N}-\text{H}$ and water $\text{O}-\text{H}$ donor groups, and O atoms of the water molecules, Cl^- anions or Cl atoms of the $[\text{ZnCl}_4]^{2-}$ anions as acceptors, leading to a three-dimensional network structure.

1. Chemical context

The cyclam macrocycle (1,4,8,11-tetraazacyclotetradecane, $\text{C}_{10}\text{H}_{24}\text{N}_4$) can adopt both planar (*trans*) and folded (*cis*) configurations (Poon & Pun, 1980). There are five conformational *trans* isomers for the macrocycle, which differ in the chirality of the *sec*-NH groups (Choi, 2009). The *trans*-I, *trans*-II and *trans*-V conformations can fold to form *cis*-I, *cis*-II and *cis*-V isomers, respectively (Subhan *et al.*, 2011). Recently, it has been reported that cyclam derivatives and their metal complexes exhibit *anti*-HIV activity (Ronconi & Sadler, 2007; De Clercq, 2010; Ross *et al.*, 2012) whereby the strength of binding to the CXCR4 receptor correlates with the anti-HIV activity. The conformation of the macrocyclic ligand and the orientations of the $\text{N}-\text{H}$ bonds are very important factors for co-receptor recognition. Therefore, a deeper knowledge of the conformation and crystal packing of metal complexes containing the cyclam ligand has become important in the development of new highly effective anti-HIV drugs that specifically target alternative events in the HIV replicative cycle (De Clercq, 2010). In addition, counter-anionic species play an important role in chemistry, pharmacy and biology



(Flores-Velez *et al.*, 1991; Fabbrizzi & Poggi, 2013). As part of a study on the structural and supramolecular features of chromium(III) complex cations with a macrocyclic ligand and with different anions, we report here the structural characterization of *trans*-[Cr(NH₃)₂(cyclam)][ZnCl₄]Cl·H₂O, (I).



2. Structural commentary

Compound (I) is another example containing a *trans*-[Cr(NH₃)₂(cyclam)]³⁺ moiety but with a different double counter-anion (Derwahl *et al.*, 1999). The asymmetric unit of (I) comprises four halves of the Cr^{III} complex cations, two tetrachloridozincate anions, two chloride anions and two water molecules. The four Cr atoms are located on crystallographic centers of symmetry. Since the complex cations have molecular *C_i* symmetry, the cyclam ligand has a *trans*-III conformation (Fig. 1). In each of the complex cations, the Cr^{III} ion is coordinated by the nitrogen atoms of the cyclam ligand occupying the equatorial sites. Two ammine ligands complete the distorted *trans*-configured octahedral coordination sphere at the axial positions. The Cr–N bond lengths including the donor atoms of the cyclam ligand range from 2.0501 (15) to 2.0615 (15) Å, comparable to those determined for *trans*-[CrCl₂(cyclam)]₂[ZnCl₄] (Flores-Velez *et al.*, 1991), *trans*-




Figure 1

The molecular structures (drawn with displacement ellipsoids at the 50% probability level) of one independent chromium(III) complex cation, one tetrachloridozincate anion, one chloride anion and one water molecule in compound (I). The primed atoms are related by symmetry code ($-x, -y + 1, -z + 2$).

Table 1

Hydrogen-bond geometry (Å, °).


<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N1A–H1NA···Cl3F ⁱ	0.90	2.80	3.3602 (16)	121
N1A–H1NA···Cl2W ⁱⁱ	0.90	2.60	3.3363 (18)	140
N1A–H3NA···Cl3E ⁱⁱⁱ	0.90	2.58	3.384 (2)	149
N2A–H1A···Cl2W ⁱⁱ	0.99	2.20	3.1754 (15)	170
N3A–H2A···Cl3F ⁱ	0.99	2.30	3.2752 (14)	170
C1A–H1A2···Cl4E ⁱⁱ	0.98	2.59	3.4739 (19)	150
N1B–H2NB···Cl1W ^{iv}	0.90	2.45	3.2683 (19)	152
N1B–H2NB···O1W	0.90	2.46	3.034 (2)	122
N1B–H3NB···Cl2E ^{iv}	0.90	2.57	3.3556 (15)	147
N2B–H1B···Cl1W	0.99	2.20	3.1704 (17)	165
N3B–H2B···O1W ^{iv}	0.99	1.98	2.968 (2)	177
N1C–H2NC···Cl3F ^v	0.90	2.68	3.4211 (19)	140
N1C–H3NC···Cl2W ^{vi}	0.90	2.38	3.2673 (17)	167
N1C–H3NC···O2W ^{vi}	0.90	2.54	2.975 (2)	111
N2C–H1C···O2W ^{vii}	0.99	1.96	2.932 (2)	167
N3C–H2C···Cl2W ^{viii}	0.99	2.23	3.2082 (16)	171
C5C–H5C2···Cl4F	0.98	2.79	3.761 (2)	174
N1D–H2ND···Cl1W ^{viii}	0.90	2.45	3.2796 (15)	154
N1D–H3ND···Cl1E ^{viii}	0.90	2.71	3.5516 (16)	156
N2D–H1D···Cl1W ^{ix}	0.99	2.19	3.1589 (16)	166
N3D–H2D···Cl2E ^{ix}	0.99	2.36	3.3276 (17)	166
C1D–H1D1···Cl1F ⁱ	0.98	2.66	3.6278 (17)	168
C1D–H1D2···Cl1E ^{viii}	0.98	2.83	3.803 (2)	172
O1W–H1O1···Cl1W ^{iv}	0.85 (1)	2.70 (2)	3.341 (2)	134 (2)
O1W–H2O1···Cl2F	0.84 (1)	2.39 (1)	3.2066 (17)	167 (2)
O2W–H1O2···Cl3E	0.83 (1)	2.37 (1)	3.1763 (19)	162 (2)
O2W–H2O2···Cl2W	0.84 (1)	2.55 (2)	3.2009 (19)	135 (2)

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $x, y, z + 1$; (iii) $x - 1, y, z + 1$; (iv) $-x + 1, -y + 1, -z + 1$; (v) $x + 1, y, z$; (vi) $-x + 1, -y + 1, -z$; (vii) $x, y + 1, z$; (viii) $-x + 1, -y, -z + 1$; (ix) $x - 1, y, z$.

[Cr(nic-O)₂(cyclam)]ClO₄ (nic-O = O-coordinating nicotinate; Choi, 2009), *trans*-[CrF₂(2,2,3-tet)]ClO₄ (2,2,3-tet = 1,4,7,11-tetrazaundecane; Choi & Moon, 2014), [Cr(ox)(cyclam)]ClO₄ (ox = oxalate; Choi *et al.*, 2004) or [Cr(acac)(cyclam)](ClO₄)₂·0.5H₂O (acac = acetylacetonate; Subhan *et al.*, 2011). However, the Cr–N bond lengths of the secondary amine group of the cyclam ligands are slightly shorter than those of the primary amine group as determined for *trans*-[CrCl₂(Me₂tn)₂]₂ZnCl₄ (Me₂tn = 2,2-dimethylpropane-1,3-diamine; Choi *et al.*, 2011), *trans*-[Cr(N₃)₂(Me₂tn)₂]ClO₄·2H₂O (Moon & Choi, 2015), or *trans*-[Cr(NCS)₂(Me₂tn)₂]SCN·0.5H₂O (Choi & Lee, 2009). The Cr–(NH₃) bond lengths range from 2.0976 (13) to 2.1062 (13) Å, similar to the average value of 2.095 (3) Å found in *trans*-[Cr(NH₃)₂(cyclam)]-(ClO₄)₂ (Derwahl *et al.*, 1999). The five-membered chelate rings of the cyclam ligands adopt *gauche* and six-membered ring chair conformations. The tetrahedral [ZnCl₄]²⁻ anion is distorted due to its involvement in hydrogen-bonding interactions. It exhibits Zn–Cl bond lengths ranging from 2.2238 (10) to 2.3232 (8) Å and Cl–Zn–Cl angles from 105.67 (3) to 115.38 (3)°.

3. Supramolecular features

In the crystal, the complex cations are stacked parallel to the *a*-axis direction. A series of N–H···Cl and C–H···Cl hydrogen bonds link the cations to neighboring anions. An extensive array of additional N–H···O and O–H···Cl


Figure 2

The crystal packing in compound (I), viewed perpendicular to the *bc* plane. Dashed lines represent hydrogen bonding interactions N—H...Cl (cyan), N—H...O (red) and O—H...Cl (purple), respectively. H atoms on C atoms have been omitted.

contacts including the lattice water molecule generates a three-dimensional network (Table 1, Fig. 2).

4. Database survey

A search in the Cambridge Structural Database (Version 5.36, last update May 2015; Groom & Allen, 2014) gave just one hit for a $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})]^{3+}$ unit, *viz.* the crystal structure of *trans*- $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})](\text{ClO}_4)\text{Cl}_2$ (Derwahl *et al.*, 1999). This dichloride perchlorate double salt and the title compound show the same *trans*-III conformation of the cyclam ligand, however with different hydrogen-bonding and crystal packing networks. The crystal structure of *cis*- $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})]\text{I}_3\cdot\text{H}_2\text{O}$ was also found (Kukina *et al.*, 1991), but no structure of any double salt of *trans*- $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})]^{3+}$ with an additional $[\text{ZnCl}_4]^{2-}$ anion.

5. Synthesis and crystallization

Cyclam and $\text{CrCl}_3(\text{THF})_3$ were purchased from Stream Chemicals and used as provided. All chemicals were reagent grade materials and used without further purification. The starting material, *trans*- $[\text{Cr}(\text{NH}_3)_2(\text{cyclam})](\text{PF}_6)(\text{NO}_3)\cdot 0.5\text{H}_2\text{O}$, was prepared according to a previously described procedure (Kane-Maguire *et al.*, 1985). The hexafluoridophosphate nitrate double salt (0.042 g) was dissolved in 5 ml of 0.01 M HCl and added to 2 ml of 1 M HCl containing 0.12 g of solid ZnCl_2 . The resulting solution was filtered and allowed to stand at room temperature for five days to give block-like yellow crystals of (I) suitable for X-ray structural analysis.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were placed in geometrically idealized positions and constrained to ride on

Table 2

Experimental details.

Crystal data	
Chemical formula	$[\text{Cr}(\text{C}_{10}\text{H}_{24}\text{N}_4)(\text{NH}_3)_2][\text{ZnCl}_4]\text{Cl}\cdot\text{H}_2\text{O}$
M_r	547.03
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	243
a, b, c (Å)	9.3980 (19), 14.876 (3), 17.981 (4)
α, β, γ (°)	66.03 (3), 76.03 (3), 78.74 (3)
V (Å ³)	2215.6 (10)
Z	4
Radiation type	Synchrotron, $\lambda = 0.620$ Å
μ (mm ⁻¹)	1.50
Crystal size (mm)	0.11 × 0.08 × 0.04
Data collection	
Diffractometer	ADSC Q210 CCD area-detector
Absorption correction	Empirical (using intensity measurements) (<i>HKL3000sm SCALEPACK</i> ; Otwinowski & Minor, 1997)
$T_{\text{min}}, T_{\text{max}}$	0.850, 0.938
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	23883, 12905, 11123
R_{int}	0.029
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.707
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.029, 0.084, 1.07
No. of reflections	12905
No. of parameters	455
No. of restraints	6
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.66, -0.76

Computer programs: *PAL BL2D-SMDC Program* (Shin *et al.*, 2016), *HKL3000sm* (Otwinowski & Minor, 1997), *SHELXT2014* (Sheldrick, 2015a), *SHELXL2014* (Sheldrick, 2015b), *DIAMOND* (Putz & Brandenburg, 2014) and *publCIF* (Westrip, 2010).

their parent atoms, with C—H = 0.98 Å and N—H = 0.90–0.99 Å and with $U_{\text{iso}}(\text{H})$ values of 1.2 or 1.5 U_{eq} of the parent atoms. The hydrogen atoms of water molecules were located in difference maps and restrained with O—H = 0.84 Å using DFIX and DANG commands (Sheldrick, 2015b).

Acknowledgements

This work was supported by a grant from 2016 Research Funds of Andong National University. The X-ray crystallography experiment at the PLS-II BL2D-SMC beamline was supported in part by MSIP and POSTECH.

References

- Choi, J.-H. (2009). *Inorg. Chim. Acta*, **362**, 4231–4236.
 Choi, J.-H., Joshi, T. & Spiccia, L. (2011). *Z. Anorg. Allg. Chem.* **637**, 1194–1198.
 Choi, J.-H. & Lee, S. H. (2009). *J. Mol. Struct.* **932**, 84–89.
 Choi, J.-H. & Moon, D. (2014). *J. Mol. Struct.* **1059**, 325–331.
 Choi, J.-H., Oh, I.-G., Suzuki, T. & Kaizaki, S. (2004). *J. Mol. Struct.* **694**, 39–44.
 De Clercq, E. (2010). *J. Med. Chem.* **53**, 1438–1450.
 Derwahl, A., Wasgestian, F., House, D. A. & Edwards, R. A. (1999). *Inorg. Chim. Acta*, **285**, 313–317.
 Fabbrizzi, L. & Poggi, A. (2013). *Chem. Soc. Rev.* **42**, 1681–1699.

- Flores-Velez, L. M., Sosa-Rivadeneira, J., Sosa-Torres, M. E., Rosales-Hoz, M. J. & Toscano, R. A. (1991). *J. Chem. Soc. Dalton Trans.* pp. 3243–3247.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Kane-Maguire, A. A. P., Wallace, K. C. & Miller, D. B. (1985). *Inorg. Chem.* **24**, 597–605.
- Kukina, G. A., Porai-Koshits, M. A., Gerda, V. I., Shevchenko, Y. N. & Shchurkina, V. N. (1991). *Sov. Phys. Crystallogr.* **36**, 739–740.
- Moon, D. & Choi, J.-H. (2015). *Spectrochim. Acta Part A*, **138**, 774–779.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Poon, C. K. & Pun, K. C. (1980). *Inorg. Chem.* **19**, 568–569.
- Putz, H. & Brandenburg, K. (2014). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Ronconi, L. & Sadler, P. J. (2007). *Coord. Chem. Rev.* **251**, 1633–1648.
- Ross, A., Choi, J.-H., Hunter, T. M., Pannecouque, C., Moggach, S. A., Parsons, S., De Clercq, E. & Sadler, P. J. (2012). *Dalton Trans.* **41**, 6408–6418.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Shin, J. W., Eom, K. & Moon, D. (2016). *J. Synchrotron Rad.* **23**, 369–373.
- Subhan, M. A., Choi, J. H. & Ng, S. W. (2011). *Z. Anorg. Allg. Chem.* **637**, 2193–2197.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2016). E72, 456–459 [https://doi.org/10.1107/S205698901600356X]

Crystal structure of *trans*-diammine(1,4,8,11-tetraazacyclotetradecane- κ^4N)chromium(III) tetrachloridozincate chloride monohydrate from synchrotron data

Dohyun Moon and Jong-Ha Choi

Computing details

Data collection: *PAL BL2D-SMDC Program* (Shin *et al.*, 2016); cell refinement: *HKL3000sm* (Otwinowski & Minor, 1997); data reduction: *HKL3000sm* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXT2014* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015b); molecular graphics: *DIAMOND* (Putz & Brandenburg, 2014); software used to prepare material for publication: *publCIF* (Westrip, 2010).

trans-Diammine(1,4,8,11-tetraazacyclotetradecane- κ^4N)chromium(III) tetrachloridozincate chloride monohydrate

Crystal data

[Cr(C₁₀H₂₄N₄)(NH₃)₂][ZnCl₄]Cl·H₂O

$M_r = 547$

Triclinic, $P\bar{1}$

$a = 9.3980$ (19) Å

$b = 14.876$ (3) Å

$c = 17.981$ (4) Å

$\alpha = 66.03$ (3)°

$\beta = 76.03$ (3)°

$\gamma = 78.74$ (3)°

$V = 2215.6$ (10) Å³

$Z = 4$

$F(000) = 1124$

$D_x = 1.640$ Mg m⁻³

Synchrotron radiation, $\lambda = 0.620$ Å

Cell parameters from 76389 reflections

$\theta = 0.4$ – 33.6 °

$\mu = 1.50$ mm⁻¹

$T = 243$ K

Block, yellow

$0.11 \times 0.08 \times 0.04$ mm

Data collection

ADSC Q210 CCD area-detector

diffractometer

Radiation source: PLSII 2D bending magnet

ω scan

Absorption correction: empirical (using intensity measurements)

(*HKL3000sm SCALEPACK*; Otwinowski & Minor, 1997)

$T_{\min} = 0.850$, $T_{\max} = 0.938$

23883 measured reflections

12905 independent reflections

11123 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 26.0$ °, $\theta_{\min} = 1.1$ °

$h = -13 \rightarrow 13$

$k = -21 \rightarrow 21$

$l = -25 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.029$

$wR(F^2) = 0.084$

$S = 1.07$

12905 reflections

455 parameters

6 restraints

Hydrogen site location: mixed
H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0513P)^2 + 0.1347P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.66 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.76 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cr1A	0.0000	0.5000	1.0000	0.00997 (6)
N1A	-0.05398 (13)	0.36162 (9)	1.01590 (7)	0.0164 (2)
H1NA	0.0293	0.3205	1.0125	0.025*
H2NA	-0.1031	0.3695	0.9761	0.025*
H3NA	-0.1111	0.3358	1.0658	0.025*
N2A	0.17264 (13)	0.42870 (9)	1.06157 (7)	0.0167 (2)
H1A	0.1790	0.3582	1.0695	0.020*
N3A	0.12143 (13)	0.51146 (9)	0.88534 (7)	0.0157 (2)
H2A	0.1219	0.4474	0.8806	0.019*
C1A	0.12606 (17)	0.43212 (12)	1.14593 (9)	0.0227 (3)
H1A1	0.1385	0.4971	1.1438	0.027*
H1A2	0.1870	0.3815	1.1838	0.027*
C2A	0.32146 (16)	0.46018 (12)	1.01888 (10)	0.0238 (3)
H2A1	0.3946	0.4181	1.0528	0.029*
H2A2	0.3230	0.5286	1.0127	0.029*
C3A	0.36314 (16)	0.45343 (13)	0.93376 (11)	0.0273 (3)
H3A1	0.4686	0.4605	0.9136	0.033*
H3A2	0.3483	0.3871	0.9400	0.033*
C4A	0.27855 (16)	0.52922 (12)	0.86811 (9)	0.0238 (3)
H4A1	0.2832	0.5956	0.8656	0.029*
H4A2	0.3257	0.5267	0.8140	0.029*
C5A	0.03428 (17)	0.58654 (12)	0.82368 (8)	0.0218 (3)
H5A1	0.0693	0.5810	0.7696	0.026*
H5A2	0.0457	0.6533	0.8177	0.026*
Cr2B	0.5000	0.5000	0.5000	0.01185 (6)
N1B	0.30918 (13)	0.57334 (9)	0.54783 (7)	0.0188 (2)
H1NB	0.2410	0.5307	0.5761	0.028*
H2NB	0.2731	0.6239	0.5060	0.028*
H3NB	0.3315	0.5969	0.5820	0.028*
N2B	0.44033 (14)	0.36582 (9)	0.58615 (8)	0.0204 (2)
H1B	0.5253	0.3162	0.5812	0.024*
N3B	0.62099 (14)	0.51975 (10)	0.57270 (8)	0.0218 (2)
H2B	0.7178	0.4803	0.5666	0.026*
C1B	0.3178 (2)	0.34242 (13)	0.55956 (12)	0.0317 (4)

H1B1	0.2242	0.3773	0.5768	0.038*
H1B2	0.3095	0.2712	0.5856	0.038*
C2B	0.40555 (19)	0.35503 (13)	0.67422 (10)	0.0299 (4)
H2B1	0.3851	0.2869	0.7092	0.036*
H2B2	0.3163	0.3990	0.6828	0.036*
C3B	0.5303 (2)	0.37936 (14)	0.70072 (10)	0.0318 (4)
H3B1	0.6206	0.3389	0.6873	0.038*
H3B2	0.5079	0.3591	0.7610	0.038*
C4B	0.5622 (2)	0.48701 (14)	0.66301 (10)	0.0300 (4)
H4B1	0.4712	0.5291	0.6725	0.036*
H4B2	0.6341	0.4950	0.6907	0.036*
C5B	0.6501 (2)	0.62542 (13)	0.53436 (12)	0.0315 (4)
H5B1	0.7347	0.6335	0.5531	0.038*
H5B2	0.5640	0.6667	0.5511	0.038*
Cr3C	0.5000	1.0000	0.0000	0.01576 (6)
N1C	0.54753 (15)	0.84509 (9)	0.04233 (8)	0.0240 (3)
H1NC	0.4628	0.8170	0.0595	0.036*
H2NC	0.5966	0.8240	0.0847	0.036*
H3NC	0.6036	0.8280	0.0009	0.036*
N2C	0.63796 (15)	1.01044 (11)	0.06889 (8)	0.0256 (3)
H1C	0.6218	1.0801	0.0645	0.031*
N3C	0.31688 (15)	0.98526 (10)	0.09225 (8)	0.0243 (3)
H2C	0.2794	1.0527	0.0913	0.029*
C1C	0.79153 (19)	0.99551 (16)	0.02512 (12)	0.0362 (4)
H1C1	0.8224	0.9246	0.0388	0.043*
H1C2	0.8590	1.0222	0.0429	0.043*
C2C	0.6153 (2)	0.94799 (15)	0.15942 (11)	0.0342 (4)
H2C1	0.6799	0.9657	0.1856	0.041*
H2C2	0.6432	0.8783	0.1669	0.041*
C3C	0.4561 (2)	0.96092 (15)	0.20207 (10)	0.0352 (4)
H3C1	0.4520	0.9271	0.2619	0.042*
H3C2	0.4278	1.0316	0.1908	0.042*
C4C	0.3413 (2)	0.92345 (13)	0.17816 (10)	0.0323 (4)
H4C1	0.3738	0.8550	0.1827	0.039*
H4C2	0.2479	0.9238	0.2168	0.039*
C5C	0.20231 (19)	0.95212 (15)	0.06752 (12)	0.0338 (4)
H5C1	0.1045	0.9676	0.0970	0.041*
H5C2	0.2198	0.8803	0.0822	0.041*
Cr4D	0.0000	0.0000	0.5000	0.00919 (6)
N1D	0.22787 (12)	-0.04570 (9)	0.48598 (7)	0.0160 (2)
H1ND	0.2753	-0.0002	0.4410	0.024*
H2ND	0.2450	-0.1042	0.4798	0.024*
H3ND	0.2609	-0.0523	0.5311	0.024*
N2D	0.02941 (13)	0.09509 (9)	0.55052 (8)	0.0177 (2)
H1D	-0.0667	0.1356	0.5558	0.021*
N3D	0.02438 (12)	0.10266 (9)	0.38037 (7)	0.0155 (2)
H2D	-0.0727	0.1423	0.3738	0.019*
C1D	0.05317 (18)	0.03371 (13)	0.63680 (10)	0.0255 (3)

H1D1	0.0337	0.0752	0.6695	0.031*
H1D2	0.1555	0.0031	0.6363	0.031*
C2D	0.13974 (19)	0.16666 (12)	0.50160 (11)	0.0284 (3)
H2D1	0.2385	0.1305	0.4978	0.034*
H2D2	0.1382	0.2108	0.5301	0.034*
C3D	0.1080 (2)	0.22836 (12)	0.41438 (12)	0.0320 (4)
H3D1	0.0048	0.2573	0.4193	0.038*
H3D2	0.1694	0.2832	0.3893	0.038*
C4D	0.13418 (17)	0.17441 (12)	0.35548 (10)	0.0264 (3)
H4D1	0.1281	0.2229	0.2994	0.032*
H4D2	0.2337	0.1390	0.3545	0.032*
C5D	0.05122 (17)	0.04541 (12)	0.32576 (9)	0.0238 (3)
H5D1	0.1536	0.0149	0.3213	0.029*
H5D2	0.0340	0.0896	0.2702	0.029*
Zn1E	0.61151 (2)	0.20650 (2)	0.29082 (2)	0.02403 (5)
Cl1E	0.53804 (5)	0.05903 (4)	0.38072 (3)	0.04285 (12)
Cl2E	0.72540 (4)	0.26657 (3)	0.35901 (2)	0.02630 (8)
Cl3E	0.79024 (5)	0.18205 (4)	0.18756 (3)	0.03439 (9)
Cl4E	0.43263 (5)	0.32125 (4)	0.23927 (3)	0.04760 (14)
Zn2F	0.01806 (2)	0.67654 (2)	0.21094 (2)	0.02221 (5)
Cl1F	-0.02844 (7)	0.79302 (4)	0.26421 (3)	0.04689 (13)
Cl2F	0.00187 (5)	0.52354 (3)	0.31592 (2)	0.02954 (8)
Cl3F	-0.16681 (4)	0.70070 (3)	0.13702 (2)	0.02651 (8)
Cl4F	0.23650 (5)	0.67718 (4)	0.12345 (3)	0.03785 (10)
Cl1W	0.72406 (5)	0.20867 (3)	0.60004 (2)	0.02949 (9)
Cl2W	0.20883 (5)	0.19663 (3)	0.10707 (3)	0.03019 (9)
O1W	0.09042 (18)	0.60251 (13)	0.43830 (10)	0.0463 (4)
H1O1	0.103 (3)	0.6621 (9)	0.4071 (13)	0.056*
H2O1	0.055 (3)	0.5784 (17)	0.4128 (14)	0.056*
O2W	0.55603 (19)	0.20704 (13)	0.07745 (10)	0.0473 (4)
H1O2	0.6248 (18)	0.211 (2)	0.0971 (15)	0.057*
H2O2	0.4729 (15)	0.215 (2)	0.1061 (14)	0.057*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cr1A	0.00887 (12)	0.00962 (12)	0.00992 (12)	-0.00051 (9)	-0.00151 (9)	-0.00252 (10)
N1A	0.0176 (5)	0.0146 (5)	0.0162 (5)	-0.0023 (4)	-0.0015 (4)	-0.0056 (4)
N2A	0.0132 (5)	0.0156 (5)	0.0202 (5)	0.0000 (4)	-0.0071 (4)	-0.0043 (5)
N3A	0.0147 (5)	0.0159 (5)	0.0150 (5)	-0.0033 (4)	0.0015 (4)	-0.0061 (4)
C1A	0.0261 (7)	0.0253 (7)	0.0168 (6)	-0.0044 (6)	-0.0111 (5)	-0.0031 (6)
C2A	0.0131 (6)	0.0249 (7)	0.0320 (8)	-0.0014 (5)	-0.0076 (5)	-0.0078 (6)
C3A	0.0127 (6)	0.0316 (8)	0.0361 (8)	0.0008 (6)	0.0006 (6)	-0.0155 (7)
C4A	0.0163 (6)	0.0289 (8)	0.0241 (7)	-0.0065 (5)	0.0062 (5)	-0.0117 (6)
C5A	0.0280 (7)	0.0240 (7)	0.0113 (6)	-0.0066 (6)	-0.0044 (5)	-0.0021 (5)
Cr2B	0.01081 (12)	0.01190 (13)	0.01273 (13)	-0.00091 (10)	0.00002 (10)	-0.00589 (11)
N1B	0.0173 (5)	0.0188 (6)	0.0183 (5)	0.0006 (4)	0.0005 (4)	-0.0081 (5)
N2B	0.0189 (6)	0.0160 (6)	0.0209 (6)	-0.0024 (4)	0.0027 (4)	-0.0048 (5)

N3B	0.0206 (6)	0.0259 (6)	0.0243 (6)	-0.0011 (5)	-0.0058 (5)	-0.0147 (5)
C1B	0.0276 (8)	0.0266 (8)	0.0400 (9)	-0.0135 (6)	0.0015 (7)	-0.0115 (7)
C2B	0.0306 (8)	0.0269 (8)	0.0184 (7)	0.0007 (6)	0.0054 (6)	-0.0018 (6)
C3B	0.0370 (9)	0.0351 (9)	0.0174 (7)	0.0065 (7)	-0.0056 (6)	-0.0081 (7)
C4B	0.0353 (9)	0.0363 (9)	0.0230 (7)	0.0062 (7)	-0.0104 (6)	-0.0175 (7)
C5B	0.0338 (9)	0.0286 (8)	0.0420 (10)	-0.0100 (7)	-0.0075 (7)	-0.0199 (8)
Cr3C	0.01463 (14)	0.01256 (14)	0.01522 (14)	0.00175 (11)	-0.00218 (11)	-0.00211 (11)
N1C	0.0272 (6)	0.0172 (6)	0.0221 (6)	0.0031 (5)	-0.0050 (5)	-0.0042 (5)
N2C	0.0237 (6)	0.0261 (7)	0.0250 (6)	-0.0005 (5)	-0.0076 (5)	-0.0069 (6)
N3C	0.0221 (6)	0.0203 (6)	0.0228 (6)	0.0000 (5)	0.0018 (5)	-0.0048 (5)
C1C	0.0201 (7)	0.0428 (11)	0.0411 (10)	0.0004 (7)	-0.0091 (7)	-0.0111 (9)
C2C	0.0407 (10)	0.0337 (9)	0.0252 (8)	0.0004 (8)	-0.0146 (7)	-0.0051 (7)
C3C	0.0482 (11)	0.0321 (9)	0.0191 (7)	-0.0011 (8)	-0.0037 (7)	-0.0061 (7)
C4C	0.0388 (9)	0.0270 (8)	0.0192 (7)	-0.0024 (7)	0.0040 (6)	-0.0024 (6)
C5C	0.0198 (7)	0.0373 (10)	0.0372 (9)	-0.0077 (7)	0.0014 (6)	-0.0087 (8)
Cr4D	0.00721 (11)	0.00831 (12)	0.01150 (12)	-0.00106 (9)	-0.00070 (9)	-0.00365 (10)
N1D	0.0103 (5)	0.0159 (5)	0.0192 (5)	-0.0007 (4)	-0.0014 (4)	-0.0050 (4)
N2D	0.0163 (5)	0.0159 (5)	0.0249 (6)	-0.0011 (4)	-0.0047 (4)	-0.0116 (5)
N3D	0.0123 (5)	0.0140 (5)	0.0156 (5)	-0.0012 (4)	-0.0018 (4)	-0.0015 (4)
C1D	0.0275 (7)	0.0327 (8)	0.0244 (7)	0.0014 (6)	-0.0093 (6)	-0.0183 (7)
C2D	0.0268 (8)	0.0224 (7)	0.0427 (9)	-0.0117 (6)	-0.0071 (7)	-0.0145 (7)
C3D	0.0342 (9)	0.0152 (7)	0.0440 (10)	-0.0122 (6)	-0.0086 (7)	-0.0037 (7)
C4D	0.0209 (7)	0.0220 (7)	0.0255 (7)	-0.0094 (6)	-0.0011 (6)	0.0032 (6)
C5D	0.0231 (7)	0.0316 (8)	0.0140 (6)	0.0018 (6)	-0.0018 (5)	-0.0090 (6)
Zn1E	0.01617 (8)	0.02918 (10)	0.01970 (9)	-0.00577 (7)	-0.00537 (6)	0.00052 (7)
Cl1E	0.02366 (19)	0.0392 (2)	0.0439 (3)	-0.01488 (18)	-0.00507 (17)	0.0108 (2)
Cl2E	0.02669 (18)	0.02502 (18)	0.02753 (18)	0.00389 (14)	-0.01067 (14)	-0.01004 (15)
Cl3E	0.0306 (2)	0.0393 (2)	0.02640 (19)	-0.00741 (17)	0.00146 (15)	-0.00763 (18)
Cl4E	0.02096 (19)	0.0463 (3)	0.0538 (3)	-0.00297 (18)	-0.01940 (19)	0.0098 (2)
Zn2F	0.02487 (9)	0.02355 (9)	0.02018 (9)	-0.00586 (7)	0.00014 (6)	-0.01115 (7)
Cl1F	0.0642 (3)	0.0419 (3)	0.0483 (3)	-0.0180 (2)	0.0050 (2)	-0.0338 (2)
Cl2F	0.0346 (2)	0.02632 (19)	0.02337 (17)	-0.00454 (15)	-0.00178 (15)	-0.00646 (15)
Cl3F	0.02952 (18)	0.02757 (18)	0.02467 (17)	0.00596 (14)	-0.00829 (14)	-0.01429 (15)
Cl4F	0.0297 (2)	0.0382 (2)	0.0430 (2)	-0.01085 (17)	0.01168 (17)	-0.0196 (2)
Cl1W	0.0332 (2)	0.02203 (17)	0.02665 (18)	0.01129 (15)	-0.00334 (15)	-0.01016 (15)
Cl2W	0.0311 (2)	0.01957 (17)	0.0317 (2)	0.00488 (14)	-0.00070 (15)	-0.00768 (15)
O1W	0.0420 (8)	0.0565 (10)	0.0491 (9)	0.0026 (7)	-0.0231 (7)	-0.0239 (8)
O2W	0.0468 (9)	0.0571 (10)	0.0496 (9)	-0.0097 (8)	-0.0099 (7)	-0.0294 (8)

Geometric parameters (Å, °)

Cr1A—N2A ⁱ	2.0553 (13)	N1C—H1NC	0.9000
Cr1A—N2A	2.0553 (13)	N1C—H2NC	0.9000
Cr1A—N3A ⁱ	2.0582 (13)	N1C—H3NC	0.9000
Cr1A—N3A	2.0582 (13)	N2C—C1C	1.490 (2)
Cr1A—N1A	2.1062 (13)	N2C—C2C	1.496 (2)
Cr1A—N1A ⁱ	2.1062 (13)	N2C—H1C	0.9900
N1A—H1NA	0.9000	N3C—C5C	1.489 (2)

N1A—H2NA	0.9000	N3C—C4C	1.490 (2)
N1A—H3NA	0.9000	N3C—H2C	0.9900
N2A—C2A	1.4866 (19)	C1C—C5C ⁱⁱⁱ	1.517 (3)
N2A—C1A	1.4928 (19)	C1C—H1C1	0.9800
N2A—H1A	0.9900	C1C—H1C2	0.9800
N3A—C4A	1.4869 (19)	C2C—C3C	1.522 (3)
N3A—C5A	1.491 (2)	C2C—H2C1	0.9800
N3A—H2A	0.9900	C2C—H2C2	0.9800
C1A—C5A ⁱ	1.513 (2)	C3C—C4C	1.521 (3)
C1A—H1A1	0.9800	C3C—H3C1	0.9800
C1A—H1A2	0.9800	C3C—H3C2	0.9800
C2A—C3A	1.525 (2)	C4C—H4C1	0.9800
C2A—H2A1	0.9800	C4C—H4C2	0.9800
C2A—H2A2	0.9800	C5C—C1C ⁱⁱⁱ	1.517 (3)
C3A—C4A	1.523 (2)	C5C—H5C1	0.9800
C3A—H3A1	0.9800	C5C—H5C2	0.9800
C3A—H3A2	0.9800	Cr4D—N2D ^{iv}	2.0544 (12)
C4A—H4A1	0.9800	Cr4D—N2D	2.0544 (12)
C4A—H4A2	0.9800	Cr4D—N3D	2.0593 (14)
C5A—C1A ⁱ	1.513 (2)	Cr4D—N3D ^{iv}	2.0593 (14)
C5A—H5A1	0.9800	Cr4D—N1D ^{iv}	2.1029 (12)
C5A—H5A2	0.9800	Cr4D—N1D	2.1029 (12)
Cr2B—N2B ⁱⁱ	2.0501 (15)	N1D—H1ND	0.9000
Cr2B—N2B	2.0502 (15)	N1D—H2ND	0.9000
Cr2B—N3B	2.0611 (13)	N1D—H3ND	0.9000
Cr2B—N3B ⁱⁱ	2.0611 (13)	N2D—C2D	1.487 (2)
Cr2B—N1B	2.0976 (13)	N2D—C1D	1.492 (2)
Cr2B—N1B ⁱⁱ	2.0977 (13)	N2D—H1D	0.9900
N1B—H1NB	0.9000	N3D—C4D	1.491 (2)
N1B—H2NB	0.9000	N3D—C5D	1.4932 (19)
N1B—H3NB	0.9000	N3D—H2D	0.9900
N2B—C2B	1.485 (2)	C1D—C5D ^{iv}	1.513 (2)
N2B—C1B	1.494 (2)	C1D—H1D1	0.9800
N2B—H1B	0.9900	C1D—H1D2	0.9800
N3B—C4B	1.486 (2)	C2D—C3D	1.530 (3)
N3B—C5B	1.489 (2)	C2D—H2D1	0.9800
N3B—H2B	0.9900	C2D—H2D2	0.9800
C1B—C5B ⁱⁱ	1.525 (3)	C3D—C4D	1.521 (3)
C1B—H1B1	0.9800	C3D—H3D1	0.9800
C1B—H1B2	0.9800	C3D—H3D2	0.9800
C2B—C3B	1.519 (3)	C4D—H4D1	0.9800
C2B—H2B1	0.9800	C4D—H4D2	0.9800
C2B—H2B2	0.9800	C5D—C1D ^{iv}	1.513 (2)
C3B—C4B	1.524 (3)	C5D—H5D1	0.9800
C3B—H3B1	0.9800	C5D—H5D2	0.9800
C3B—H3B2	0.9800	Zn1E—C14E	2.2238 (10)
C4B—H4B1	0.9800	Zn1E—C11E	2.2523 (11)
C4B—H4B2	0.9800	Zn1E—C13E	2.2817 (9)

C5B—C1B ⁱⁱ	1.525 (3)	Zn1E—C12E	2.3118 (7)
C5B—H5B1	0.9800	Zn2F—C11F	2.2275 (7)
C5B—H5B2	0.9800	Zn2F—C14F	2.2640 (9)
Cr3C—N3C ⁱⁱⁱ	2.0579 (15)	Zn2F—C12F	2.2960 (11)
Cr3C—N3C	2.0579 (15)	Zn2F—C13F	2.3232 (8)
Cr3C—N2C ⁱⁱⁱ	2.0615 (15)	O1W—H1O1	0.848 (9)
Cr3C—N2C	2.0615 (15)	O1W—H2O1	0.836 (9)
Cr3C—N1C	2.1039 (14)	O2W—H1O2	0.832 (9)
Cr3C—N1C ⁱⁱⁱ	2.1039 (14)	O2W—H2O2	0.843 (9)
N2A ⁱ —Cr1A—N2A	180.0	N3C—Cr3C—N1C	90.06 (6)
N2A ⁱ —Cr1A—N3A ⁱ	94.49 (5)	N2C ⁱⁱⁱ —Cr3C—N1C	88.08 (6)
N2A—Cr1A—N3A ⁱ	85.51 (5)	N2C—Cr3C—N1C	91.92 (6)
N2A ⁱ —Cr1A—N3A	85.51 (5)	N3C ⁱⁱⁱ —Cr3C—N1C ⁱⁱⁱ	90.05 (6)
N2A—Cr1A—N3A	94.49 (5)	N3C—Cr3C—N1C ⁱⁱⁱ	89.94 (6)
N3A ⁱ —Cr1A—N3A	180.00 (4)	N2C ⁱⁱⁱ —Cr3C—N1C ⁱⁱⁱ	91.92 (6)
N2A ⁱ —Cr1A—N1A	90.76 (5)	N2C—Cr3C—N1C ⁱⁱⁱ	88.08 (6)
N2A—Cr1A—N1A	89.24 (5)	N1C—Cr3C—N1C ⁱⁱⁱ	180.00 (8)
N3A ⁱ —Cr1A—N1A	91.07 (6)	Cr3C—N1C—H1NC	109.5
N3A—Cr1A—N1A	88.93 (6)	Cr3C—N1C—H2NC	109.5
N2A ⁱ —Cr1A—N1A ⁱ	89.24 (5)	H1NC—N1C—H2NC	109.5
N2A—Cr1A—N1A ⁱ	90.76 (5)	Cr3C—N1C—H3NC	109.5
N3A ⁱ —Cr1A—N1A ⁱ	88.93 (6)	H1NC—N1C—H3NC	109.5
N3A—Cr1A—N1A ⁱ	91.07 (6)	H2NC—N1C—H3NC	109.5
N1A—Cr1A—N1A ⁱ	180.00 (6)	C1C—N2C—C2C	113.16 (14)
Cr1A—N1A—H1NA	109.5	C1C—N2C—Cr3C	106.52 (11)
Cr1A—N1A—H2NA	109.5	C2C—N2C—Cr3C	117.72 (12)
H1NA—N1A—H2NA	109.5	C1C—N2C—H1C	106.2
Cr1A—N1A—H3NA	109.5	C2C—N2C—H1C	106.2
H1NA—N1A—H3NA	109.5	Cr3C—N2C—H1C	106.2
H2NA—N1A—H3NA	109.5	C5C—N3C—C4C	113.35 (14)
C2A—N2A—C1A	114.29 (12)	C5C—N3C—Cr3C	107.00 (10)
C2A—N2A—Cr1A	117.18 (9)	C4C—N3C—Cr3C	116.36 (11)
C1A—N2A—Cr1A	106.44 (9)	C5C—N3C—H2C	106.5
C2A—N2A—H1A	106.0	C4C—N3C—H2C	106.5
C1A—N2A—H1A	106.0	Cr3C—N3C—H2C	106.5
Cr1A—N2A—H1A	106.0	N2C—C1C—C5C ⁱⁱⁱ	109.33 (14)
C4A—N3A—C5A	113.52 (12)	N2C—C1C—H1C1	109.8
C4A—N3A—Cr1A	117.51 (9)	C5C ⁱⁱⁱ —C1C—H1C1	109.8
C5A—N3A—Cr1A	106.14 (9)	N2C—C1C—H1C2	109.8
C4A—N3A—H2A	106.3	C5C ⁱⁱⁱ —C1C—H1C2	109.8
C5A—N3A—H2A	106.3	H1C1—C1C—H1C2	108.3
Cr1A—N3A—H2A	106.3	N2C—C2C—C3C	112.24 (15)
N2A—C1A—C5A ⁱ	108.42 (12)	N2C—C2C—H2C1	109.2
N2A—C1A—H1A1	110.0	C3C—C2C—H2C1	109.2
C5A ⁱ —C1A—H1A1	110.0	N2C—C2C—H2C2	109.2
N2A—C1A—H1A2	110.0	C3C—C2C—H2C2	109.2
C5A ⁱ —C1A—H1A2	110.0	H2C1—C2C—H2C2	107.9

H1A1—C1A—H1A2	108.4	C4C—C3C—C2C	116.90 (15)
N2A—C2A—C3A	111.60 (13)	C4C—C3C—H3C1	108.1
N2A—C2A—H2A1	109.3	C2C—C3C—H3C1	108.1
C3A—C2A—H2A1	109.3	C4C—C3C—H3C2	108.1
N2A—C2A—H2A2	109.3	C2C—C3C—H3C2	108.1
C3A—C2A—H2A2	109.3	H3C1—C3C—H3C2	107.3
H2A1—C2A—H2A2	108.0	N3C—C4C—C3C	111.92 (15)
C4A—C3A—C2A	115.86 (13)	N3C—C4C—H4C1	109.2
C4A—C3A—H3A1	108.3	C3C—C4C—H4C1	109.2
C2A—C3A—H3A1	108.3	N3C—C4C—H4C2	109.2
C4A—C3A—H3A2	108.3	C3C—C4C—H4C2	109.2
C2A—C3A—H3A2	108.3	H4C1—C4C—H4C2	107.9
H3A1—C3A—H3A2	107.4	N3C—C5C—C1C ⁱⁱⁱ	109.26 (15)
N3A—C4A—C3A	112.23 (13)	N3C—C5C—H5C1	109.8
N3A—C4A—H4A1	109.2	C1C ⁱⁱⁱ —C5C—H5C1	109.8
C3A—C4A—H4A1	109.2	N3C—C5C—H5C2	109.8
N3A—C4A—H4A2	109.2	C1C ⁱⁱⁱ —C5C—H5C2	109.8
C3A—C4A—H4A2	109.2	H5C1—C5C—H5C2	108.3
H4A1—C4A—H4A2	107.9	N2D ^{iv} —Cr4D—N2D	180.0
N3A—C5A—C1A ⁱ	108.15 (12)	N2D ^{iv} —Cr4D—N3D	85.25 (5)
N3A—C5A—H5A1	110.1	N2D—Cr4D—N3D	94.75 (5)
C1A ⁱ —C5A—H5A1	110.1	N2D ^{iv} —Cr4D—N3D ^{iv}	94.76 (5)
N3A—C5A—H5A2	110.1	N2D—Cr4D—N3D ^{iv}	85.24 (5)
C1A ⁱ —C5A—H5A2	110.1	N3D—Cr4D—N3D ^{iv}	180.0
H5A1—C5A—H5A2	108.4	N2D ^{iv} —Cr4D—N1D ^{iv}	89.83 (5)
N2B ⁱⁱ —Cr2B—N2B	180.0	N2D—Cr4D—N1D ^{iv}	90.17 (5)
N2B ⁱⁱ —Cr2B—N3B	85.91 (6)	N3D—Cr4D—N1D ^{iv}	88.87 (6)
N2B—Cr2B—N3B	94.09 (6)	N3D ^{iv} —Cr4D—N1D ^{iv}	91.13 (6)
N2B ⁱⁱ —Cr2B—N3B ⁱⁱ	94.09 (6)	N2D ^{iv} —Cr4D—N1D	90.17 (5)
N2B—Cr2B—N3B ⁱⁱ	85.91 (6)	N2D—Cr4D—N1D	89.83 (5)
N3B—Cr2B—N3B ⁱⁱ	180.00 (7)	N3D—Cr4D—N1D	91.13 (6)
N2B ⁱⁱ —Cr2B—N1B	89.22 (6)	N3D ^{iv} —Cr4D—N1D	88.87 (6)
N2B—Cr2B—N1B	90.78 (6)	N1D ^{iv} —Cr4D—N1D	180.0
N3B—Cr2B—N1B	91.36 (5)	Cr4D—N1D—H1ND	109.5
N3B ⁱⁱ —Cr2B—N1B	88.64 (5)	Cr4D—N1D—H2ND	109.5
N2B ⁱⁱ —Cr2B—N1B ⁱⁱ	90.78 (6)	H1ND—N1D—H2ND	109.5
N2B—Cr2B—N1B ⁱⁱ	89.22 (6)	Cr4D—N1D—H3ND	109.5
N3B—Cr2B—N1B ⁱⁱ	88.64 (5)	H1ND—N1D—H3ND	109.5
N3B ⁱⁱ —Cr2B—N1B ⁱⁱ	91.36 (5)	H2ND—N1D—H3ND	109.5
N1B—Cr2B—N1B ⁱⁱ	180.00 (6)	C2D—N2D—C1D	114.07 (12)
Cr2B—N1B—H1NB	109.5	C2D—N2D—Cr4D	117.13 (10)
Cr2B—N1B—H2NB	109.5	C1D—N2D—Cr4D	107.05 (9)
H1NB—N1B—H2NB	109.5	C2D—N2D—H1D	105.9
Cr2B—N1B—H3NB	109.5	C1D—N2D—H1D	105.9
H1NB—N1B—H3NB	109.5	Cr4D—N2D—H1D	105.9
H2NB—N1B—H3NB	109.5	C4D—N3D—C5D	113.03 (12)
C2B—N2B—C1B	112.86 (13)	C4D—N3D—Cr4D	117.41 (10)
C2B—N2B—Cr2B	116.96 (11)	C5D—N3D—Cr4D	106.16 (9)

C1B—N2B—Cr2B	106.88 (10)	C4D—N3D—H2D	106.5
C2B—N2B—H1B	106.5	C5D—N3D—H2D	106.5
C1B—N2B—H1B	106.5	Cr4D—N3D—H2D	106.5
Cr2B—N2B—H1B	106.5	N2D—C1D—C5D ^{iv}	108.07 (12)
C4B—N3B—C5B	112.83 (13)	N2D—C1D—H1D1	110.1
C4B—N3B—Cr2B	117.59 (11)	C5D ^{iv} —C1D—H1D1	110.1
C5B—N3B—Cr2B	106.77 (10)	N2D—C1D—H1D2	110.1
C4B—N3B—H2B	106.3	C5D ^{iv} —C1D—H1D2	110.1
C5B—N3B—H2B	106.3	H1D1—C1D—H1D2	108.4
Cr2B—N3B—H2B	106.3	N2D—C2D—C3D	111.55 (13)
N2B—C1B—C5B ⁱⁱ	109.12 (14)	N2D—C2D—H2D1	109.3
N2B—C1B—H1B1	109.9	C3D—C2D—H2D1	109.3
C5B ⁱⁱ —C1B—H1B1	109.9	N2D—C2D—H2D2	109.3
N2B—C1B—H1B2	109.9	C3D—C2D—H2D2	109.3
C5B ⁱⁱ —C1B—H1B2	109.9	H2D1—C2D—H2D2	108.0
H1B1—C1B—H1B2	108.3	C4D—C3D—C2D	116.59 (14)
N2B—C2B—C3B	112.58 (14)	C4D—C3D—H3D1	108.1
N2B—C2B—H2B1	109.1	C2D—C3D—H3D1	108.1
C3B—C2B—H2B1	109.1	C4D—C3D—H3D2	108.1
N2B—C2B—H2B2	109.1	C2D—C3D—H3D2	108.1
C3B—C2B—H2B2	109.1	H3D1—C3D—H3D2	107.3
H2B1—C2B—H2B2	107.8	N3D—C4D—C3D	111.82 (13)
C2B—C3B—C4B	116.95 (15)	N3D—C4D—H4D1	109.3
C2B—C3B—H3B1	108.1	C3D—C4D—H4D1	109.3
C4B—C3B—H3B1	108.1	N3D—C4D—H4D2	109.3
C2B—C3B—H3B2	108.1	C3D—C4D—H4D2	109.3
C4B—C3B—H3B2	108.1	H4D1—C4D—H4D2	107.9
H3B1—C3B—H3B2	107.3	N3D—C5D—C1D ^{iv}	108.31 (12)
N3B—C4B—C3B	112.03 (13)	N3D—C5D—H5D1	110.0
N3B—C4B—H4B1	109.2	C1D ^{iv} —C5D—H5D1	110.0
C3B—C4B—H4B1	109.2	N3D—C5D—H5D2	110.0
N3B—C4B—H4B2	109.2	C1D ^{iv} —C5D—H5D2	110.0
C3B—C4B—H4B2	109.2	H5D1—C5D—H5D2	108.4
H4B1—C4B—H4B2	107.9	Cl4E—Zn1E—Cl1E	115.38 (3)
N3B—C5B—C1B ⁱⁱ	109.08 (13)	Cl4E—Zn1E—Cl3E	110.99 (3)
N3B—C5B—H5B1	109.9	Cl1E—Zn1E—Cl3E	108.75 (4)
C1B ⁱⁱ —C5B—H5B1	109.9	Cl4E—Zn1E—Cl2E	107.79 (3)
N3B—C5B—H5B2	109.9	Cl1E—Zn1E—Cl2E	107.77 (3)
C1B ⁱⁱ —C5B—H5B2	109.9	Cl3E—Zn1E—Cl2E	105.67 (3)
H5B1—C5B—H5B2	108.3	Cl1F—Zn2F—Cl4F	115.32 (3)
N3C ⁱⁱⁱ —Cr3C—N3C	180.00 (6)	Cl1F—Zn2F—Cl2F	109.63 (3)
N3C ⁱⁱⁱ —Cr3C—N2C ⁱⁱⁱ	94.32 (6)	Cl4F—Zn2F—Cl2F	109.79 (4)
N3C—Cr3C—N2C ⁱⁱⁱ	85.68 (6)	Cl1F—Zn2F—Cl3F	107.62 (3)
N3C ⁱⁱⁱ —Cr3C—N2C	85.68 (6)	Cl4F—Zn2F—Cl3F	107.31 (2)
N3C—Cr3C—N2C	94.32 (6)	Cl2F—Zn2F—Cl3F	106.77 (4)
N2C ⁱⁱⁱ —Cr3C—N2C	180.0	H1O1—O1W—H2O1	107.9 (19)
N3C ⁱⁱⁱ —Cr3C—N1C	89.94 (6)	H1O2—O2W—H2O2	112.1 (19)

C2A—N2A—C1A—C5A ⁱ	171.57 (12)	C2C—N2C—C1C—C5C ⁱⁱⁱ	170.49 (16)
Cr1A—N2A—C1A—C5A ⁱ	40.59 (14)	Cr3C—N2C—C1C—C5C ⁱⁱⁱ	39.61 (18)
C1A—N2A—C2A—C3A	179.11 (12)	C1C—N2C—C2C—C3C	-177.36 (16)
Cr1A—N2A—C2A—C3A	-55.38 (15)	Cr3C—N2C—C2C—C3C	-52.33 (19)
N2A—C2A—C3A—C4A	70.45 (18)	N2C—C2C—C3C—C4C	67.1 (2)
C5A—N3A—C4A—C3A	178.01 (12)	C5C—N3C—C4C—C3C	-178.55 (14)
Cr1A—N3A—C4A—C3A	53.37 (15)	Cr3C—N3C—C4C—C3C	56.74 (17)
C2A—C3A—C4A—N3A	-69.39 (18)	C2C—C3C—C4C—N3C	-69.9 (2)
C4A—N3A—C5A—C1A ⁱ	-172.82 (12)	C4C—N3C—C5C—C1C ⁱⁱⁱ	-167.98 (14)
Cr1A—N3A—C5A—C1A ⁱ	-42.24 (13)	Cr3C—N3C—C5C—C1C ⁱⁱⁱ	-38.36 (16)
C2B—N2B—C1B—C5B ⁱⁱ	169.01 (14)	C2D—N2D—C1D—C5D ^{iv}	-171.39 (12)
Cr2B—N2B—C1B—C5B ⁱⁱ	39.06 (15)	Cr4D—N2D—C1D—C5D ^{iv}	-40.14 (13)
C1B—N2B—C2B—C3B	-179.75 (14)	C1D—N2D—C2D—C3D	-179.09 (13)
Cr2B—N2B—C2B—C3B	-55.15 (16)	Cr4D—N2D—C2D—C3D	54.77 (16)
N2B—C2B—C3B—C4B	68.20 (19)	N2D—C2D—C3D—C4D	-70.12 (19)
C5B—N3B—C4B—C3B	178.83 (14)	C5D—N3D—C4D—C3D	-177.52 (13)
Cr2B—N3B—C4B—C3B	53.86 (17)	Cr4D—N3D—C4D—C3D	-53.38 (16)
C2B—C3B—C4B—N3B	-67.16 (19)	C2D—C3D—C4D—N3D	69.25 (19)
C4B—N3B—C5B—C1B ⁱⁱ	-169.58 (14)	C4D—N3D—C5D—C1D ^{iv}	172.59 (12)
Cr2B—N3B—C5B—C1B ⁱⁱ	-38.92 (16)	Cr4D—N3D—C5D—C1D ^{iv}	42.50 (13)

Symmetry codes: (i) $-x, -y+1, -z+2$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x+1, -y+2, -z$; (iv) $-x, -y, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1A—H1NA \cdots Cl3F ^v	0.90	2.80	3.3602 (16)	121
N1A—H1NA \cdots Cl2W ^{vi}	0.90	2.60	3.3363 (18)	140
N1A—H3NA \cdots Cl3E ^{vii}	0.90	2.58	3.384 (2)	149
N2A—H1A \cdots Cl2W ^{vi}	0.99	2.20	3.1754 (15)	170
N3A—H2A \cdots Cl3F ^v	0.99	2.30	3.2752 (14)	170
C1A—H1A2 \cdots Cl4E ^{vi}	0.98	2.59	3.4739 (19)	150
N1B—H2NB \cdots Cl1W ⁱⁱ	0.90	2.45	3.2683 (19)	152
N1B—H2NB \cdots O1W	0.90	2.46	3.034 (2)	122
N1B—H3NB \cdots Cl2E ⁱⁱ	0.90	2.57	3.3556 (15)	147
N2B—H1B \cdots Cl1W	0.99	2.20	3.1704 (17)	165
N3B—H2B \cdots O1W ⁱⁱ	0.99	1.98	2.968 (2)	177
N1C—H2NC \cdots Cl3F ^{viii}	0.90	2.68	3.4211 (19)	140
N1C—H3NC \cdots Cl2W ^{ix}	0.90	2.38	3.2673 (17)	167
N1C—H3NC \cdots O2W ^{ix}	0.90	2.54	2.975 (2)	111
N2C—H1C \cdots O2W ^x	0.99	1.96	2.932 (2)	167
N3C—H2C \cdots Cl2W ^x	0.99	2.23	3.2082 (16)	171
C5C—H5C2 \cdots Cl4F	0.98	2.79	3.761 (2)	174
N1D—H2ND \cdots Cl1W ^{xi}	0.90	2.45	3.2796 (15)	154
N1D—H3ND \cdots Cl1E ^{xi}	0.90	2.71	3.5516 (16)	156
N2D—H1D \cdots Cl1W ^{xii}	0.99	2.19	3.1589 (16)	166
N3D—H2D \cdots Cl2E ^{xii}	0.99	2.36	3.3276 (17)	166
C1D—H1D1 \cdots Cl1F ^v	0.98	2.66	3.6278 (17)	168

$C1D-H1D2\cdots C11E^{xi}$	0.98	2.83	3.803 (2)	172
$O1W-H1O1\cdots C11W^{ii}$	0.85 (1)	2.70 (2)	3.341 (2)	134 (2)
$O1W-H2O1\cdots C12F$	0.84 (1)	2.39 (1)	3.2066 (17)	167 (2)
$O2W-H1O2\cdots C13E$	0.83 (1)	2.37 (1)	3.1763 (19)	162 (2)
$O2W-H2O2\cdots C12W$	0.84 (1)	2.55 (2)	3.2009 (19)	135 (2)

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (v) $-x, -y+1, -z+1$; (vi) $x, y, z+1$; (vii) $x-1, y, z+1$; (viii) $x+1, y, z$; (ix) $-x+1, -y+1, -z$; (x) $x, y+1, z$; (xi) $-x+1, -y, -z+1$; (xii) $x-1, y, z$.