



Received 9 October 2016
Accepted 20 October 2016

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Keywords: crystal structure; benzofuran; carbazole derivatives; inversion dimers; aggregation; C—H···O hydrogen bonds; C—H···π interactions.

CCDC references: 1479200; 1479199;
1479198

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structures of three carbazole derivatives: 12-ethyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]-carbazole, (1), 2-(4,5-dimethoxy-2-nitrophenyl)-4-hydroxy-9-phenylsulfonyl-9*H*-carbazole-3-carbaldehyde, (2), and 12-phenyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole, (3)

Rajeswari Gangadharan,^a P. Narayanan,^b K. Sethusankar,^{b*} Velu Saravanan^c and Arasambattu K. Mohanakrishnan^c


^aDepartment of Physics, Ethiraj College for Women (Autonomous), Chennai 600 008, India, ^bDepartment of Physics, RKM Vivekananda College (Autonomous), Chennai 600 004, India, and ^cDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India. *Correspondence e-mail: ksethusankar@yahoo.co.in

The three title compounds, $C_{26}H_{19}NO_3S$, (1), $C_{27}H_{20}N_2O_8S$, (2), and $C_{30}H_{19}NO_3S$, (3), are carbazole derivatives, where (1) and (3) are heterocycle-containing carbazoles with a benzofuran moiety fused to a carbazole unit. In (2), a dimethoxynitrophenyl ring is attached to the carbazole moiety. In the three derivatives, a phenylsulfonyl group is attached to the N atom of the carbazole unit. Compound (1) crystallizes with two independent molecules in the asymmetric unit (*A* and *B*). The carbazole skeleton in the three compounds is essentially planar. In compound (1), the benzene ring of the phenylsulfonyl moiety is almost orthogonal to the carbazole moiety, with dihedral angles of 85.42 (9) and 84.52 (9) $^\circ$ in molecules *A* and *B*, respectively. The benzene ring of the phenylsulfonyl group in compounds (2) and (3) are inclined to the carbazole moiety, making dihedral angles of 70.73 (13) and 81.73 (12) $^\circ$, respectively. The S atom has a distorted tetrahedral configuration in all three compounds. In the crystals, C—H···O hydrogen bonds give rise to $R_2^2(12)$ inversion dimers for compound (1), and to $R_2^2(24)$ inversion dimers and $R_4^4(40)$ ring motifs for compound (2). The crystal packing in (1) also features C—H···π and π—π interactions [shortest intercentroid distance = 3.684 (1) Å], leading to supramolecular three-dimensional aggregation. In the crystal of compound (2), the combination of the various C—H···O hydrogen bonds leads to the formation of a three-dimensional network. In the crystal of compound (3), molecules are linked by C—H···O hydrogen bonds, forming chains running parallel to the *a* axis, and the chains are linked by C—H···π interactions, forming corrugated sheets parallel to the *ab* plane.

1. Chemical context


Carbazoles are widely used as building blocks for new organic materials and play an important role in electroactive and photoactive devices. Carbazole derivatives have also been used as luminescent and hole-transporting materials (Dijken *et al.*, 2004). These compounds are also thermally and photochemically stable which makes them useful materials for technological applications (Diaz *et al.*, 2002).

Heterocycle-containing carbazole derivatives are embodied in many natural products (Itoigawa *et al.*, 2000) and display a broad spectrum of useful biological activities, such as anti-tumour, antimitotic and antioxidative activities (Prudhomme, 2003; Tachibana *et al.*, 2003; Hu *et al.*, 2006). A number of



OPEN ACCESS

benzo-annulated carbazole ring systems containing an aromatic ring fused to the carbazole nucleus are potential candidates for cancer treatment as a result of their DNA intercalative binding properties. They have been shown to bind to estrogen receptors and exhibit a pronounced anti-tumor activity against leukemia, renal tumor, colon cancer and malignant melanoma tumor cell lines (Pindur & Lemster, 1997).



Most heterocycle-containing carbazoles reported in the literature comprise a common heterocyclic ring moiety fused with a carbazole ring system, such as pyridocarbazoles and indolocarbazoles. In this context, we discuss here three carbazole derivatives, two of which have benzofuran moieties fused with the carbazole unit.

2. Structural commentary

The three title compounds $C_{26}H_{19}NO_3S$, (1), $C_{27}H_{20}N_2O_8S$, (2), and $C_{30}H_{19}NO_3S$, (3), are carbazole derivatives, where (1) and (3) are heterocycle-containing carbazoles with a benzofuran fused to the carbazole skeleton (Figs. 1 and 3, respectively). In (2), a dimethoxynitrophenyl ring is attached to the carbazole moiety (Fig. 2). In the three derivatives, a phenylsulfonyl group is attached to the N atom of the carbazole unit. Compound (1) crystallizes with two independent molecules (*A* and *B*) in the asymmetric unit, as shown in Fig. 1. The carbazole skeleton in the three compounds is essentially planar [maximum deviations of 0.052 (2) Å for atom C12 in molecule *A* and 0.080 (2) Å for atom C12' in molecule *B* of (1), -0.034 (2) Å for atom C10 in (2), and -0.042 (4) Å for atom C3 in (3)]. The carbazole benzofuran fused pentacyclic unit is almost planar in (1) and (3), with dihedral angles between the benzofuran and carbazole units being 2.48 (6) and 4.16 (6)° in molecules *A* and *B*, respectively of (1), and




Figure 1
The molecular structure of compound (1), showing the atom-numbering scheme and displacement ellipsoids drawn at the 30% probability level. The intramolecular C—H···O hydrogen bonds, which generate two S(6) ring motifs, are shown as dashed lines (see Table 1).

Table 1
Hydrogen-bond geometry (\AA , $^\circ$) for (1).

Cg1, *Cg4*, *Cg6*, *Cg17* and *Cg20* are the centroids of rings O3/C10/C11/C13/C14, C7–C12, C21–C26, O3'/C10'/C11'/C13'/C14' and C7'-C12', respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4—H4···O2	0.93	2.35	2.954 (3)	122
C4'—H4'···O1'	0.93	2.36	2.966 (3)	122
C9—H9···O1	0.93	2.29	2.881 (2)	121
C9'—H9'···O2'	0.93	2.28	2.875 (2)	121
C4—H4···O2 ⁱ	0.93	2.53	3.277 (3)	137
C20'—H20A···Cg17 ⁱⁱ	0.96	2.82	3.449 (3)	124
C20'—H20C···Cg20 ⁱⁱ	0.96	2.79	3.427 (3)	125
C20—H20D···Cg4 ⁱⁱⁱ	0.96	2.83	3.464 (3)	124
C20'—H20C···Cg1 ⁱⁱⁱ	0.96	2.85	3.478 (3)	124
C25'—H25'···Cg6 ^{iv}	0.93	2.90	3.762 (3)	155

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x, -y+1, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x-1, y, z$.

2.33 (8)° in compound (3). In compound (1), the benzene ring of the phenylsulfonyl group is almost orthogonal to the carbazole moiety, with the dihedral angles between their mean planes being 85.42 (9) and 84.52 (9)° in molecules *A* and *B*, respectively. The benzene ring of the phenylsulfonyl group in compounds (2) and (3) are inclined to the carbazole moiety making dihedral angles of 70.73 (12) and 81.73 (12)°, respectively.

In all three compounds, there are two intramolecular C—H···O hydrogen bonds, involving the sulfonyl ring O atoms forming two cyclic $S(6)$ motifs (Tables 1, 2 and 3). In compound (2), an O—H···O hydrogen bond generates an additional $S(6)$ ring motif (Table 2). Atom S1 has a distorted tetrahedral geometry in all three compounds. The widening of angle O2=S1=O1 [119.55 (10) and 119.46 (10) $^{\circ}$ in molecules *A* and *B*, respectively, of (1), 119.78 (10) $^{\circ}$ in (2) and

Table 2

Hydrogen-bond geometry (\AA , $^\circ$) for (2).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3A…O4	0.82	1.83	2.554 (3)	146
C4—H4…O1	0.93	2.29	2.866 (3)	119
C9—H9…O2	0.93	2.47	3.054 (2)	121
C2—H2…O5 ⁱ	0.93	2.50	3.281 (3)	142
C17—H17…O8 ⁱⁱ	0.93	2.59	3.481 (5)	161
C18—H18…O2 ⁱⁱⁱ	0.93	2.51	3.384 (5)	157
C26—H26C…O4 ^{iv}	0.96	2.50	3.265 (4)	137

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (iv) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Table 3

Hydrogen-bond geometry (\AA , $^\circ$) for (3).

Cg3 and Cg4 are the centroids of rings C1–C6 and C7–C12, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4—H4…O1	0.93	2.34	2.924 (4)	121
C9—H9…O2	0.93	2.34	2.926 (3)	121
C2—H2…O3 ⁱ	0.93	2.57	3.464 (4)	160
C17—H17…Cg4 ⁱⁱ	0.93	2.81	3.683 (3)	156
C22—H22…Cg3 ⁱⁱⁱ	0.93	2.95	3.722 (3)	141

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$.

119.99 (13) $^\circ$ in (3)] and narrowing of angle N—S—C [104.85 (9) and 104.82 (9) $^\circ$ in molecules *A* and *B*, respectively, of (1), 102.92 (9) $^\circ$ in (2) and 105.79 (12) $^\circ$ in (3)] from the ideal tetrahedral value are attributed to the Thorpe–Ingold effect (Bassindale, 1984). As a result of the electron-withdrawing character of the phenylsulfonyl group, the bond lengths N1—C5 [1.430 (2) and 1.431 (2) \AA in molecules *A* and *B* of (1), 1.429 (3) \AA in (2) and 1.432 (4) \AA in (3)] and N1—C8 [1.428 (2) and 1.425 (2) \AA in molecules *A* and *B* of (1), 1.414 (2) \AA in (2) and 1.432 (3) \AA in (3)] in all three compounds are longer than the normal value of 1.355 (14) \AA [Cambridge Structural Database (CSD), Version 5.37; last update May 2016.; Groom *et al.*, 2016].




Figure 2

The molecular structure of compound (2), showing the atom-numbering scheme and displacement ellipsoids are drawn at the 30% probability level. The intramolecular O—H…O and C—H…O hydrogen bonds, which generate three S(6) ring motifs, are shown as dashed lines (see Table 2).




Figure 3

The molecular structure of compound (3), showing the atom-numbering scheme and displacement ellipsoids drawn at the 30% probability level. The intramolecular C—H…O hydrogen bonds, which generate two S(6) ring motifs, are shown as dashed lines (see Table 3).


In compound (2), the dimethoxynitrophenyl ring makes a dihedral angle of 76.63 (8) $^\circ$ with the carbazole moiety. The nitro group in (2) is (+) syn-periplanar to the phenyl ring (atoms C20–C25), as indicated by the values of the torsion angles C24—C25—N2—O6 = 21.4 (3) $^\circ$ and C20—C25—N2—O5 = 19.9 (3) $^\circ$. The torsion angles C22—C23—O8—C27 = -174.6 (2) $^\circ$ and C23—C22—O7—C26 = 175.9 (2) $^\circ$ indicate that the two methoxy substituents at C23 and C22 are almost coplanar with the phenyl ring.

In compound (3), the phenyl ring attached at C12 is oriented at a dihedral angle of 78.39 (11) $^\circ$ to the carbazole unit.

3. Supramolecular features

In the crystal of compound (1), molecules are linked via C4—H4…O2 and C4’—H4’…O1’ hydrogen bonds, generating two $R_2^2(12)$ inversion dimers (Table 1 and Fig. 4). The crystal packing also features C—H… π (Table 1) and π — π interactions leading to supramolecular three-dimensional aggregation. The π — π interactions involve inversion related *A* molecules with an intercentroid distance $Cg4\cdots Cg4^i$ = 3.703 (2) \AA [where $Cg4$ is the centroid of ring C7–C12; symmetry code: (i) $-x + 2, -y + 1, -z + 1$], and inversion related *B* molecules, with an intercentroid distance $Cg20\cdots Cg20^{ii}$ = 3.684 (2) \AA [where $Cg20$ is the centroid of ring C7’–C12’; symmetry code: (ii) $-x + 1, -y + 1, -z$].

In the crystal of compound (2), neighbouring molecules are linked by C18—H18…O2ⁱⁱⁱ and C26—H26C…O4^{iv} hydrogen bonds forming $R_4^4(40)$ ring motifs resulting in the formation of sheets parallel to the *bc* plane (Table 2 and Fig. 5). Molecules

**Figure 4**


The crystal packing of compound (1), viewed along the a -axis, showing the formation of centrosymmetric A–A dimers, with descriptor $R_2^2(12)$. The dashed lines indicate the intermolecular C–H \cdots O hydrogen bonds (Table 1) and H atoms not involved in hydrogen bonding, and the phenyl ring of the phenylsulfonate groups, have been excluded for clarity.

are also linked via $C2-H2\cdots O5^i$ hydrogen bonds which form $R_2^2(24)$ inversion dimers. These dimers are further crosslinked by $C17-H17\cdots O8^{ii}$ hydrogen bonds (Table 2), forming sheets parallel to plane (102); as shown in Fig. 6. The sum of these interactions is the formation of a three-dimensional hydrogen-bonded framework.

In the crystal of compound (3), molecules are linked through $C2-H2\cdots O3^i$ hydrogen bonds (Table 3), that generate infinite one-dimensional $C(9)$ chains running parallel to the a axis (Fig. 7). The chains are further crosslinked by $C17-H17\cdots Cg4^{ii}$ and $C22-H22\cdots Cg3^{iii}$ interactions (Table 3), which results in the formation corrugated sheets parallel to the ab plane.

4. Database survey

A search of the CSD (Groom *et al.*, 2016) revealed two closely related structures including the parent compound 7*H*-1-benzofuro[2,3-*b*]carbazole (Panchatcharam *et al.*, 2011a). This carbazole–benzofuran fused pentacyclic unit crystallizes in the space group $Pca_{2}1$. However, compound 7-phenylsulfonyl-7*H*-benzofuran[2,3-*b*]carbazole (Panchatcharam *et al.*, 2011b) is the closest analogue to the title compounds (1) and (3), and crystallizes in the space group $P2_1/c$. The presence of an ethyl or phenyl substituent attached to the carbazole unit does not cause much variation in the structural parameters. The packing of the title compounds are consolidated by C–H \cdots O

**Figure 5**

The crystal packing of compound (2), viewed along the a axis, showing the formation of $R_4^4(40)$ graph-set ring motifs, resulting in the formation of sheets parallel to the bc plane. The dashed lines indicate the C–H \cdots O hydrogen bonds (Table 2), and H atoms not involved in the hydrogen bonding have been excluded for clarity.

Table 4
Experimental details.

	(1)	(2)	(3)
Crystal data			
Chemical formula	C ₂₆ H ₁₉ NO ₃ S	C ₂₇ H ₂₀ N ₂ O ₈ S	C ₃₀ H ₁₉ NO ₃ S
M _r	425.48	532.51	473.52
Crystal system, space group	Triclinic, P\bar{1}	Monoclinic, P2 ₁ /c	Orthorhombic, P2 ₁ 2 ₁ 2 ₁
Temperature (K)	296	296	296
a, b, c (Å)	8.3037 (2), 14.3468 (3), 18.4068 (5)	11.2133 (3), 14.5811 (4), 15.1509 (4)	10.6461 (10), 11.8994 (11), 18.2418 (16)
α, β, γ (°)	70.594 (1), 78.139 (1), 85.356 (1)	90, 102.320 (1), 90	90, 90, 90
V (Å ³)	2023.90 (8)	2420.16 (11)	2310.9 (4)
Z	4	4	4
Radiation type	Mo K α	Mo K α	Mo K α
μ (mm ⁻¹)	0.19	0.19	0.17
Crystal size (mm)	0.35 × 0.30 × 0.25	0.35 × 0.30 × 0.25	0.35 × 0.30 × 0.25
Data collection			
Diffractometer	Bruker Kappa APEXII CCD	Bruker Kappa APEXII CCD	Bruker Kappa APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2008)	Multi-scan (<i>SADABS</i> ; Bruker, 2008)	Multi-scan (<i>SADABS</i> ; Bruker, 2008)
T _{min} , T _{max}	0.936, 0.954	0.935, 0.953	0.941, 0.957
No. of measured, independent and observed [I > 2σ(I)] reflections	32388, 8968, 7032	42635, 6535, 4517	29269, 5055, 3194
R _{int}	0.022	0.025	0.061
(sin θ/λ) _{max} (Å ⁻¹)	0.643	0.690	0.640
Refinement			
R[F ² > 2σ(F ²)], wR(F ²), S	0.046, 0.125, 1.01	0.047, 0.150, 1.00	0.042, 0.101, 1.00
No. of reflections	8968	6535	5055
No. of parameters	561	345	316
No. of restraints	0	0	1
H-atom treatment	H-atom parameters constrained	H-atom parameters constrained	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.67, -0.37	0.33, -0.46	0.16, -0.25
Absolute structure	—	—	Flack (1983), 2189 Friedel pairs
Absolute structure parameter	—	—	0.08 (9)

Computer programs: *APEX2* (Bruker, 2008), *SAINT* (Bruker, 2008), *SHELXS97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia, 2012) and *Mercury* (Macrae *et al.*, 2008), *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).





Figure 6

The crystal packing of compound (2), viewed normal to plane (2̄04), showing the formation of R₂²(24) graph-set ring motifs, resulting in the formation of sheets parallel to plane (2̄04). The dashed lines indicate the intermolecular C—H···O hydrogen bonds (Table 2), and H atoms not involved in the hydrogen bonding have been excluded for clarity.

**Figure 7**

The crystal packing of compound (3), viewed along the *b* axis, showing the C–H···O hydrogen bonds (dashed lines; Table 3), which generate *C*(9) chains running parallel to the *a* axis. H atoms not involved in the hydrogen bonding, and the phenylsulfonate groups, have been excluded for clarity.

interactions, but the related compounds exhibit only C–H···π and π–π interactions.

A similar search conducted for compound (2) gave 10 hits of compounds having a phenyl ring attached to a 7-phenylsulfonyl-7*H*-benzofuran[2,3-*b*]carbazole skeleton. The closest analogues to compound (2) are 2-(4,5-dimethoxy-2-nitrophenyl)-4-methoxy-3-methyl-9-phenylsulfonyl-9*H*-carbazole (Narayanan *et al.*, 2014a) and 2-(4,5-dimethoxy-2-nitrophenyl)-4-methoxy-9-phenylsulfonyl-9*H*-carbazole-3-carbaldehyde (Narayanan *et al.*, 2014b). Both crystallize in the space group *Pca*2₁, and differ from compound (2) only in the groups attached to the substituted phenyl ring of the carbazole moiety.

5. Synthesis and crystallization

For the preparation of compound (1), a solution of [1-(phenylsulfonyl)-3-propionyl-1*H*-indol-2-yl]methyl pivalate (0.1 g, 2.34 mmol), anhydrous SnCl₄ (0.07 g, 2.81 mmol) and benzofuran (0.033 g, 2.81 mmol) in dry DCE (10 ml) was stirred at room temperature under a nitrogen atmosphere for 3 h. After the completion of the reaction (monitored by thin-layer chromatography, TLC), it was poured into ice water (100 ml), the organic layer was separated and the aqueous layer was extracted with DCM (2 × 20 ml). The combined extract was washed with water (3 × 50 ml) and dried (Na₂SO₄). Removal of solvent followed by column chromatographic purification (silica gel; hexane–ethyl acetate, 8:2 *v/v*) led to the isolation of compound (1) as a colourless solid (yield 0.064 g, 64%; m.p. 483–485 K).

For the preparation of compound (2), to a solution of 4-methoxycarbazole-3-carbaldehydes (0.82 g, 1.5 mmol) in dry DCM (20 ml), 1 *M* solution of BBr₃ (1.65 ml, 1.65 mmol) in DCM was added at 273 K. After completion of the reaction (monitored by TLC), it was poured into ice water (50 ml) containing HCl (5 ml). The organic layer was separated and the aqueous layer was then extracted with DCM (2 × 10 ml). The combined organic layer was washed with water (2 × 30 ml) and dried (Na₂SO₄). Removal of the solvent followed by tritura-

tion of the crude product with MeOH (10 ml) gave compound (2) as a pale-yellow solid (yield 0.73 g, 92%; m.p. 467–469 K).

For the preparation of compound (3), a solution of [3-benzoyl-1-(phenylsulfonyl)-1*H*-indol-2-yl]methyl pivalate (0.1 g, 2.11 mmol), anhydrous SnCl₄ (0.066 g, 2.52 mmol) and benzofuran (0.03 g, 2.52 mmol) in dry DCE (10 ml) was stirred at room temperature under a nitrogen atmosphere for 3 h. After the completion of the reaction (monitored by TLC), it was poured into ice water (100 ml), the organic layer was separated and the aqueous layer was extracted with DCM (2 × 20 ml). The combined extract was washed with water (3 × 50 ml) and dried (Na₂SO₄). Removal of solvent followed by column chromatographic purification (silica gel; hexane–ethyl acetate, 8:2 *v/v*) gave compound (3) as a colourless solid (yield 0.07 g, 70%; m.p. 491–493 K). Colourless block-like crystals were obtained by slow evaporation of a solution of (1) and (3) in ethyl acetate. Yellow block-like crystals were obtained by slow evaporation of a solution of (2) in methanol.

6. Refinement

Crystal data, data collection and structure refinement details for compounds (1), (2) and (3) are summarized in Table 4. The H atoms were included in calculated positions and treated as riding atoms: O–H = 0.82 Å, C–H = 0.93–0.97 Å, with *U*_{iso}(H) = 1.5*U*_{eq}(hydroxy O and methyl C) and 1.2*U*_{eq}(C) for other H atoms. The methyl groups were allowed to rotate, but not to tip, to best fit the electron density.

Acknowledgements

The authors thank Dr Babu Varghese, Senior Scientific Officer, SAIF, IIT Madras, Chennai, India, for the data collections.

References

- Bassindale, A. (1984). *The Third Dimension in Organic Chemistry*, ch. 1, p. 11. New York: John Wiley and Sons.
- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.

- Diaz, J. L., Villacampa, B., Lopez-Calahorra, F. & Velasco, D. (2002). *Chem. Mater.* **14**, 2240–2251.
- Dijken, A. V., Bastiaansen, J. J. A. M., Kiggen, N. M. M., Langeveld, B. M. W., Rothe, C., Monkman, A., Bach, I., Stössel, P. & Brunner, K. (2004). *J. Am. Chem. Soc.* **126**, 7718–7727.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hu, L., Li, Z., Li, Y., Qu, J., Ling, Y.-H., Jiang, J. & Boykin, D. W. (2006). *J. Med. Chem.* **49**, 6273–6282.
- Itoigawa, M., Kashiwada, Y., Ito, C., Furukawa, H., Tachibana, Y., Bastow, K. F. & Lee, K.-H. (2000). *J. Nat. Prod.* **63**, 893–897.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Narayanan, P., Sethusankar, K., Saravanan, V. & Mohanakrishnan, A. K. (2014a). *Acta Cryst.* **E70**, o336–o337.
- Narayanan, P., Sethusankar, K., Saravanan, V. & Mohanakrishnan, A. K. (2014b). *Acta Cryst.* **E70**, o424–o425.
- Panchatcharam, R., Dhayalan, V., Mohanakrishnan, A. K., Chakkavarthi, G. & Manivannan, V. (2011a). *Acta Cryst.* **E67**, o2797.
- Panchatcharam, R., Dhayalan, V., Mohanakrishnan, A. K., Chakkavarthi, G. & Manivannan, V. (2011b). *Acta Cryst.* **E67**, o2829.
- Pindur, U. & Lemster, T. (1997). In *Recent Research Developments in Organic and Bioorganic Chemistry*, Vol. 1, edited by S. G. Pandalai, pp. 33–54. Trivandrum India: Transworld Research Network.
- Prudhomme, M. (2003). *Eur. J. Med. Chem.* **38**, 123–140.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tachibana, Y., Kikuzaki, H., Lajis, N. H. & Nakatani, N. (2003). *J. Agric. Food Chem.* **51**, 6461–6467.

supporting information

Acta Cryst. (2016). E72, 1744-1750 [https://doi.org/10.1107/S2056989016016819]

Crystal structures of three carbazole derivatives: 12-ethyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole, (1), 2-(4,5-dimethoxy-2-nitrophenyl)-4-hydroxy-9-phenylsulfonyl-9*H*-carbazole-3-carbaldehyde, (2), and 12-phenyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole, (3)

Rajeswari Gangadharan, P. Narayanan, K. Sethusankar, Velu Saravanan and Arasambattu K. Mohanakrishnan

Computing details

For all compounds, data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

(1) 12-Ethyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole

Crystal data

C₂₆H₁₉NO₃S
 $M_r = 425.48$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 8.3037 (2)$ Å
 $b = 14.3468 (3)$ Å
 $c = 18.4068 (5)$ Å
 $\alpha = 70.594 (1)^\circ$
 $\beta = 78.139 (1)^\circ$
 $\gamma = 85.356 (1)^\circ$
 $V = 2023.90 (8)$ Å³

Z = 4
 $F(000) = 888$
 $D_x = 1.396 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 8968 reflections
 $\theta = 1.5\text{--}27.2^\circ$
 $\mu = 0.19 \text{ mm}^{-1}$
T = 296 K
Block, colourless
0.35 × 0.30 × 0.25 mm

Data collection

Bruker Kappa APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω & φ scans
Absorption correction: multi-scan
(SADABS; Bruker, 2008)
 $T_{\min} = 0.936$, $T_{\max} = 0.954$

32388 measured reflections
8968 independent reflections
7032 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 27.2^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -10 \rightarrow 10$
 $k = -18 \rightarrow 13$
 $l = -23 \rightarrow 23$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.046$ $wR(F^2) = 0.125$ $S = 1.01$

8968 reflections

561 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0579P)^2 + 0.955P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.67 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.37 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7000 (3)	0.23489 (15)	0.65728 (12)	0.0518 (5)
H1	0.6502	0.2770	0.6846	0.062*
C1'	0.3087 (3)	0.35673 (16)	-0.15171 (12)	0.0468 (5)
H1'	0.2811	0.4208	-0.1799	0.056*
C2	0.6984 (3)	0.13356 (17)	0.69399 (14)	0.0619 (6)
H2	0.6474	0.1077	0.7463	0.074*
C2'	0.3273 (3)	0.28301 (18)	-0.18562 (13)	0.0558 (6)
H2'	0.3112	0.2977	-0.2368	0.067*
C3'	0.3696 (3)	0.18749 (17)	-0.14461 (14)	0.0561 (6)
H3'	0.3828	0.1393	-0.1690	0.067*
C3	0.7715 (3)	0.07018 (16)	0.65380 (14)	0.0603 (6)
H3	0.7685	0.0024	0.6796	0.072*
C4'	0.3926 (3)	0.16229 (15)	-0.06827 (13)	0.0481 (5)
H4'	0.4215	0.0982	-0.0408	0.058*
C4	0.8488 (3)	0.10544 (14)	0.57628 (13)	0.0507 (5)
H4	0.8985	0.0628	0.5494	0.061*
C5'	0.3710 (2)	0.23655 (14)	-0.03417 (11)	0.0375 (4)
C5	0.8495 (2)	0.20709 (13)	0.53989 (11)	0.0395 (4)
C6'	0.3317 (2)	0.33431 (13)	-0.07473 (10)	0.0355 (4)
C6	0.7773 (2)	0.27308 (13)	0.57885 (11)	0.0390 (4)
C7'	0.3200 (2)	0.39194 (13)	-0.02196 (10)	0.0316 (4)
C7	0.8033 (2)	0.37338 (13)	0.52354 (10)	0.0351 (4)
C8	0.8888 (2)	0.36459 (12)	0.45138 (10)	0.0329 (4)
C8'	0.3470 (2)	0.32578 (12)	0.05125 (10)	0.0319 (4)
C9	0.9360 (2)	0.44497 (13)	0.38509 (10)	0.0354 (4)

H9	0.9938	0.4388	0.3379	0.043*
C9'	0.3411 (2)	0.35445 (13)	0.11646 (10)	0.0351 (4)
H9'	0.3593	0.3108	0.1644	0.042*
C10'	0.3060 (2)	0.45326 (13)	0.10416 (10)	0.0331 (4)
C10	0.8897 (2)	0.53486 (12)	0.39514 (10)	0.0337 (4)
C11	0.8055 (2)	0.54828 (13)	0.46460 (10)	0.0345 (4)
C11'	0.2811 (2)	0.52240 (12)	0.03310 (10)	0.0312 (4)
C12'	0.2920 (2)	0.49311 (12)	-0.03285 (10)	0.0311 (4)
C12	0.7653 (2)	0.46682 (13)	0.53194 (11)	0.0382 (4)
C13'	0.2429 (2)	0.61493 (13)	0.04986 (11)	0.0356 (4)
C13	0.7802 (2)	0.65476 (13)	0.44577 (11)	0.0393 (4)
C14'	0.2500 (2)	0.59338 (13)	0.12880 (11)	0.0391 (4)
C14	0.8538 (2)	0.69530 (13)	0.36753 (11)	0.0395 (4)
C15'	0.2182 (3)	0.66080 (16)	0.16837 (13)	0.0537 (5)
H15'	0.2245	0.6433	0.2211	0.064*
C15	0.8558 (3)	0.79472 (14)	0.32618 (13)	0.0501 (5)
H15	0.9080	0.8191	0.2739	0.060*
C16	0.7764 (4)	0.85641 (15)	0.36614 (14)	0.0643 (7)
H16	0.7735	0.9241	0.3403	0.077*
C16'	0.1766 (3)	0.75549 (17)	0.12644 (14)	0.0639 (7)
H16'	0.1543	0.8034	0.1511	0.077*
C17	0.7011 (4)	0.81913 (16)	0.44405 (15)	0.0712 (8)
H17	0.6484	0.8623	0.4697	0.085*
C17'	0.1677 (3)	0.78012 (16)	0.04815 (15)	0.0648 (7)
H17'	0.1395	0.8445	0.0210	0.078*
C18	0.7025 (3)	0.71908 (16)	0.48456 (13)	0.0567 (6)
H18	0.6520	0.6951	0.5371	0.068*
C18'	0.1997 (3)	0.71123 (15)	0.00921 (13)	0.0496 (5)
H18'	0.1925	0.7289	-0.0434	0.060*
C19'	0.2730 (2)	0.56711 (13)	-0.11114 (10)	0.0359 (4)
H19A	0.3424	0.5469	-0.1527	0.043*
H19B	0.3100	0.6311	-0.1145	0.043*
C19	0.6875 (3)	0.48089 (15)	0.60964 (12)	0.0453 (5)
H19C	0.7244	0.4282	0.6520	0.054*
H19D	0.7226	0.5432	0.6108	0.054*
C20'	0.0955 (3)	0.57674 (17)	-0.12336 (13)	0.0525 (5)
H20A	0.0600	0.5143	-0.1225	0.079*
H20B	0.0887	0.6256	-0.1731	0.079*
H20C	0.0262	0.5964	-0.0822	0.079*
C20	0.5034 (3)	0.48049 (19)	0.62191 (15)	0.0628 (6)
H20D	0.4666	0.5338	0.5809	0.094*
H20E	0.4572	0.4886	0.6717	0.094*
H20F	0.4685	0.4188	0.6210	0.094*
C21	0.7629 (2)	0.19838 (14)	0.37307 (11)	0.0411 (4)
C21'	0.1572 (2)	0.09554 (13)	0.13654 (11)	0.0376 (4)
C22'	0.1101 (3)	0.02676 (15)	0.10739 (13)	0.0489 (5)
H22'	0.1882	-0.0043	0.0788	0.059*
C22	0.6784 (3)	0.28012 (17)	0.33250 (13)	0.0511 (5)

H22	0.7243	0.3426	0.3143	0.061*
C23	0.5248 (3)	0.2671 (2)	0.31958 (15)	0.0630 (6)
H23	0.4661	0.3211	0.2929	0.076*
C23'	-0.0548 (3)	0.00497 (18)	0.12142 (15)	0.0598 (6)
H23'	-0.0880	-0.0417	0.1027	0.072*
C24'	-0.1696 (3)	0.05186 (18)	0.16283 (15)	0.0622 (6)
H24'	-0.2804	0.0372	0.1715	0.075*
C24	0.4587 (3)	0.1738 (2)	0.34630 (16)	0.0686 (7)
H24	0.3558	0.1653	0.3371	0.082*
C25'	-0.1225 (3)	0.12049 (18)	0.19168 (14)	0.0595 (6)
H25'	-0.2014	0.1519	0.2196	0.071*
C25	0.5434 (3)	0.0934 (2)	0.38642 (16)	0.0681 (7)
H25	0.4974	0.0310	0.4041	0.082*
C26'	0.0421 (3)	0.14267 (15)	0.17911 (12)	0.0488 (5)
H26'	0.0750	0.1884	0.1989	0.059*
C26	0.6966 (3)	0.10474 (17)	0.40068 (14)	0.0545 (5)
H26	0.7540	0.0506	0.4282	0.065*
N1	0.9229 (2)	0.26237 (11)	0.46133 (9)	0.0378 (3)
N1'	0.38545 (19)	0.23058 (11)	0.04342 (9)	0.0366 (3)
O1	1.04367 (18)	0.28737 (10)	0.32125 (9)	0.0513 (4)
O1'	0.46157 (18)	0.05262 (10)	0.09012 (9)	0.0534 (4)
O2'	0.40205 (19)	0.14614 (10)	0.18312 (9)	0.0520 (4)
O2	1.03320 (18)	0.12012 (10)	0.41762 (10)	0.0551 (4)
O3'	0.29021 (17)	0.49510 (9)	0.16273 (7)	0.0414 (3)
O3	0.92329 (17)	0.62339 (9)	0.33564 (7)	0.0411 (3)
S1	0.95872 (6)	0.21460 (3)	0.38890 (3)	0.04001 (13)
S1'	0.36624 (6)	0.12527 (3)	0.11770 (3)	0.03903 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0642 (14)	0.0404 (11)	0.0413 (11)	-0.0003 (10)	-0.0025 (10)	-0.0054 (9)
C1'	0.0568 (12)	0.0498 (11)	0.0361 (10)	-0.0062 (9)	-0.0102 (9)	-0.0147 (9)
C2	0.0790 (17)	0.0445 (12)	0.0454 (12)	-0.0047 (11)	-0.0017 (11)	0.0024 (10)
C2'	0.0675 (15)	0.0655 (15)	0.0409 (11)	-0.0130 (11)	-0.0092 (10)	-0.0237 (11)
C3'	0.0646 (14)	0.0575 (13)	0.0536 (13)	-0.0127 (11)	0.0009 (11)	-0.0322 (11)
C3	0.0820 (17)	0.0326 (10)	0.0551 (14)	-0.0020 (10)	-0.0134 (12)	0.0011 (10)
C4'	0.0541 (12)	0.0416 (11)	0.0498 (12)	-0.0054 (9)	-0.0018 (10)	-0.0201 (9)
C4	0.0675 (14)	0.0319 (10)	0.0512 (12)	0.0036 (9)	-0.0141 (10)	-0.0105 (9)
C5'	0.0358 (9)	0.0397 (10)	0.0380 (10)	-0.0059 (7)	-0.0034 (8)	-0.0147 (8)
C5	0.0461 (11)	0.0326 (9)	0.0381 (10)	-0.0009 (8)	-0.0118 (8)	-0.0069 (8)
C6'	0.0317 (9)	0.0410 (10)	0.0339 (9)	-0.0064 (7)	-0.0030 (7)	-0.0127 (8)
C6	0.0447 (10)	0.0331 (9)	0.0365 (10)	-0.0006 (8)	-0.0097 (8)	-0.0064 (8)
C7'	0.0280 (8)	0.0362 (9)	0.0296 (8)	-0.0034 (7)	-0.0052 (7)	-0.0089 (7)
C7	0.0395 (10)	0.0325 (9)	0.0318 (9)	0.0000 (7)	-0.0083 (7)	-0.0077 (7)
C8	0.0349 (9)	0.0299 (8)	0.0364 (9)	0.0006 (7)	-0.0101 (7)	-0.0122 (7)
C8'	0.0287 (8)	0.0295 (8)	0.0360 (9)	-0.0027 (6)	-0.0066 (7)	-0.0076 (7)
C9	0.0381 (9)	0.0358 (9)	0.0329 (9)	-0.0002 (7)	-0.0054 (7)	-0.0127 (7)

C9'	0.0392 (9)	0.0324 (9)	0.0323 (9)	-0.0025 (7)	-0.0117 (7)	-0.0052 (7)
C10'	0.0332 (9)	0.0363 (9)	0.0310 (9)	-0.0038 (7)	-0.0081 (7)	-0.0105 (7)
C10	0.0362 (9)	0.0317 (9)	0.0318 (9)	-0.0040 (7)	-0.0076 (7)	-0.0068 (7)
C11	0.0405 (10)	0.0316 (9)	0.0331 (9)	0.0006 (7)	-0.0093 (7)	-0.0115 (7)
C11'	0.0271 (8)	0.0317 (8)	0.0323 (9)	-0.0029 (6)	-0.0051 (7)	-0.0067 (7)
C12'	0.0262 (8)	0.0338 (8)	0.0301 (8)	-0.0033 (6)	-0.0046 (7)	-0.0059 (7)
C12	0.0456 (10)	0.0351 (9)	0.0326 (9)	0.0003 (8)	-0.0071 (8)	-0.0098 (8)
C13'	0.0331 (9)	0.0354 (9)	0.0358 (9)	-0.0028 (7)	-0.0034 (7)	-0.0097 (7)
C13	0.0501 (11)	0.0308 (9)	0.0358 (10)	-0.0010 (8)	-0.0088 (8)	-0.0086 (7)
C14'	0.0438 (10)	0.0340 (9)	0.0371 (10)	-0.0027 (8)	-0.0052 (8)	-0.0093 (8)
C14	0.0485 (11)	0.0341 (9)	0.0377 (10)	-0.0002 (8)	-0.0126 (8)	-0.0115 (8)
C15'	0.0736 (15)	0.0469 (12)	0.0424 (11)	-0.0001 (10)	-0.0061 (10)	-0.0199 (10)
C15	0.0706 (14)	0.0350 (10)	0.0415 (11)	-0.0050 (9)	-0.0123 (10)	-0.0062 (9)
C16	0.106 (2)	0.0285 (10)	0.0549 (14)	0.0030 (11)	-0.0158 (13)	-0.0090 (10)
C16'	0.0921 (19)	0.0433 (12)	0.0568 (14)	0.0079 (12)	-0.0032 (13)	-0.0253 (11)
C17	0.116 (2)	0.0370 (12)	0.0573 (15)	0.0125 (13)	-0.0063 (14)	-0.0199 (11)
C17'	0.0923 (19)	0.0353 (11)	0.0606 (15)	0.0141 (11)	-0.0116 (13)	-0.0126 (10)
C18	0.0833 (17)	0.0397 (11)	0.0430 (12)	0.0056 (11)	-0.0025 (11)	-0.0152 (9)
C18'	0.0629 (13)	0.0377 (10)	0.0436 (11)	0.0062 (9)	-0.0104 (10)	-0.0085 (9)
C19'	0.0340 (9)	0.0389 (9)	0.0301 (9)	-0.0037 (7)	-0.0047 (7)	-0.0050 (7)
C19	0.0513 (12)	0.0428 (10)	0.0399 (10)	0.0003 (9)	-0.0102 (9)	-0.0104 (9)
C20'	0.0413 (11)	0.0633 (14)	0.0463 (12)	0.0031 (10)	-0.0127 (9)	-0.0072 (10)
C20	0.0562 (14)	0.0605 (14)	0.0663 (15)	-0.0004 (11)	-0.0108 (12)	-0.0138 (12)
C21	0.0450 (11)	0.0447 (10)	0.0381 (10)	-0.0006 (8)	-0.0028 (8)	-0.0224 (9)
C21'	0.0449 (10)	0.0280 (8)	0.0351 (9)	-0.0010 (7)	-0.0104 (8)	-0.0023 (7)
C22'	0.0543 (12)	0.0393 (10)	0.0532 (12)	-0.0039 (9)	-0.0105 (10)	-0.0140 (9)
C22	0.0563 (13)	0.0547 (12)	0.0462 (12)	0.0050 (10)	-0.0114 (10)	-0.0218 (10)
C23	0.0580 (14)	0.0840 (18)	0.0567 (14)	0.0142 (13)	-0.0197 (11)	-0.0339 (13)
C23'	0.0594 (14)	0.0546 (13)	0.0662 (15)	-0.0148 (11)	-0.0174 (12)	-0.0141 (12)
C24'	0.0467 (13)	0.0615 (14)	0.0648 (15)	-0.0098 (11)	-0.0130 (11)	0.0011 (12)
C24	0.0477 (13)	0.108 (2)	0.0646 (16)	-0.0109 (14)	-0.0066 (12)	-0.0463 (16)
C25'	0.0515 (13)	0.0601 (14)	0.0533 (13)	0.0051 (11)	0.0016 (10)	-0.0086 (11)
C25	0.0609 (15)	0.0781 (18)	0.0716 (17)	-0.0225 (13)	-0.0026 (13)	-0.0333 (15)
C26'	0.0559 (13)	0.0429 (11)	0.0442 (11)	-0.0001 (9)	-0.0064 (9)	-0.0117 (9)
C26	0.0582 (13)	0.0507 (12)	0.0570 (13)	-0.0068 (10)	-0.0061 (11)	-0.0221 (11)
N1	0.0467 (9)	0.0292 (7)	0.0376 (8)	0.0022 (6)	-0.0092 (7)	-0.0107 (6)
N1'	0.0413 (8)	0.0304 (7)	0.0376 (8)	-0.0015 (6)	-0.0080 (7)	-0.0098 (6)
O1	0.0518 (8)	0.0470 (8)	0.0513 (8)	-0.0048 (6)	0.0090 (7)	-0.0211 (7)
O1'	0.0515 (9)	0.0375 (7)	0.0717 (10)	0.0136 (6)	-0.0170 (7)	-0.0185 (7)
O2'	0.0694 (10)	0.0401 (7)	0.0503 (8)	0.0021 (7)	-0.0325 (7)	-0.0077 (6)
O2	0.0536 (9)	0.0398 (8)	0.0729 (10)	0.0133 (6)	-0.0095 (8)	-0.0241 (7)
O3'	0.0576 (8)	0.0352 (7)	0.0335 (7)	-0.0001 (6)	-0.0126 (6)	-0.0117 (5)
O3	0.0530 (8)	0.0315 (6)	0.0334 (7)	-0.0025 (6)	-0.0019 (6)	-0.0067 (5)
S1	0.0404 (3)	0.0345 (2)	0.0461 (3)	0.00394 (19)	-0.0029 (2)	-0.0185 (2)
S1'	0.0431 (3)	0.0288 (2)	0.0450 (3)	0.00405 (18)	-0.0160 (2)	-0.00813 (19)

Geometric parameters (\AA , $\text{^{\circ}}$)

C1—C2	1.385 (3)	C15'—H15'	0.9300
C1—C6	1.395 (3)	C15—C16	1.378 (3)
C1—H1	0.9300	C15—H15	0.9300
C1'—C2'	1.380 (3)	C16—C17	1.382 (3)
C1'—C6'	1.395 (3)	C16—H16	0.9300
C1'—H1'	0.9300	C16'—C17'	1.381 (3)
C2—C3	1.384 (3)	C16'—H16'	0.9300
C2—H2	0.9300	C17—C18	1.381 (3)
C2'—C3'	1.384 (3)	C17—H17	0.9300
C2'—H2'	0.9300	C17'—C18'	1.382 (3)
C3'—C4'	1.380 (3)	C17'—H17'	0.9300
C3'—H3'	0.9300	C18—H18	0.9300
C3—C4	1.378 (3)	C18'—H18'	0.9300
C3—H3	0.9300	C19'—C20'	1.525 (3)
C4'—C5'	1.389 (3)	C19'—H19A	0.9700
C4'—H4'	0.9300	C19'—H19B	0.9700
C4—C5	1.388 (3)	C19—C20	1.499 (3)
C4—H4	0.9300	C19—H19C	0.9700
C5'—C6'	1.401 (3)	C19—H19D	0.9700
C5'—N1'	1.431 (2)	C20'—H20A	0.9600
C5—C6	1.397 (3)	C20'—H20B	0.9600
C5—N1	1.430 (2)	C20'—H20C	0.9600
C6'—C7'	1.455 (2)	C20—H20D	0.9600
C6—C7	1.463 (2)	C20—H20E	0.9600
C7'—C12'	1.405 (2)	C20—H20F	0.9600
C7'—C8'	1.417 (2)	C21—C26	1.385 (3)
C7—C12	1.403 (2)	C21—C22	1.388 (3)
C7—C8	1.412 (2)	C21—S1	1.756 (2)
C8—C9	1.383 (2)	C21'—C22'	1.384 (3)
C8—N1	1.428 (2)	C21'—C26'	1.388 (3)
C8'—C9'	1.383 (2)	C21'—S1'	1.757 (2)
C8'—N1'	1.424 (2)	C22'—C23'	1.382 (3)
C9—C10	1.376 (2)	C22'—H22'	0.9300
C9—H9	0.9300	C22—C23	1.383 (3)
C9'—C10'	1.376 (2)	C22—H22	0.9300
C9'—H9'	0.9300	C23—C24	1.379 (4)
C10'—O3'	1.378 (2)	C23—H23	0.9300
C10'—C11'	1.398 (2)	C23'—C24'	1.371 (4)
C10—O3	1.378 (2)	C23'—H23'	0.9300
C10—C11	1.391 (2)	C24'—C25'	1.378 (4)
C11—C12	1.394 (2)	C24'—H24'	0.9300
C11—C13	1.457 (2)	C24—C25	1.374 (4)
C11'—C12'	1.395 (2)	C24—H24	0.9300
C11'—C13'	1.457 (2)	C25'—C26'	1.383 (3)
C12'—C19'	1.509 (2)	C25'—H25'	0.9300
C12—C19	1.515 (3)	C25—C26	1.384 (3)

C13'—C18'	1.393 (3)	C25—H25	0.9300
C13'—C14'	1.394 (3)	C26'—H26'	0.9300
C13—C14	1.388 (3)	C26—H26	0.9300
C13—C18	1.389 (3)	N1—S1	1.6573 (16)
C14'—C15'	1.372 (3)	N1'—S1'	1.6596 (15)
C14'—O3'	1.383 (2)	O1—S1	1.4257 (15)
C14—C15	1.375 (3)	O1'—S1'	1.4225 (14)
C14—O3	1.381 (2)	O2'—S1'	1.4245 (15)
C15'—C16'	1.377 (3)	O2—S1	1.4219 (14)
C2—C1—C6	119.2 (2)	C15'—C16'—H16'	119.6
C2—C1—H1	120.4	C17'—C16'—H16'	119.6
C6—C1—H1	120.4	C18—C17—C16	121.4 (2)
C2'—C1'—C6'	119.4 (2)	C18—C17—H17	119.3
C2'—C1'—H1'	120.3	C16—C17—H17	119.3
C6'—C1'—H1'	120.3	C16'—C17'—C18'	121.5 (2)
C3—C2—C1	120.9 (2)	C16'—C17'—H17'	119.2
C3—C2—H2	119.6	C18'—C17'—H17'	119.2
C1—C2—H2	119.6	C17—C18—C13	119.1 (2)
C1'—C2'—C3'	121.0 (2)	C17—C18—H18	120.5
C1'—C2'—H2'	119.5	C13—C18—H18	120.5
C3'—C2'—H2'	119.5	C17'—C18'—C13'	119.1 (2)
C4'—C3'—C2'	121.3 (2)	C17'—C18'—H18'	120.4
C4'—C3'—H3'	119.3	C13'—C18'—H18'	120.4
C2'—C3'—H3'	119.3	C12'—C19'—C20'	112.29 (15)
C4—C3—C2	121.40 (19)	C12'—C19'—H19A	109.1
C4—C3—H3	119.3	C20'—C19'—H19A	109.1
C2—C3—H3	119.3	C12'—C19'—H19B	109.1
C3'—C4'—C5'	117.3 (2)	C20'—C19'—H19B	109.1
C3'—C4'—H4'	121.3	H19A—C19'—H19B	107.9
C5'—C4'—H4'	121.3	C20—C19—C12	111.13 (18)
C3—C4—C5	117.4 (2)	C20—C19—H19C	109.4
C3—C4—H4	121.3	C12—C19—H19C	109.4
C5—C4—H4	121.3	C20—C19—H19D	109.4
C4'—C5'—C6'	122.65 (18)	C12—C19—H19D	109.4
C4'—C5'—N1'	128.82 (18)	H19C—C19—H19D	108.0
C6'—C5'—N1'	108.53 (15)	C19'—C20'—H20A	109.5
C4—C5—C6	122.67 (18)	C19'—C20'—H20B	109.5
C4—C5—N1	128.59 (18)	H20A—C20'—H20B	109.5
C6—C5—N1	108.72 (15)	C19'—C20'—H20C	109.5
C1'—C6'—C5'	118.27 (17)	H20A—C20'—H20C	109.5
C1'—C6'—C7'	133.70 (18)	H20B—C20'—H20C	109.5
C5'—C6'—C7'	108.01 (15)	C19—C20—H20D	109.5
C1—C6—C5	118.48 (17)	C19—C20—H20E	109.5
C1—C6—C7	133.59 (18)	H20D—C20—H20E	109.5
C5—C6—C7	107.93 (16)	C19—C20—H20F	109.5
C12'—C7'—C8'	120.72 (16)	H20D—C20—H20F	109.5
C12'—C7'—C6'	132.15 (16)	H20E—C20—H20F	109.5

C8'—C7'—C6'	107.11 (15)	C26—C21—C22	121.4 (2)
C12—C7—C8	120.56 (16)	C26—C21—S1	119.59 (17)
C12—C7—C6	132.36 (17)	C22—C21—S1	118.97 (16)
C8—C7—C6	107.03 (15)	C22'—C21'—C26'	121.2 (2)
C9—C8—C7	123.32 (16)	C22'—C21'—S1'	119.35 (16)
C9—C8—N1	127.96 (16)	C26'—C21'—S1'	119.49 (15)
C7—C8—N1	108.64 (15)	C23'—C22'—C21'	119.0 (2)
C9'—C8'—C7'	123.27 (16)	C23'—C22'—H22'	120.5
C9'—C8'—N1'	128.19 (16)	C21'—C22'—H22'	120.5
C7'—C8'—N1'	108.50 (15)	C23—C22—C21	118.9 (2)
C10—C9—C8	114.04 (16)	C23—C22—H22	120.6
C10—C9—H9	123.0	C21—C22—H22	120.6
C8—C9—H9	123.0	C24—C23—C22	120.0 (2)
C10'—C9'—C8'	114.04 (16)	C24—C23—H23	120.0
C10'—C9'—H9'	123.0	C22—C23—H23	120.0
C8'—C9'—H9'	123.0	C24'—C23'—C22'	120.3 (2)
C9'—C10'—O3'	122.56 (15)	C24'—C23'—H23'	119.9
C9'—C10'—C11'	125.35 (16)	C22'—C23'—H23'	119.9
O3'—C10'—C11'	112.09 (15)	C23'—C24'—C25'	120.7 (2)
C9—C10—O3	122.68 (16)	C23'—C24'—H24'	119.6
C9—C10—C11	125.31 (16)	C25'—C24'—H24'	119.6
O3—C10—C11	112.01 (15)	C25—C24—C23	120.7 (2)
C10—C11—C12	120.03 (16)	C25—C24—H24	119.7
C10—C11—C13	105.15 (15)	C23—C24—H24	119.7
C12—C11—C13	134.82 (17)	C24'—C25'—C26'	120.1 (2)
C12'—C11'—C10'	120.06 (16)	C24'—C25'—H25'	120.0
C12'—C11'—C13'	134.90 (16)	C26'—C25'—H25'	120.0
C10'—C11'—C13'	105.02 (15)	C24—C25—C26	120.4 (2)
C11'—C12'—C7'	116.44 (15)	C24—C25—H25	119.8
C11'—C12'—C19'	121.11 (16)	C26—C25—H25	119.8
C7'—C12'—C19'	122.45 (16)	C25'—C26'—C21'	118.9 (2)
C11—C12—C7	116.57 (16)	C25'—C26'—H26'	120.6
C11—C12—C19	120.53 (16)	C21'—C26'—H26'	120.6
C7—C12—C19	122.89 (16)	C25—C26—C21	118.6 (2)
C18'—C13'—C14'	117.35 (17)	C25—C26—H26	120.7
C18'—C13'—C11'	136.85 (18)	C21—C26—H26	120.7
C14'—C13'—C11'	105.77 (15)	C8—N1—C5	107.60 (14)
C14—C13—C18	117.63 (17)	C8—N1—S1	124.36 (12)
C14—C13—C11	105.65 (16)	C5—N1—S1	122.50 (12)
C18—C13—C11	136.71 (18)	C8'—N1'—C5'	107.70 (14)
C15'—C14'—O3'	124.13 (18)	C8'—N1'—S1'	124.55 (13)
C15'—C14'—C13'	124.26 (18)	C5'—N1'—S1'	122.61 (12)
O3'—C14'—C13'	111.60 (16)	C10'—O3'—C14'	105.49 (13)
C15—C14—O3	123.90 (18)	C10—O3—C14	105.43 (14)
C15—C14—C13	124.36 (18)	O2—S1—O1	119.55 (9)
O3—C14—C13	111.74 (15)	O2—S1—N1	106.87 (9)
C14'—C15'—C16'	117.0 (2)	O1—S1—N1	107.03 (8)
C14'—C15'—H15'	121.5	O2—S1—C21	108.71 (9)

C16'—C15'—H15'	121.5	O1—S1—C21	108.85 (10)
C14—C15—C16	116.6 (2)	N1—S1—C21	104.86 (8)
C14—C15—H15	121.7	O1'—S1'—O2'	119.46 (9)
C16—C15—H15	121.7	O1'—S1'—N1'	107.03 (9)
C15—C16—C17	121.0 (2)	O2'—S1'—N1'	106.76 (8)
C15—C16—H16	119.5	O1'—S1'—C21'	108.59 (9)
C17—C16—H16	119.5	O2'—S1'—C21'	109.19 (9)
C15'—C16'—C17'	120.7 (2)	N1'—S1'—C21'	104.82 (8)
C6—C1—C2—C3	−0.3 (4)	C18—C13—C14—O3	−178.70 (19)
C6'—C1'—C2'—C3'	−0.5 (3)	C11—C13—C14—O3	0.0 (2)
C1'—C2'—C3'—C4'	0.8 (4)	O3'—C14'—C15'—C16'	−178.7 (2)
C1—C2—C3—C4	0.2 (4)	C13'—C14'—C15'—C16'	0.1 (3)
C2'—C3'—C4'—C5'	0.2 (3)	O3—C14—C15—C16	178.0 (2)
C2—C3—C4—C5	−0.4 (4)	C13—C14—C15—C16	−1.0 (3)
C3'—C4'—C5'—C6'	−1.6 (3)	C14—C15—C16—C17	0.8 (4)
C3'—C4'—C5'—N1'	179.34 (19)	C14'—C15'—C16'—C17'	0.1 (4)
C3—C4—C5—C6	0.6 (3)	C15—C16—C17—C18	0.0 (5)
C3—C4—C5—N1	178.7 (2)	C15'—C16'—C17'—C18'	0.1 (4)
C2'—C1'—C6'—C5'	−0.8 (3)	C16—C17—C18—C13	−0.5 (4)
C2'—C1'—C6'—C7'	−179.0 (2)	C14—C13—C18—C17	0.3 (4)
C4'—C5'—C6'—C1'	1.9 (3)	C11—C13—C18—C17	−177.8 (2)
N1'—C5'—C6'—C1'	−178.86 (16)	C16'—C17'—C18'—C13'	−0.4 (4)
C4'—C5'—C6'—C7'	−179.43 (17)	C14'—C13'—C18'—C17'	0.5 (3)
N1'—C5'—C6'—C7'	−0.2 (2)	C11'—C13'—C18'—C17'	178.1 (2)
C2—C1—C6—C5	0.5 (3)	C11'—C12'—C19'—C20'	−91.9 (2)
C2—C1—C6—C7	−179.2 (2)	C7'—C12'—C19'—C20'	87.6 (2)
C4—C5—C6—C1	−0.7 (3)	C11—C12—C19—C20	92.2 (2)
N1—C5—C6—C1	−179.08 (18)	C7—C12—C19—C20	−88.9 (2)
C4—C5—C6—C7	179.08 (19)	C26'—C21'—C22'—C23'	−0.2 (3)
N1—C5—C6—C7	0.7 (2)	S1'—C21'—C22'—C23'	−178.99 (16)
C1'—C6'—C7'—C12'	−5.1 (3)	C26—C21—C22—C23	−0.1 (3)
C5'—C6'—C7'—C12'	176.55 (18)	S1—C21—C22—C23	179.50 (16)
C1'—C6'—C7'—C8'	176.2 (2)	C21—C22—C23—C24	−0.5 (3)
C5'—C6'—C7'—C8'	−2.18 (19)	C21'—C22'—C23'—C24'	0.8 (3)
C1—C6—C7—C12	3.5 (4)	C22'—C23'—C24'—C25'	−0.6 (4)
C5—C6—C7—C12	−176.2 (2)	C22—C23—C24—C25	0.6 (4)
C1—C6—C7—C8	−179.1 (2)	C23'—C24'—C25'—C26'	−0.1 (4)
C5—C6—C7—C8	1.2 (2)	C23—C24—C25—C26	0.0 (4)
C12—C7—C8—C9	−1.8 (3)	C24'—C25'—C26'—C21'	0.7 (3)
C6—C7—C8—C9	−179.51 (17)	C22'—C21'—C26'—C25'	−0.5 (3)
C12—C7—C8—N1	175.14 (16)	S1'—C21'—C26'—C25'	178.27 (16)
C6—C7—C8—N1	−2.6 (2)	C24—C25—C26—C21	−0.6 (4)
C12'—C7'—C8'—C9'	2.7 (3)	C22—C21—C26—C25	0.6 (3)
C6'—C7'—C8'—C9'	−178.41 (16)	S1—C21—C26—C25	−178.94 (18)
C12'—C7'—C8'—N1'	−175.16 (15)	C9—C8—N1—C5	179.77 (18)
C6'—C7'—C8'—N1'	3.75 (19)	C7—C8—N1—C5	3.0 (2)
C7—C8—C9—C10	−1.0 (3)	C9—C8—N1—S1	−26.1 (3)


N1—C8—C9—C10	−177.35 (17)	C7—C8—N1—S1	157.12 (13)
C7'—C8'—C9'—C10'	0.2 (3)	C4—C5—N1—C8	179.5 (2)
N1'—C8'—C9'—C10'	177.62 (16)	C6—C5—N1—C8	−2.3 (2)
C8'—C9'—C10'—O3'	178.07 (15)	C4—C5—N1—S1	24.8 (3)
C8'—C9'—C10'—C11'	−1.4 (3)	C6—C5—N1—S1	−156.96 (14)
C8—C9—C10—O3	−179.04 (16)	C9'—C8'—N1'—C5'	178.38 (17)
C8—C9—C10—C11	1.0 (3)	C7'—C8'—N1'—C5'	−3.91 (19)
C9—C10—C11—C12	1.8 (3)	C9'—C8'—N1'—S1'	23.4 (3)
O3—C10—C11—C12	−178.11 (16)	C7'—C8'—N1'—S1'	−158.95 (13)
C9—C10—C11—C13	−178.23 (17)	C4'—C5'—N1'—C8'	−178.32 (19)
O3—C10—C11—C13	1.8 (2)	C6'—C5'—N1'—C8'	2.53 (19)
C9'—C10'—C11'—C12'	−0.4 (3)	C4'—C5'—N1'—S1'	−22.7 (3)
O3'—C10'—C11'—C12'	−179.87 (15)	C6'—C5'—N1'—S1'	158.16 (13)
C9'—C10'—C11'—C13'	178.22 (17)	C9'—C10'—O3'—C14'	−178.07 (17)
O3'—C10'—C11'—C13'	−1.29 (19)	C11'—C10'—O3'—C14'	1.46 (19)
C10'—C11'—C12'—C7'	3.2 (2)	C15'—C14'—O3'—C10'	177.89 (19)
C13'—C11'—C12'—C7'	−174.88 (18)	C13'—C14'—O3'—C10'	−1.0 (2)
C10'—C11'—C12'—C19'	−177.26 (15)	C9—C10—O3—C14	178.21 (17)
C13'—C11'—C12'—C19'	4.7 (3)	C11—C10—O3—C14	−1.9 (2)
C8'—C7'—C12'—C11'	−4.3 (2)	C15—C14—O3—C10	−178.02 (19)
C6'—C7'—C12'—C11'	177.12 (17)	C13—C14—O3—C10	1.1 (2)
C8'—C7'—C12'—C19'	176.17 (15)	C8—N1—S1—O2	163.27 (15)
C6'—C7'—C12'—C19'	−2.4 (3)	C5—N1—S1—O2	−46.32 (17)
C10—C11—C12—C7	−4.5 (3)	C8—N1—S1—O1	34.08 (17)
C13—C11—C12—C7	175.6 (2)	C5—N1—S1—O1	−175.50 (15)
C10—C11—C12—C19	174.46 (17)	C8—N1—S1—C21	−81.44 (16)
C13—C11—C12—C19	−5.5 (3)	C5—N1—S1—C21	68.98 (16)
C8—C7—C12—C11	4.5 (3)	C26—C21—S1—O2	10.56 (19)
C6—C7—C12—C11	−178.43 (19)	C22—C21—S1—O2	−169.02 (15)
C8—C7—C12—C19	−174.43 (17)	C26—C21—S1—O1	142.30 (16)
C6—C7—C12—C19	2.6 (3)	C22—C21—S1—O1	−37.27 (18)
C12'—C11'—C13'—C18'	1.1 (4)	C26—C21—S1—N1	−103.44 (17)
C10'—C11'—C13'—C18'	−177.2 (2)	C22—C21—S1—N1	76.98 (17)
C12'—C11'—C13'—C14'	178.86 (19)	C8'—N1'—S1'—O1'	−162.49 (14)
C10'—C11'—C13'—C14'	0.59 (19)	C5'—N1'—S1'—O1'	46.03 (16)
C10—C11—C13—C14	−1.1 (2)	C8'—N1'—S1'—O2'	−33.47 (17)
C12—C11—C13—C14	178.9 (2)	C5'—N1'—S1'—O2'	175.04 (14)
C10—C11—C13—C18	177.2 (3)	C8'—N1'—S1'—C21'	82.31 (16)
C12—C11—C13—C18	−2.8 (4)	C5'—N1'—S1'—C21'	−69.18 (16)
C18'—C13'—C14'—C15'	−0.4 (3)	C22'—C21'—S1'—O1'	−14.57 (18)
C11'—C13'—C14'—C15'	−178.65 (19)	C26'—C21'—S1'—O1'	166.59 (15)
C18'—C13'—C14'—O3'	178.54 (17)	C22'—C21'—S1'—O2'	−146.37 (15)
C11'—C13'—C14'—O3'	0.3 (2)	C26'—C21'—S1'—O2'	34.79 (17)
C18—C13—C14—C15	0.4 (3)	C22'—C21'—S1'—N1'	99.55 (16)
C11—C13—C14—C15	179.11 (19)	C26'—C21'—S1'—N1'	−79.29 (16)

Hydrogen-bond geometry (Å, °)

Cg1, Cg4, Cg6, Cg17 and Cg20 are the centroids of rings O3/C10/C11/C13/C14, C7–C12, C21–C26, O3'/C10'/C11'/C13'/C14' and C7'–C12', respectively.

$D\text{--H}\cdots A$	$D\text{--H}$	$H\cdots A$	$D\cdots A$	$D\text{--H}\cdots A$
C4—H4···O2	0.93	2.35	2.954 (3)	122
C4'—H4'···O1'	0.93	2.36	2.966 (3)	122
C9—H9···O1	0.93	2.29	2.881 (2)	121
C9'—H9'···O2'	0.93	2.28	2.875 (2)	121
C4—H4···O2 ⁱ	0.93	2.53	3.277 (3)	137
C20'—H20A···Cg17 ⁱⁱ	0.96	2.82	3.449 (3)	124
C20'—H20C···Cg20 ⁱⁱ	0.96	2.79	3.427 (3)	125
C20—H20D···Cg4 ⁱⁱⁱ	0.96	2.83	3.464 (3)	124
C20'—H20C···Cg1 ⁱⁱⁱ	0.96	2.85	3.478 (3)	124
C25'—H25'···Cg6 ^{iv}	0.93	2.90	3.762 (3)	155

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x, -y+1, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x-1, y, z$.

(2) 2-(4,5-Dimethoxy-2-nitrophenyl)-4-hydroxy-9-phenylsulfonyl-9*H*-carbazole-3-carbaldehyde*Crystal data*

$$M_r = 532.51$$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$$a = 11.2133 (3) \text{ \AA}$$

$$b = 14.5811 (4) \text{ \AA}$$

$$c = 15.1509 (4) \text{ \AA}$$

$$\beta = 102.320 (1)^\circ$$

$$V = 2420.16 (11) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 1104$$

$$D_x = 1.461 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6535 reflections

$$\theta = 2.3\text{--}29.4^\circ$$

$$\mu = 0.19 \text{ mm}^{-1}$$

$$T = 296 \text{ K}$$

Block, colourless

$$0.35 \times 0.30 \times 0.25 \text{ mm}$$

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω & φ scans

Absorption correction: multi-scan
(SADABS; Bruker, 2008)

$$T_{\min} = 0.935, T_{\max} = 0.953$$

42635 measured reflections

6535 independent reflections

4517 reflections with $I > 2\sigma(I)$

$$R_{\text{int}} = 0.025$$

$$\theta_{\max} = 29.4^\circ, \theta_{\min} = 2.3^\circ$$

$$h = -15 \rightarrow 13$$

$$k = -19 \rightarrow 20$$

$$l = -20 \rightarrow 20$$

Refinement

Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.047$$

$$wR(F^2) = 0.150$$

$$S = 1.00$$

6535 reflections

345 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.068P)^2 + 1.0361P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.33 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.46 \text{ e \AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6804 (3)	0.56057 (17)	0.45636 (16)	0.0713 (6)
H1	0.6535	0.6007	0.4957	0.086*
C2	0.8008 (3)	0.5346 (2)	0.4706 (2)	0.0874 (9)
H2	0.8559	0.5577	0.5204	0.105*
C3	0.8422 (3)	0.4750 (2)	0.4128 (2)	0.0891 (8)
H3	0.9245	0.4595	0.4242	0.107*
C4	0.7646 (2)	0.43809 (19)	0.33878 (18)	0.0713 (6)
H4	0.7923	0.3977	0.3001	0.086*
C5	0.64295 (18)	0.46384 (15)	0.32434 (13)	0.0534 (5)
C6	0.5994 (2)	0.52489 (14)	0.38090 (13)	0.0549 (5)
C7	0.47098 (19)	0.53781 (13)	0.34519 (13)	0.0494 (4)
C8	0.43752 (17)	0.48471 (12)	0.26750 (12)	0.0447 (4)
C9	0.31852 (17)	0.47932 (12)	0.21721 (12)	0.0456 (4)
H9	0.2992	0.4433	0.1655	0.055*
C10	0.23043 (18)	0.52915 (12)	0.24661 (13)	0.0478 (4)
C11	0.2616 (2)	0.58639 (13)	0.32419 (14)	0.0543 (5)
C12	0.3826 (2)	0.59004 (13)	0.37285 (13)	0.0557 (5)
C13	0.1699 (3)	0.63906 (17)	0.35558 (17)	0.0733 (7)
H13	0.0893	0.6345	0.3240	0.088*
C14	0.45277 (18)	0.27073 (13)	0.25225 (14)	0.0514 (4)
C15	0.3416 (2)	0.23769 (17)	0.2092 (2)	0.0797 (7)
H15	0.3097	0.2509	0.1487	0.096*
C16	0.2785 (3)	0.1842 (2)	0.2586 (4)	0.1212 (15)
H16	0.2032	0.1596	0.2308	0.145*
C17	0.3251 (5)	0.1667 (2)	0.3478 (4)	0.1320 (19)
H17	0.2809	0.1311	0.3804	0.158*
C18	0.4361 (4)	0.2011 (3)	0.3895 (3)	0.1163 (13)
H18	0.4667	0.1892	0.4505	0.140*
C19	0.5023 (3)	0.2529 (2)	0.34196 (18)	0.0833 (8)
H19	0.5788	0.2756	0.3694	0.100*
C20	0.10327 (17)	0.52421 (13)	0.19144 (13)	0.0485 (4)
C21	0.06126 (19)	0.59499 (13)	0.13086 (14)	0.0535 (5)
H21	0.1094	0.6470	0.1314	0.064*
C22	-0.04914 (18)	0.59032 (14)	0.07036 (14)	0.0526 (5)
C23	-0.12246 (18)	0.51187 (15)	0.06833 (13)	0.0523 (5)

C24	-0.08426 (19)	0.44225 (14)	0.12820 (14)	0.0546 (5)
H24	-0.1326	0.3905	0.1283	0.066*
C25	0.02706 (18)	0.44943 (13)	0.18874 (13)	0.0504 (4)
C26	-0.0294 (3)	0.73900 (18)	0.0125 (2)	0.0880 (8)
H26A	-0.0178	0.7652	0.0718	0.132*
H26B	-0.0744	0.7811	-0.0309	0.132*
H26C	0.0486	0.7271	-0.0017	0.132*
C27	-0.3111 (3)	0.4409 (2)	0.0050 (2)	0.0998 (11)
H27A	-0.2738	0.3845	-0.0074	0.150*
H27B	-0.3831	0.4518	-0.0410	0.150*
H27C	-0.3328	0.4370	0.0628	0.150*
N1	0.54318 (14)	0.43996 (11)	0.25235 (11)	0.0480 (4)
N2	0.05889 (18)	0.37377 (13)	0.25278 (15)	0.0636 (5)
O1	0.65435 (13)	0.31079 (11)	0.20126 (11)	0.0646 (4)
O2	0.46032 (15)	0.36250 (12)	0.10740 (9)	0.0624 (4)
O3	0.41687 (18)	0.64230 (12)	0.44680 (11)	0.0771 (5)
H3A	0.3575	0.6700	0.4567	0.116*
O4	0.1928 (2)	0.68940 (14)	0.42164 (13)	0.0960 (6)
O5	0.1327 (2)	0.38570 (13)	0.32219 (12)	0.0859 (6)
O6	0.0090 (2)	0.30090 (13)	0.23384 (19)	0.1202 (9)
O7	-0.09553 (15)	0.65540 (11)	0.00978 (11)	0.0679 (4)
O8	-0.22814 (14)	0.51372 (12)	0.00565 (11)	0.0660 (4)
S1	0.53315 (4)	0.34260 (3)	0.19364 (3)	0.04694 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0855 (17)	0.0674 (14)	0.0553 (12)	-0.0180 (12)	0.0023 (11)	-0.0065 (10)
C2	0.0765 (18)	0.095 (2)	0.0761 (17)	-0.0219 (15)	-0.0172 (14)	-0.0047 (15)
C3	0.0605 (15)	0.107 (2)	0.0890 (19)	-0.0087 (15)	-0.0084 (14)	-0.0067 (17)
C4	0.0517 (12)	0.0842 (17)	0.0736 (15)	-0.0044 (11)	0.0037 (11)	-0.0054 (13)
C5	0.0510 (11)	0.0575 (11)	0.0500 (10)	-0.0113 (9)	0.0067 (8)	0.0014 (9)
C6	0.0637 (12)	0.0503 (10)	0.0491 (10)	-0.0145 (9)	0.0081 (9)	0.0002 (8)
C7	0.0634 (12)	0.0407 (9)	0.0457 (9)	-0.0081 (8)	0.0151 (8)	-0.0036 (7)
C8	0.0517 (10)	0.0404 (9)	0.0445 (9)	-0.0055 (7)	0.0157 (8)	-0.0031 (7)
C9	0.0505 (10)	0.0422 (9)	0.0458 (9)	-0.0029 (7)	0.0138 (8)	-0.0055 (7)
C10	0.0563 (11)	0.0393 (9)	0.0520 (10)	-0.0005 (8)	0.0206 (8)	-0.0002 (7)
C11	0.0712 (13)	0.0420 (9)	0.0557 (11)	0.0045 (9)	0.0268 (10)	-0.0036 (8)
C12	0.0802 (14)	0.0406 (9)	0.0481 (10)	-0.0052 (9)	0.0176 (10)	-0.0067 (8)
C13	0.0913 (18)	0.0641 (14)	0.0723 (15)	0.0170 (12)	0.0346 (13)	-0.0068 (11)
C14	0.0555 (11)	0.0454 (9)	0.0568 (11)	0.0013 (8)	0.0201 (9)	-0.0033 (8)
C15	0.0688 (15)	0.0590 (14)	0.110 (2)	-0.0134 (12)	0.0169 (14)	-0.0089 (14)
C16	0.081 (2)	0.074 (2)	0.220 (5)	-0.0199 (16)	0.056 (3)	0.011 (3)
C17	0.145 (4)	0.072 (2)	0.218 (5)	0.004 (2)	0.127 (4)	0.038 (3)
C18	0.171 (4)	0.100 (2)	0.098 (2)	0.005 (3)	0.073 (3)	0.0353 (19)
C19	0.107 (2)	0.0841 (18)	0.0604 (14)	-0.0114 (16)	0.0204 (14)	0.0180 (13)
C20	0.0506 (10)	0.0433 (9)	0.0573 (11)	0.0046 (8)	0.0243 (8)	-0.0006 (8)
C21	0.0561 (11)	0.0452 (10)	0.0658 (12)	0.0020 (8)	0.0279 (10)	0.0046 (9)

C22	0.0547 (11)	0.0524 (10)	0.0589 (11)	0.0114 (9)	0.0301 (9)	0.0077 (9)
C23	0.0489 (10)	0.0616 (12)	0.0526 (10)	0.0066 (9)	0.0247 (8)	0.0034 (9)
C24	0.0531 (11)	0.0535 (11)	0.0623 (12)	-0.0032 (9)	0.0236 (9)	0.0028 (9)
C25	0.0547 (11)	0.0444 (10)	0.0564 (11)	0.0037 (8)	0.0217 (9)	0.0037 (8)
C26	0.108 (2)	0.0609 (14)	0.097 (2)	0.0041 (14)	0.0266 (17)	0.0282 (14)
C27	0.0640 (16)	0.125 (3)	0.105 (2)	-0.0266 (16)	0.0042 (15)	0.041 (2)
N1	0.0430 (8)	0.0530 (9)	0.0485 (8)	-0.0045 (7)	0.0110 (6)	-0.0055 (7)
N2	0.0637 (11)	0.0504 (10)	0.0770 (13)	-0.0043 (8)	0.0156 (10)	0.0102 (9)
O1	0.0527 (8)	0.0739 (10)	0.0719 (10)	0.0082 (7)	0.0235 (7)	-0.0076 (8)
O2	0.0753 (10)	0.0752 (10)	0.0372 (7)	0.0096 (8)	0.0127 (6)	-0.0036 (6)
O3	0.1079 (14)	0.0637 (10)	0.0587 (9)	0.0009 (9)	0.0157 (9)	-0.0251 (7)
O4	0.1246 (17)	0.0882 (13)	0.0835 (12)	0.0285 (12)	0.0407 (12)	-0.0271 (10)
O5	0.1136 (15)	0.0743 (11)	0.0630 (10)	-0.0168 (10)	0.0036 (10)	0.0174 (9)
O6	0.1217 (17)	0.0587 (11)	0.152 (2)	-0.0302 (11)	-0.0351 (15)	0.0344 (12)
O7	0.0683 (10)	0.0655 (9)	0.0742 (10)	0.0102 (7)	0.0246 (8)	0.0226 (8)
O8	0.0533 (8)	0.0826 (11)	0.0638 (9)	-0.0019 (8)	0.0168 (7)	0.0140 (8)
S1	0.0484 (3)	0.0542 (3)	0.0409 (2)	0.00208 (19)	0.01574 (18)	-0.00321 (18)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.374 (4)	C16—H16	0.9300
C1—C6	1.400 (3)	C17—C18	1.365 (6)
C1—H1	0.9300	C17—H17	0.9300
C2—C3	1.382 (4)	C18—C19	1.368 (4)
C2—H2	0.9300	C18—H18	0.9300
C3—C4	1.374 (4)	C19—H19	0.9300
C3—H3	0.9300	C20—C25	1.381 (3)
C4—C5	1.386 (3)	C20—C21	1.395 (3)
C4—H4	0.9300	C21—C22	1.376 (3)
C5—C6	1.394 (3)	C21—H21	0.9300
C5—N1	1.429 (2)	C22—O7	1.345 (2)
C6—C7	1.439 (3)	C22—C23	1.405 (3)
C7—C12	1.384 (3)	C23—O8	1.352 (3)
C7—C8	1.392 (3)	C23—C24	1.368 (3)
C8—C9	1.390 (3)	C24—C25	1.386 (3)
C8—N1	1.414 (2)	C24—H24	0.9300
C9—C10	1.375 (3)	C25—N2	1.462 (3)
C9—H9	0.9300	C26—O7	1.423 (3)
C10—C11	1.423 (3)	C26—H26A	0.9600
C10—C20	1.493 (3)	C26—H26B	0.9600
C11—C12	1.400 (3)	C26—H26C	0.9600
C11—C13	1.443 (3)	C27—O8	1.410 (3)
C12—O3	1.342 (2)	C27—H27A	0.9600
C13—O4	1.224 (3)	C27—H27B	0.9600
C13—H13	0.9300	C27—H27C	0.9600
C14—C15	1.365 (3)	N1—S1	1.6663 (16)
C14—C19	1.379 (3)	N2—O5	1.204 (2)
C14—S1	1.744 (2)	N2—O6	1.207 (3)

C15—C16	1.377 (5)	O1—S1	1.4172 (15)
C15—H15	0.9300	O2—S1	1.4154 (14)
C16—C17	1.365 (6)	O3—H3A	0.8200
C2—C1—C6	118.0 (3)	C17—C18—C19	120.2 (4)
C2—C1—H1	121.0	C17—C18—H18	119.9
C6—C1—H1	121.0	C19—C18—H18	119.9
C1—C2—C3	121.7 (2)	C18—C19—C14	118.3 (3)
C1—C2—H2	119.1	C18—C19—H19	120.9
C3—C2—H2	119.1	C14—C19—H19	120.9
C4—C3—C2	121.6 (3)	C25—C20—C21	116.42 (19)
C4—C3—H3	119.2	C25—C20—C10	124.47 (17)
C2—C3—H3	119.2	C21—C20—C10	118.75 (17)
C3—C4—C5	117.0 (3)	C22—C21—C20	122.10 (19)
C3—C4—H4	121.5	C22—C21—H21	118.9
C5—C4—H4	121.5	C20—C21—H21	118.9
C4—C5—C6	122.5 (2)	O7—C22—C21	125.56 (19)
C4—C5—N1	129.7 (2)	O7—C22—C23	114.83 (19)
C6—C5—N1	107.75 (18)	C21—C22—C23	119.60 (18)
C5—C6—C1	119.2 (2)	O8—C23—C24	125.75 (19)
C5—C6—C7	107.83 (17)	O8—C23—C22	114.86 (18)
C1—C6—C7	133.0 (2)	C24—C23—C22	119.4 (2)
C12—C7—C8	118.74 (19)	C23—C24—C25	119.54 (19)
C12—C7—C6	133.03 (18)	C23—C24—H24	120.2
C8—C7—C6	108.22 (17)	C25—C24—H24	120.2
C9—C8—C7	123.28 (17)	C20—C25—C24	122.93 (18)
C9—C8—N1	128.65 (16)	C20—C25—N2	120.95 (19)
C7—C8—N1	108.07 (16)	C24—C25—N2	116.09 (18)
C10—C9—C8	117.68 (17)	O7—C26—H26A	109.5
C10—C9—H9	121.2	O7—C26—H26B	109.5
C8—C9—H9	121.2	H26A—C26—H26B	109.5
C9—C10—C11	120.71 (19)	O7—C26—H26C	109.5
C9—C10—C20	117.36 (17)	H26A—C26—H26C	109.5
C11—C10—C20	121.86 (17)	H26B—C26—H26C	109.5
C12—C11—C10	119.81 (18)	O8—C27—H27A	109.5
C12—C11—C13	119.0 (2)	O8—C27—H27B	109.5
C10—C11—C13	121.2 (2)	H27A—C27—H27B	109.5
O3—C12—C7	118.0 (2)	O8—C27—H27C	109.5
O3—C12—C11	122.32 (19)	H27A—C27—H27C	109.5
C7—C12—C11	119.72 (17)	H27B—C27—H27C	109.5
O4—C13—C11	123.3 (3)	C8—N1—C5	108.08 (15)
O4—C13—H13	118.4	C8—N1—S1	121.10 (12)
C11—C13—H13	118.4	C5—N1—S1	124.59 (14)
C15—C14—C19	122.6 (2)	O5—N2—O6	122.2 (2)
C15—C14—S1	119.18 (19)	O5—N2—C25	119.70 (18)
C19—C14—S1	118.15 (18)	O6—N2—C25	118.1 (2)
C14—C15—C16	117.6 (3)	C12—O3—H3A	109.5
C14—C15—H15	121.2	C22—O7—C26	117.48 (19)

C16—C15—H15	121.2	C23—O8—C27	117.93 (19)
C17—C16—C15	120.7 (4)	O2—S1—O1	119.78 (9)
C17—C16—H16	119.7	O2—S1—N1	106.14 (9)
C15—C16—H16	119.7	O1—S1—N1	106.32 (9)
C18—C17—C16	120.6 (3)	O2—S1—C14	109.52 (10)
C18—C17—H17	119.7	O1—S1—C14	110.66 (10)
C16—C17—H17	119.7	N1—S1—C14	102.92 (9)
C6—C1—C2—C3	0.0 (4)	C9—C10—C20—C21	-99.0 (2)
C1—C2—C3—C4	-0.7 (5)	C11—C10—C20—C21	78.0 (2)
C2—C3—C4—C5	0.5 (4)	C25—C20—C21—C22	-1.1 (3)
C3—C4—C5—C6	0.5 (4)	C10—C20—C21—C22	172.28 (17)
C3—C4—C5—N1	177.3 (2)	C20—C21—C22—O7	-179.71 (17)
C4—C5—C6—C1	-1.3 (3)	C20—C21—C22—C23	-0.5 (3)
N1—C5—C6—C1	-178.70 (18)	O7—C22—C23—O8	0.2 (2)
C4—C5—C6—C7	178.6 (2)	C21—C22—C23—O8	-179.13 (17)
N1—C5—C6—C7	1.2 (2)	O7—C22—C23—C24	-179.03 (17)
C2—C1—C6—C5	1.0 (3)	C21—C22—C23—C24	1.7 (3)
C2—C1—C6—C7	-178.9 (2)	O8—C23—C24—C25	179.69 (18)
C5—C6—C7—C12	-179.9 (2)	C22—C23—C24—C25	-1.2 (3)
C1—C6—C7—C12	0.0 (4)	C21—C20—C25—C24	1.6 (3)
C5—C6—C7—C8	0.0 (2)	C10—C20—C25—C24	-171.36 (18)
C1—C6—C7—C8	179.9 (2)	C21—C20—C25—N2	-176.43 (17)
C12—C7—C8—C9	-1.9 (3)	C10—C20—C25—N2	10.6 (3)
C6—C7—C8—C9	178.26 (17)	C23—C24—C25—C20	-0.5 (3)
C12—C7—C8—N1	178.64 (16)	C23—C24—C25—N2	177.65 (18)
C6—C7—C8—N1	-1.3 (2)	C9—C8—N1—C5	-177.46 (18)
C7—C8—C9—C10	-0.1 (3)	C7—C8—N1—C5	2.0 (2)
N1—C8—C9—C10	179.28 (17)	C9—C8—N1—S1	-24.0 (3)
C8—C9—C10—C11	2.0 (3)	C7—C8—N1—S1	155.45 (13)
C8—C9—C10—C20	179.03 (16)	C4—C5—N1—C8	-179.2 (2)
C9—C10—C11—C12	-1.9 (3)	C6—C5—N1—C8	-2.0 (2)
C20—C10—C11—C12	-178.83 (18)	C4—C5—N1—S1	28.5 (3)
C9—C10—C11—C13	179.94 (19)	C6—C5—N1—S1	-154.29 (14)
C20—C10—C11—C13	3.1 (3)	C20—C25—N2—O5	19.9 (3)
C8—C7—C12—O3	-178.63 (18)	C24—C25—N2—O5	-158.3 (2)
C6—C7—C12—O3	1.2 (3)	C20—C25—N2—O6	-160.5 (2)
C8—C7—C12—C11	1.9 (3)	C24—C25—N2—O6	21.3 (3)
C6—C7—C12—C11	-178.2 (2)	C21—C22—O7—C26	-4.8 (3)
C10—C11—C12—O3	-179.54 (18)	C23—C22—O7—C26	176.0 (2)
C13—C11—C12—O3	-1.4 (3)	C24—C23—O8—C27	4.6 (3)
C10—C11—C12—C7	-0.1 (3)	C22—C23—O8—C27	-174.6 (2)
C13—C11—C12—C7	178.08 (19)	C8—N1—S1—O2	58.73 (16)
C12—C11—C13—O4	2.2 (4)	C5—N1—S1—O2	-152.36 (15)
C10—C11—C13—O4	-179.6 (2)	C8—N1—S1—O1	-172.69 (14)
C19—C14—C15—C16	-0.4 (4)	C5—N1—S1—O1	-23.78 (18)
S1—C14—C15—C16	-177.4 (2)	C8—N1—S1—C14	-56.31 (16)
C14—C15—C16—C17	1.3 (5)	C5—N1—S1—C14	92.60 (16)

C15—C16—C17—C18	−0.8 (6)	C15—C14—S1—O2	5.3 (2)
C16—C17—C18—C19	−0.6 (6)	C19—C14—S1—O2	−171.91 (19)
C17—C18—C19—C14	1.4 (5)	C15—C14—S1—O1	−128.93 (19)
C15—C14—C19—C18	−0.9 (4)	C19—C14—S1—O1	53.9 (2)
S1—C14—C19—C18	176.1 (2)	C15—C14—S1—N1	117.83 (18)
C9—C10—C20—C25	73.8 (2)	C19—C14—S1—N1	−59.3 (2)
C11—C10—C20—C25	−109.2 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3A···O4	0.82	1.83	2.554 (3)	146
C4—H4···O1	0.93	2.29	2.866 (3)	119
C9—H9···O2	0.93	2.47	3.054 (2)	121
C2—H2···O5 ⁱ	0.93	2.50	3.281 (3)	142
C17—H17···O8 ⁱⁱ	0.93	2.59	3.481 (5)	161
C18—H18···O2 ⁱⁱⁱ	0.93	2.51	3.384 (5)	157
C26—H26C···O4 ^{iv}	0.96	2.50	3.265 (4)	137

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, y-1/2, -z+1/2$; (iii) $x, -y+1/2, z+1/2$; (iv) $x, -y+3/2, z-1/2$.

(3) 12-Phenyl-7-phenylsulfonyl-7*H*-benzofuro[2,3-*b*]carbazole*Crystal data*

$C_{30}H_{19}NO_3S$	$F(000) = 984$
$M_r = 473.52$	$D_x = 1.361 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 5055 reflections
$a = 10.6461 (10) \text{ \AA}$	$\theta = 2.2\text{--}27.1^\circ$
$b = 11.8994 (11) \text{ \AA}$	$\mu = 0.17 \text{ mm}^{-1}$
$c = 18.2418 (16) \text{ \AA}$	$T = 296 \text{ K}$
$V = 2310.9 (4) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.35 \times 0.30 \times 0.25 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD	29269 measured reflections
diffractometer	5055 independent reflections
Radiation source: fine-focus sealed tube	3194 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.061$
ω & φ scans	$\theta_{\max} = 27.1^\circ, \theta_{\min} = 2.2^\circ$
Absorption correction: multi-scan	$h = -13 \rightarrow 13$
(SADABS; Bruker, 2008)	$k = -15 \rightarrow 15$
$T_{\min} = 0.941, T_{\max} = 0.957$	$l = -23 \rightarrow 22$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.101$	H-atom parameters constrained
$S = 1.00$	
5055 reflections	
316 parameters	
1 restraint	

$$w = 1/[\sigma^2(F_o^2) + (0.0488P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack (1983), 2189 Friedel pairs

Absolute structure parameter: 0.08 (9)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.0093 (3)	0.6577 (2)	0.59295 (16)	0.0546 (7)
H1	-0.0202	0.7107	0.6298	0.066*
C2	-0.1114 (3)	0.6067 (3)	0.56158 (18)	0.0668 (9)
H2	-0.1917	0.6246	0.5779	0.080*
C3	-0.0964 (3)	0.5288 (3)	0.50594 (18)	0.0697 (9)
H3	-0.1672	0.4959	0.4852	0.084*
C4	0.0190 (3)	0.4993 (2)	0.48087 (16)	0.0628 (8)
H4	0.0282	0.4472	0.4433	0.075*
C5	0.1227 (3)	0.5493 (2)	0.51323 (14)	0.0485 (7)
C6	0.1098 (3)	0.6296 (2)	0.56937 (14)	0.0419 (6)
C7	0.2349 (2)	0.6687 (2)	0.58813 (13)	0.0398 (6)
C8	0.3212 (3)	0.6114 (2)	0.54335 (12)	0.0427 (6)
C9	0.4492 (3)	0.6300 (2)	0.54535 (14)	0.0510 (7)
H9	0.5054	0.5918	0.5153	0.061*
C10	0.4868 (3)	0.7095 (2)	0.59553 (14)	0.0455 (7)
C11	0.4059 (3)	0.7687 (2)	0.64193 (13)	0.0413 (6)
C12	0.2773 (3)	0.7488 (2)	0.63911 (12)	0.0404 (6)
C13	0.4852 (3)	0.8433 (2)	0.68342 (14)	0.0452 (7)
C14	0.6070 (3)	0.8217 (2)	0.66050 (15)	0.0513 (7)
C15	0.7122 (3)	0.8755 (3)	0.68680 (18)	0.0633 (9)
H15	0.7925	0.8582	0.6702	0.076*
C16	0.6910 (3)	0.9561 (3)	0.73888 (18)	0.0681 (9)
H16	0.7591	0.9949	0.7584	0.082*
C17	0.5714 (3)	0.9816 (3)	0.76328 (16)	0.0626 (8)
H17	0.5607	1.0376	0.7983	0.075*
C18	0.4674 (3)	0.9256 (2)	0.73661 (14)	0.0554 (7)
H18	0.3873	0.9425	0.7538	0.067*
C19	0.1912 (2)	0.8073 (2)	0.69074 (13)	0.0405 (6)
C20	0.1210 (3)	0.9003 (2)	0.66997 (15)	0.0507 (7)
H20	0.1255	0.9270	0.6221	0.061*
C21	0.0447 (3)	0.9532 (2)	0.72040 (17)	0.0592 (8)

H21	-0.0021	1.0153	0.7058	0.071*
C22	0.0362 (3)	0.9174 (3)	0.78952 (15)	0.0619 (8)
H22	-0.0153	0.9546	0.8227	0.074*
C23	0.1041 (3)	0.8252 (3)	0.81136 (16)	0.0708 (9)
H23	0.0978	0.7995	0.8594	0.085*
C24	0.1812 (3)	0.7709 (2)	0.76269 (13)	0.0562 (8)
H24	0.2274	0.7089	0.7781	0.067*
C25	0.3170 (3)	0.3277 (2)	0.54143 (14)	0.0493 (7)
C26	0.4280 (3)	0.3073 (3)	0.57863 (18)	0.0666 (9)
H26	0.5024	0.3417	0.5641	0.080*
C27	0.4265 (4)	0.2342 (3)	0.6382 (2)	0.0808 (11)
H27	0.5000	0.2199	0.6641	0.097*
C28	0.3177 (5)	0.1840 (3)	0.6584 (2)	0.0906 (12)
H28	0.3176	0.1342	0.6977	0.109*
C29	0.2091 (4)	0.2050 (3)	0.6224 (2)	0.0904 (12)
H29	0.1352	0.1707	0.6377	0.108*
C30	0.2075 (3)	0.2761 (3)	0.56366 (18)	0.0686 (9)
H30	0.1327	0.2897	0.5388	0.082*
N1	0.2531 (2)	0.53725 (18)	0.49596 (11)	0.0497 (6)
O1	0.2250 (2)	0.37314 (17)	0.41398 (10)	0.0763 (7)
O2	0.4380 (2)	0.43983 (18)	0.44385 (11)	0.0795 (7)
O3	0.60967 (18)	0.73994 (16)	0.60631 (10)	0.0573 (5)
S1	0.31294 (8)	0.41674 (6)	0.46577 (4)	0.0583 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0488 (19)	0.0502 (17)	0.0649 (19)	0.0022 (15)	-0.0009 (16)	0.0002 (15)
C2	0.0497 (19)	0.0610 (19)	0.090 (2)	0.0060 (17)	-0.0077 (18)	0.0020 (18)
C3	0.060 (2)	0.0595 (19)	0.089 (2)	-0.0023 (17)	-0.0197 (18)	-0.0047 (19)
C4	0.079 (2)	0.0473 (17)	0.0623 (19)	0.0004 (17)	-0.0142 (18)	-0.0081 (15)
C5	0.0597 (19)	0.0395 (15)	0.0463 (15)	0.0038 (15)	-0.0043 (14)	0.0030 (13)
C6	0.0483 (17)	0.0370 (13)	0.0404 (14)	0.0004 (13)	-0.0020 (13)	0.0078 (12)
C7	0.0504 (17)	0.0337 (14)	0.0353 (14)	0.0036 (12)	0.0024 (12)	0.0028 (12)
C8	0.0583 (18)	0.0354 (14)	0.0345 (13)	0.0007 (14)	0.0085 (14)	0.0028 (11)
C9	0.060 (2)	0.0431 (15)	0.0499 (17)	0.0039 (14)	0.0170 (15)	-0.0010 (14)
C10	0.0444 (18)	0.0460 (16)	0.0462 (15)	0.0007 (13)	0.0047 (14)	0.0045 (13)
C11	0.0469 (18)	0.0361 (13)	0.0409 (14)	0.0024 (13)	0.0024 (13)	0.0047 (12)
C12	0.0542 (18)	0.0341 (13)	0.0327 (13)	0.0069 (13)	0.0028 (12)	0.0026 (11)
C13	0.0516 (19)	0.0437 (16)	0.0404 (15)	0.0000 (14)	0.0002 (13)	0.0060 (13)
C14	0.057 (2)	0.0433 (16)	0.0539 (17)	-0.0014 (16)	-0.0010 (16)	0.0037 (14)
C15	0.053 (2)	0.060 (2)	0.076 (2)	-0.0077 (17)	-0.0024 (17)	0.0049 (18)
C16	0.069 (2)	0.057 (2)	0.079 (2)	-0.0150 (19)	-0.0148 (19)	0.0028 (18)
C17	0.080 (2)	0.0466 (17)	0.0614 (18)	-0.0068 (17)	-0.0117 (18)	-0.0040 (15)
C18	0.063 (2)	0.0449 (16)	0.0587 (17)	0.0014 (15)	-0.0005 (15)	0.0022 (15)
C19	0.0445 (16)	0.0357 (13)	0.0413 (14)	0.0019 (13)	-0.0014 (13)	-0.0010 (11)
C20	0.0557 (17)	0.0481 (16)	0.0482 (15)	0.0028 (16)	0.0013 (14)	0.0050 (14)
C21	0.0557 (18)	0.0439 (17)	0.078 (2)	0.0151 (15)	0.0026 (17)	-0.0054 (16)

C22	0.070 (2)	0.0649 (19)	0.0505 (17)	0.0154 (18)	0.0111 (15)	-0.0121 (16)
C23	0.085 (2)	0.083 (2)	0.0438 (16)	0.021 (2)	0.0115 (17)	0.0046 (16)
C24	0.068 (2)	0.0584 (18)	0.0420 (15)	0.0161 (17)	0.0077 (15)	0.0045 (13)
C25	0.0552 (18)	0.0403 (14)	0.0523 (16)	0.0042 (14)	0.0050 (16)	-0.0114 (13)
C26	0.060 (2)	0.065 (2)	0.074 (2)	0.0094 (16)	-0.0077 (18)	-0.0208 (19)
C27	0.102 (3)	0.067 (2)	0.073 (2)	0.034 (2)	-0.034 (2)	-0.019 (2)
C28	0.131 (4)	0.060 (2)	0.081 (2)	0.022 (3)	0.004 (3)	0.007 (2)
C29	0.097 (3)	0.066 (2)	0.108 (3)	0.005 (2)	0.012 (3)	0.027 (2)
C30	0.068 (2)	0.0540 (18)	0.084 (2)	0.0061 (17)	0.0014 (18)	0.0101 (17)
N1	0.0652 (16)	0.0427 (13)	0.0412 (12)	0.0012 (12)	0.0038 (11)	-0.0069 (10)
O1	0.1128 (18)	0.0644 (14)	0.0517 (12)	0.0049 (12)	-0.0123 (12)	-0.0223 (11)
O2	0.0867 (16)	0.0718 (15)	0.0801 (14)	-0.0049 (12)	0.0470 (12)	-0.0182 (12)
O3	0.0487 (12)	0.0556 (12)	0.0675 (12)	-0.0014 (11)	0.0091 (10)	-0.0024 (11)
S1	0.0794 (6)	0.0499 (4)	0.0455 (4)	0.0012 (4)	0.0136 (4)	-0.0115 (4)

Geometric parameters (\AA , $^{\circ}$)

C1—C2	1.370 (4)	C16—H16	0.9300
C1—C6	1.379 (4)	C17—C18	1.381 (4)
C1—H1	0.9300	C17—H17	0.9300
C2—C3	1.383 (4)	C18—H18	0.9300
C2—H2	0.9300	C19—C24	1.386 (3)
C3—C4	1.358 (4)	C19—C20	1.388 (3)
C3—H3	0.9300	C20—C21	1.379 (4)
C4—C5	1.386 (4)	C20—H20	0.9300
C4—H4	0.9300	C21—C22	1.334 (4)
C5—C6	1.407 (4)	C21—H21	0.9300
C5—N1	1.431 (4)	C22—C23	1.373 (4)
C6—C7	1.452 (4)	C22—H22	0.9300
C7—C8	1.406 (3)	C23—C24	1.371 (4)
C7—C12	1.406 (3)	C23—H23	0.9300
C8—C9	1.381 (4)	C24—H24	0.9300
C8—N1	1.432 (3)	C25—C30	1.379 (4)
C9—C10	1.376 (4)	C25—C26	1.385 (4)
C9—H9	0.9300	C25—S1	1.740 (3)
C10—O3	1.372 (3)	C26—C27	1.392 (5)
C10—C11	1.398 (3)	C26—H26	0.9300
C11—C12	1.391 (4)	C27—C28	1.354 (5)
C11—C13	1.440 (4)	C27—H27	0.9300
C12—C19	1.487 (3)	C28—C29	1.352 (5)
C13—C14	1.386 (4)	C28—H28	0.9300
C13—C18	1.392 (4)	C29—C30	1.365 (4)
C14—C15	1.376 (4)	C29—H29	0.9300
C14—O3	1.387 (3)	C30—H30	0.9300
C15—C16	1.369 (4)	N1—S1	1.663 (2)
C15—H15	0.9300	O1—S1	1.427 (2)
C16—C17	1.383 (4)	O2—S1	1.417 (2)

C2—C1—C6	119.5 (3)	C16—C17—H17	119.4
C2—C1—H1	120.3	C17—C18—C13	118.4 (3)
C6—C1—H1	120.3	C17—C18—H18	120.8
C1—C2—C3	120.7 (3)	C13—C18—H18	120.8
C1—C2—H2	119.6	C24—C19—C20	117.8 (2)
C3—C2—H2	119.6	C24—C19—C12	120.0 (2)
C4—C3—C2	121.7 (3)	C20—C19—C12	122.1 (2)
C4—C3—H3	119.1	C21—C20—C19	119.9 (2)
C2—C3—H3	119.1	C21—C20—H20	120.0
C3—C4—C5	117.7 (3)	C19—C20—H20	120.0
C3—C4—H4	121.1	C22—C21—C20	121.6 (3)
C5—C4—H4	121.1	C22—C21—H21	119.2
C4—C5—C6	121.6 (3)	C20—C21—H21	119.2
C4—C5—N1	129.5 (3)	C21—C22—C23	119.6 (3)
C6—C5—N1	108.8 (2)	C21—C22—H22	120.2
C1—C6—C5	118.7 (3)	C23—C22—H22	120.2
C1—C6—C7	133.8 (2)	C24—C23—C22	120.3 (3)
C5—C6—C7	107.4 (2)	C24—C23—H23	119.9
C8—C7—C12	120.3 (2)	C22—C23—H23	119.9
C8—C7—C6	107.9 (2)	C23—C24—C19	120.8 (3)
C12—C7—C6	131.9 (2)	C23—C24—H24	119.6
C9—C8—C7	123.5 (2)	C19—C24—H24	119.6
C9—C8—N1	127.9 (2)	C30—C25—C26	120.0 (3)
C7—C8—N1	108.6 (2)	C30—C25—S1	118.9 (2)
C10—C9—C8	114.5 (2)	C26—C25—S1	121.1 (2)
C10—C9—H9	122.8	C25—C26—C27	118.8 (3)
C8—C9—H9	122.8	C25—C26—H26	120.6
O3—C10—C9	123.7 (2)	C27—C26—H26	120.6
O3—C10—C11	111.6 (2)	C28—C27—C26	119.9 (3)
C9—C10—C11	124.8 (3)	C28—C27—H27	120.1
C12—C11—C10	119.9 (2)	C26—C27—H27	120.1
C12—C11—C13	134.6 (2)	C29—C28—C27	121.2 (4)
C10—C11—C13	105.5 (2)	C29—C28—H28	119.4
C11—C12—C7	117.1 (2)	C27—C28—H28	119.4
C11—C12—C19	120.3 (2)	C28—C29—C30	120.4 (4)
C7—C12—C19	122.6 (2)	C28—C29—H29	119.8
C14—C13—C18	117.9 (3)	C30—C29—H29	119.8
C14—C13—C11	106.0 (2)	C29—C30—C25	119.7 (3)
C18—C13—C11	136.1 (3)	C29—C30—H30	120.1
C15—C14—C13	124.7 (3)	C25—C30—H30	120.1
C15—C14—O3	123.9 (3)	C5—N1—C8	107.3 (2)
C13—C14—O3	111.3 (3)	C5—N1—S1	122.09 (19)
C16—C15—C14	115.7 (3)	C8—N1—S1	122.47 (18)
C16—C15—H15	122.1	C10—O3—C14	105.5 (2)
C14—C15—H15	122.1	O2—S1—O1	119.98 (13)
C15—C16—C17	121.9 (3)	O2—S1—N1	106.63 (13)
C15—C16—H16	119.0	O1—S1—N1	106.34 (13)
C17—C16—H16	119.0	O2—S1—C25	108.57 (14)

C18—C17—C16	121.3 (3)	O1—S1—C25	108.66 (13)
C18—C17—H17	119.4	N1—S1—C25	105.76 (11)
C6—C1—C2—C3	-0.9 (4)	C16—C17—C18—C13	-0.9 (4)
C1—C2—C3—C4	0.6 (5)	C14—C13—C18—C17	0.4 (4)
C2—C3—C4—C5	0.4 (4)	C11—C13—C18—C17	-179.1 (3)
C3—C4—C5—C6	-1.1 (4)	C11—C12—C19—C24	-77.7 (3)
C3—C4—C5—N1	-177.3 (3)	C7—C12—C19—C24	99.5 (3)
C2—C1—C6—C5	0.2 (4)	C11—C12—C19—C20	100.7 (3)
C2—C1—C6—C7	177.5 (3)	C7—C12—C19—C20	-82.1 (3)
C4—C5—C6—C1	0.8 (4)	C24—C19—C20—C21	0.0 (4)
N1—C5—C6—C1	177.7 (2)	C12—C19—C20—C21	-178.5 (2)
C4—C5—C6—C7	-177.2 (2)	C19—C20—C21—C22	0.2 (4)
N1—C5—C6—C7	-0.2 (3)	C20—C21—C22—C23	-0.5 (5)
C1—C6—C7—C8	-177.7 (3)	C21—C22—C23—C24	0.7 (5)
C5—C6—C7—C8	-0.2 (3)	C22—C23—C24—C19	-0.6 (5)
C1—C6—C7—C12	1.5 (5)	C20—C19—C24—C23	0.2 (4)
C5—C6—C7—C12	179.0 (2)	C12—C19—C24—C23	178.7 (3)
C12—C7—C8—C9	-0.9 (4)	C30—C25—C26—C27	0.0 (4)
C6—C7—C8—C9	178.4 (2)	S1—C25—C26—C27	-179.2 (2)
C12—C7—C8—N1	-178.8 (2)	C25—C26—C27—C28	0.5 (5)
C6—C7—C8—N1	0.5 (3)	C26—C27—C28—C29	-1.2 (5)
C7—C8—C9—C10	0.4 (4)	C27—C28—C29—C30	1.2 (6)
N1—C8—C9—C10	177.9 (2)	C28—C29—C30—C25	-0.6 (5)
C8—C9—C10—O3	179.7 (2)	C26—C25—C30—C29	0.0 (4)
C8—C9—C10—C11	0.1 (4)	S1—C25—C30—C29	179.2 (2)
O3—C10—C11—C12	-179.8 (2)	C4—C5—N1—C8	177.2 (3)
C9—C10—C11—C12	-0.1 (4)	C6—C5—N1—C8	0.6 (3)
O3—C10—C11—C13	1.3 (3)	C4—C5—N1—S1	-33.7 (4)
C9—C10—C11—C13	-179.0 (2)	C6—C5—N1—S1	149.69 (18)
C10—C11—C12—C7	-0.4 (3)	C9—C8—N1—C5	-178.4 (2)
C13—C11—C12—C7	178.1 (2)	C7—C8—N1—C5	-0.7 (3)
C10—C11—C12—C19	177.0 (2)	C9—C8—N1—S1	32.6 (3)
C13—C11—C12—C19	-4.5 (4)	C7—C8—N1—S1	-149.66 (18)
C8—C7—C12—C11	0.9 (3)	C9—C10—O3—C14	179.7 (2)
C6—C7—C12—C11	-178.3 (2)	C11—C10—O3—C14	-0.7 (3)
C8—C7—C12—C19	-176.5 (2)	C15—C14—O3—C10	-179.2 (3)
C6—C7—C12—C19	4.4 (4)	C13—C14—O3—C10	-0.3 (3)
C12—C11—C13—C14	179.9 (3)	C5—N1—S1—O2	171.21 (19)
C10—C11—C13—C14	-1.4 (3)	C8—N1—S1—O2	-44.3 (2)
C12—C11—C13—C18	-0.5 (5)	C5—N1—S1—O1	42.1 (2)
C10—C11—C13—C18	178.1 (3)	C8—N1—S1—O1	-173.43 (19)
C18—C13—C14—C15	0.4 (4)	C5—N1—S1—C25	-73.3 (2)
C11—C13—C14—C15	180.0 (3)	C8—N1—S1—C25	71.2 (2)
C18—C13—C14—O3	-178.6 (2)	C30—C25—S1—O2	-167.4 (2)
C11—C13—C14—O3	1.1 (3)	C26—C25—S1—O2	11.8 (3)
C13—C14—C15—C16	-0.5 (4)	C30—C25—S1—O1	-35.4 (3)
O3—C14—C15—C16	178.3 (2)	C26—C25—S1—O1	143.8 (2)

C14—C15—C16—C17	−0.1 (5)	C30—C25—S1—N1	78.5 (2)
C15—C16—C17—C18	0.8 (5)	C26—C25—S1—N1	−102.3 (2)

Hydrogen-bond geometry (Å, °)

Cg3 and Cg4 are the centroids of rings C1—C6 and C7—C12, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
C4—H4···O1	0.93	2.34	2.924 (4)	121
C9—H9···O2	0.93	2.34	2.926 (3)	121
C2—H2···O3 ⁱ	0.93	2.57	3.464 (4)	160
C17—H17···Cg4 ⁱⁱ	0.93	2.81	3.683 (3)	156
C22—H22···Cg3 ⁱⁱⁱ	0.93	2.95	3.722 (3)	141

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, y+1/2, -z+3/2$; (iii) $-x, y+1/2, -z+3/2$.