

Received 26 November 2016

Accepted 27 November 2016

Edited by H. Stoeckli-Evans, University of
Neuchâtel, Switzerland**Keywords:** crystal structure; cone-shaped
conformation; hydrogen bonds; crown ether;
supramolecular compounds..**CCDC reference:** 1519443**Supporting information:** this article has
supporting information at journals.iucr.org/e

Crystal structure of (6*E*,20*E*)-3,24-difluoro-13,14,28,29-tetrahydro-5*H*,22*H*-tetrabenzo-[*e,j,p,u*][1,4,12,15]tetraoxacyclodocosine-5,22-dione

Shaaban K. Mohamed,^a Mehmet Akkurt,^b Farouq E. Hawaiz,^{c*} Mzgin M Ayoob^c
and Eric Hosten^d^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester, M1 5GD, England, and, Chemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^cChemistry Department, College of Education, Salahaddin University-Hawler, Erbil, Kurdistan Region, Iraq, and ^dChemistry Department, Summerstrand Campus (South), PO Box 77000, Nelson Mandela, Metropolitan University, South Africa. *Correspondence e-mail: shaabankamel@yahoo.com


The conformation of the title compound, C₃₄H₂₆F₂O₆, is cone-shaped, partially determined by intramolecular C—H···O short contacts. The benzene rings at the top of the cone are inclined to one another by 73.10 (7)°, while the benzene rings at the bottom of the cone are inclined to one another by 35.49 (8)°. In the crystal, molecules are linked by C—H···O and C—H···F hydrogen bonds, forming a three-dimensional supramolecular structure. There are also C—H···π contacts present within the framework structure.

1. Chemical context


Macrocyclic compounds are known for their various applications, particularly in coordination chemistry (Delgado; 1995). The study of synthetic macrocyclic compounds is an important area of chemistry in view of their presence in many biologically significant naturally occurring metal complexes. Such compounds have received special attention because of their presence in many important biological systems such as metallo-porphyrins (for example haemoglobin, myoglobin, cytochromes, chlorophylls), corrins (vitamin B12) and antibiotics (valinomycin, nonactin) with antibiotic, antifungal, anticancer and immunosuppressive activities as seen for erythromycin (McGuire *et al.*, 1952; Woodward *et al.*; 1981), amphotericin B (Vandeputte *et al.*, 1956; Nicolaou *et al.*, 1988), epithilone B (Gerth *et al.*, 1996; Bode & Carreira; 2001) and rapamycin (Vezina *et al.*, 1975; Smith *et al.*, 1997). In addition, macrocyclic compounds having ether linkages and chalcone moieties have important applications (Rina *et al.*, 2012, Matsushima *et al.*, 2001). In this context the title compound was prepared and herein we report on its synthesis and crystal structure.

2. Structural commentary

The title compound, Fig. 1, has a cone-shaped conformation, partially determined by intramolecular C—H···O short contacts (Table 1 and Fig. 1). The benzene rings at the top of the cone (C11–C16 and C31–C36) are inclined to one another by 73.10 (7)°, while the benzene rings at the bottom of the cone (C21–C26 and C41–C46) are inclined to one another by



35.49 (8)° (Fig. 2). The bond lengths and angles are similar to those observed in one of the starting materials for the synthesis of the title compound, *viz.* 2,2'-[ethane-1,2-diylbis(oxy)]-dibenzaldehyde (Aravindan *et al.*, 2003; Zhang *et al.*, 2003); both measured at room temperature. A low temperature (120 K) structure analysis of the same compound has also been reported (Akkurt *et al.*, 2013).



3. Supramolecular features

In the crystal, molecules are linked by C—H...O and C—H...F hydrogen bonds, forming a three-dimensional supramolecular structure (Fig. 3 and Table 1). There are also C—H... π interactions present, involving inversion-related molecules, within the three-dimensional framework (Table 1).

4. Database survey

A search of the Cambridge Structural Database (Version 5.37, update May 2016; Groom *et al.*, 2016) indicated the presence of the subunit 1,2-bis(2-vinylphenoxy)ethane in a number of macrocyclic-type compounds. However, no macrocyclic-type compounds were found containing the subunit 2,2'-[ethane-




Figure 2
A CPK model of the title compound, illustrating the cone-shaped conformation.

1,2-diylbis(oxy)]dibenzaldehyde. The title compound, which contains both these subunits, is unique; no other reports of molecules of this type were found.




Figure 1
A view of the molecular structure of the title compound, with atom labelling and 50% probability displacement ellipsoids. The short intramolecular C—H...O contacts are shown as dashed lines (see Table 1).




Figure 3
The crystal packing of the title compound, viewed along the *b* axis. Hydrogen bonds are shown as dashed lines (see Table 1), and for clarity only the H atoms involved in hydrogen bonding have been included.

Table 1
Hydrogen-bond geometry (Å, °).

Cg3 is the centroid of the C31–C36 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
C5–H5...O4	0.95	2.35	2.7023 (16)	101
C9–H9...O1	0.95	2.40	2.7281 (16)	100
C4–H4...F2 ⁱ	0.95	2.37	3.1387 (17)	138
C15–H15...O3 ⁱⁱ	0.95	2.51	3.3211 (19)	143
C33–H33...F2 ⁱⁱⁱ	0.95	2.53	3.483 (2)	176
C34–H34...O6 ^{iv}	0.95	2.51	3.3649 (17)	150
C36–H36...F2 ⁱ	0.95	2.43	3.3380 (19)	161
C44–H44...O1 ^v	0.95	2.58	3.4986 (18)	163
C46–H46...Cg3 ⁱⁱⁱ	0.95	2.84	3.6829 (16)	149

Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (v) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

5. Synthesis and crystallization

The title compound has been synthesized by two methods, illustrated in Fig. 4.

Method (a): High-dilution method

A mixture of 2,2'-(ethane-1,2-diylbis(oxy))dibenzaldehyde (**A**) (67.6 mg; 0.25 mmol) and 1,1'-((ethane-1,2-diylbis(oxy))-bis(5-fluoro-2,1-phenylene))bis(ethan-1-one) (**B**) (83.6 mg; 0.25 mmol) was dissolved in a KOH solution (10%, 130–160 ml) in MeOH/H₂O (3:1) and the mixture was refluxed for 6 h. The reaction mixture was left at room temperature with stirring for *ca* four days, then the solvent was reduced to nearly half volume under reduced pressure. The resulting precipitate was collected by filtration, dried and recrystallized from chloroform/methanol solution (1:1) to give yellow block-shaped crystals, suitable for x-ray diffraction (yield 80%, m.p. 553–554 K).

Method (b): Ultrasound-assisted synthesis

Compound **A** (0.55 mmol, 0.15 gm) was dissolved in ethanol (5 ml) and added to a solution of compound **B** (0.55 mmol) in ethanol (5 ml), and solid NaOH (0.3 gm) was added to the mixture. The mixture was then irradiated in the water bath of




Figure 4
Reaction scheme.

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₃₄ H ₂₆ F ₂ O ₆
<i>M_r</i>	568.55
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁ / <i>c</i>
Temperature (K)	200
<i>a</i> , <i>b</i> , <i>c</i> (Å)	16.2618 (7), 11.6708 (5), 14.7359 (7)
β (°)	96.945 (2)
<i>V</i> (Å ³)	2776.2 (2)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.10
Crystal size (mm)	0.63 × 0.29 × 0.15
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2009)
<i>T_{min}</i> , <i>T_{max}</i>	0.894, 1.000
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	38582, 6911, 5302
<i>R_{int}</i>	0.020
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.668
Refinement	
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.040, 0.106, 1.01
No. of reflections	6911
No. of parameters	379
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.32, -0.39

Computer programs: *APEX2* and *SAINT* (Bruker, 2009), *SHELXT2014/7* (Sheldrick, 2015a), *SHELXL2014/7* (Sheldrick, 2015b) and *PLATON* (Spek, 2009).

an ultrasonic cleaner at room temperature for 20 min. The mixture solidified and the product was separated by filtration under vacuum, washed with ethanol, dried and purified by recrystallization from chloroform solution (yield 74%). Single crystals were obtained by slow evaporation of a dilute solution of the title compound in chloroform over 13 days at room temperature (m.p. 553–554 K).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were positioned geometrically and refined using a riding model: C–H = 0.95–0.99 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Acknowledgements

The authors gratefully acknowledge the X-ray staff at Nelson Mandela Metropolitan University, South Africa for providing the X-ray data.

References

- Akkurt, M., Mohamed, S. K., Horton, P. N., Abdel-Raheem, E. M. M. & Albayati, M. R. (2013). *Acta Cryst.* **E69**, o1260.
 Aravindan, P. G., Yogavel, M., Thirumavalavan, M., Akilan, P., Velmurugan, D., Kandaswamy, M., Shanmuga Sundara Raj, S. & Fun, H.-K. (2003). *Acta Cryst.* **E59**, o806–o807.
 Bode, J. W. & Carreira, E. M. (2001). *J. Org. Chem.* **66**, 6410–6424.
 Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.

- Delgado, R. (1995). *Revista Portuguesa de Química*, **2**, 18–29.
- Gerth, K., Bedorf, N., Höfle, G., Irschik, H. & Reichenbach, H. (1996). *J. Antibiot.* **49**, 560–563.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Matsushima, R., Fujimoto, S. & Tokumura, K. (2001). *Bull. Chem. Soc. Jpn.*, **74**, 827–832.
- McGuire, J. M., Bunch, R. L., Anderson, R. C., Boaz, H. E., Flynn, E. H., Powell, H. M. & Smith, J. W. (1952). *Antibiot. Chemother.* **2**, 281–283.
- Nicolaou, K. C., Daines, O. Y., Ogawa, Y. & Chakraborty, T. K. (1988). *J. Am. Chem. Soc.* **110**, 4696–4705.
- Rina, M., Mandal, T. K., Asok, K. & Mallik, A. K. (2012). *ARKIVOC* (ix), pp. 95–110.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Smith, A. B., Condon, S. M., McCauley, J. A., Leazer, J. L., Leahy, J. W. & Maleczka, R. E. (1997). *J. Am. Chem. Soc.* **119**, 962–973.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Vandeputte, J., Wachtel, J. L. & Stiller, E. T. (1956). *Antibiot. Annu.* 587–591.
- Vézina, C., Kudelski, A. & Sehgal, S. N. (1975). *J. Antibiot.* **28**, 721–726.
- Woodward, R. B., Logusch, E., Nambiar, K. P., Sakan, K., Ward, D. E., Au-Yeung, B. W., Balaram, P., Browne, L. J., Card, P. J. & Chen, C. H. (1981). *J. Am. Chem. Soc.* **103**, 3215–3217.
- Zhang, J., Han, Q., Yang, X., Xu, X. & Wang, X. (2003). *Nanjing Ligong Daxue Xuebao*, **27**, 128.

supporting information

Acta Cryst. (2017). E73, 13-16 [https://doi.org/10.1107/S2056989016018934]

Crystal structure of (6*E*,20*E*)-3,24-difluoro-13,14,28,29-tetrahydro-5*H*,22*H*-tetrabenzo[*e,j,p,u*][1,4,12,15]tetraoxacyclodocosine-5,22-dione

Shaaban K. Mohamed, Mehmet Akkurt, Farouq E. Hawaiz, Mzgin M Ayoob and Eric Hosten

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE* (Bruker, 2009); program(s) used to solve structure: *SHELXT2014/7* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015b); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2014/7* (Sheldrick, 2015b) and *PLATON* (Spek, 2009).

(6*E*,20*E*)-3,24-Difluoro-13,14,28,29-tetrahydro-5*H*,22*H*-tetrabenzo[*e,j,p,u*][1,4,12,15]tetraoxacyclodocosine-5,22-dione

Crystal data

C₃₄H₂₆F₂O₆

M_r = 568.55

Monoclinic, *P2₁/c*

a = 16.2618 (7) Å

b = 11.6708 (5) Å

c = 14.7359 (7) Å

β = 96.945 (2)°

V = 2776.2 (2) Å³

Z = 4

F(000) = 1184

D_x = 1.360 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 9878 reflections

θ = 2.5–28.2°

μ = 0.10 mm⁻¹

T = 200 K

Block, yellow

0.63 × 0.29 × 0.15 mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: sealed tube

Graphite monochromator

Detector resolution: 8.3333 pixels mm⁻¹

φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2009)

T_{min} = 0.894, *T_{max}* = 1.000

38582 measured reflections

6911 independent reflections

5302 reflections with *I* > 2σ(*I*)

R_{int} = 0.020

θ_{max} = 28.3°, θ_{min} = 2.5°

h = -21 → 20

k = -15 → 15

l = -19 → 19

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.040

wR(*F*²) = 0.106

S = 1.01

6911 reflections

379 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0392P)^2 + 1.1029P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	-0.00518 (7)	-0.16078 (10)	0.73153 (8)	0.0712 (3)
F2	0.30286 (7)	0.77217 (11)	0.36051 (9)	0.0823 (4)
O1	0.07789 (6)	0.21042 (8)	0.53657 (8)	0.0426 (2)
O2	0.12188 (7)	0.44266 (9)	0.50741 (7)	0.0479 (3)
O3	0.13697 (6)	0.63181 (11)	0.65378 (8)	0.0567 (3)
O4	0.45695 (6)	0.39443 (9)	0.59069 (6)	0.0389 (2)
O5	0.41025 (6)	0.24383 (8)	0.44309 (6)	0.0365 (2)
O6	0.23814 (6)	-0.05205 (9)	0.57921 (8)	0.0473 (3)
C1	0.02059 (8)	0.30158 (12)	0.51429 (11)	0.0423 (3)
H1A	0.0096	0.3429	0.5703	0.051*
H1B	-0.0325	0.2714	0.4835	0.051*
C2	0.05993 (9)	0.37973 (13)	0.45186 (11)	0.0433 (3)
H2A	0.0851	0.3349	0.4053	0.052*
H2B	0.0181	0.4324	0.4202	0.052*
C3	0.20104 (8)	0.60305 (12)	0.62506 (10)	0.0376 (3)
C4	0.27425 (8)	0.57029 (12)	0.68694 (10)	0.0388 (3)
H4	0.2730	0.5805	0.7507	0.047*
C5	0.34280 (8)	0.52688 (11)	0.65906 (9)	0.0335 (3)
H5	0.3429	0.5177	0.5950	0.040*
C6	0.51460 (8)	0.32791 (12)	0.54755 (10)	0.0368 (3)
H6A	0.5213	0.2511	0.5760	0.044*
H6B	0.5693	0.3661	0.5535	0.044*
C7	0.48031 (8)	0.31774 (12)	0.44922 (9)	0.0372 (3)
H7A	0.4638	0.3942	0.4242	0.045*
H7B	0.5228	0.2859	0.4135	0.045*
C8	0.26989 (8)	0.11337 (11)	0.44357 (9)	0.0337 (3)
H8	0.3169	0.0888	0.4834	0.040*
C9	0.19546 (8)	0.08387 (12)	0.46483 (9)	0.0352 (3)
H9	0.1477	0.1089	0.4264	0.042*
C10	0.18427 (8)	0.01401 (11)	0.54551 (9)	0.0336 (3)
C11	0.05441 (8)	0.11931 (11)	0.58468 (9)	0.0341 (3)
C12	0.10509 (8)	0.02187 (11)	0.58816 (9)	0.0321 (3)
C13	0.08374 (9)	-0.07270 (12)	0.63846 (10)	0.0387 (3)
H13	0.1166	-0.1402	0.6413	0.046*
C14	0.01517 (10)	-0.06726 (14)	0.68353 (10)	0.0454 (4)
C15	-0.03344 (9)	0.02861 (14)	0.68325 (11)	0.0468 (4)

H15	-0.0795	0.0305	0.7173	0.056*
C16	-0.01442 (9)	0.12229 (13)	0.63273 (10)	0.0421 (3)
H16	-0.0482	0.1889	0.6306	0.050*
C21	0.16576 (9)	0.52274 (13)	0.46597 (10)	0.0399 (3)
C22	0.20682 (8)	0.60381 (12)	0.52387 (10)	0.0352 (3)
C23	0.25285 (8)	0.68864 (13)	0.48752 (12)	0.0452 (4)
H23	0.2808	0.7455	0.5257	0.054*
C24	0.25707 (9)	0.68851 (16)	0.39503 (13)	0.0559 (5)
C25	0.21969 (11)	0.6085 (2)	0.33750 (12)	0.0684 (6)
H25	0.2256	0.6099	0.2742	0.082*
C26	0.17280 (11)	0.52466 (18)	0.37308 (11)	0.0607 (5)
H26	0.1454	0.4683	0.3339	0.073*
C31	0.41776 (8)	0.49208 (11)	0.71727 (9)	0.0329 (3)
C32	0.47646 (8)	0.42363 (11)	0.68041 (9)	0.0332 (3)
C33	0.54911 (8)	0.39044 (13)	0.73364 (10)	0.0393 (3)
H33	0.5881	0.3429	0.7085	0.047*
C34	0.56427 (9)	0.42722 (14)	0.82355 (10)	0.0442 (3)
H34	0.6142	0.4056	0.8597	0.053*
C35	0.50782 (10)	0.49464 (14)	0.86091 (10)	0.0455 (4)
H35	0.5188	0.5196	0.9225	0.055*
C36	0.43489 (9)	0.52605 (13)	0.80844 (10)	0.0409 (3)
H36	0.3957	0.5716	0.8350	0.049*
C41	0.35808 (8)	0.24427 (11)	0.36284 (9)	0.0333 (3)
C42	0.28472 (8)	0.18095 (11)	0.36335 (9)	0.0337 (3)
C43	0.22821 (9)	0.18261 (13)	0.28370 (10)	0.0429 (3)
H43	0.1776	0.1418	0.2828	0.051*
C44	0.24391 (10)	0.24171 (15)	0.20655 (11)	0.0503 (4)
H44	0.2045	0.2416	0.1534	0.060*
C45	0.31735 (11)	0.30101 (14)	0.20740 (11)	0.0501 (4)
H45	0.3289	0.3408	0.1542	0.060*
C46	0.37440 (9)	0.30306 (13)	0.28523 (10)	0.0419 (3)
H46	0.4246	0.3447	0.2854	0.050*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0696 (7)	0.0713 (7)	0.0754 (7)	-0.0146 (6)	0.0193 (6)	0.0290 (6)
F2	0.0549 (6)	0.0891 (8)	0.1024 (9)	-0.0128 (6)	0.0076 (6)	0.0586 (7)
O1	0.0308 (5)	0.0332 (5)	0.0649 (7)	0.0023 (4)	0.0108 (4)	0.0063 (5)
O2	0.0516 (6)	0.0491 (6)	0.0411 (6)	-0.0208 (5)	-0.0019 (5)	0.0028 (5)
O3	0.0340 (5)	0.0756 (8)	0.0623 (7)	0.0065 (5)	0.0129 (5)	-0.0085 (6)
O4	0.0311 (5)	0.0489 (6)	0.0351 (5)	0.0073 (4)	-0.0020 (4)	-0.0084 (4)
O5	0.0345 (5)	0.0411 (5)	0.0330 (5)	-0.0087 (4)	0.0010 (4)	-0.0006 (4)
O6	0.0344 (5)	0.0437 (6)	0.0630 (7)	0.0062 (4)	0.0029 (5)	0.0083 (5)
C1	0.0292 (6)	0.0323 (7)	0.0637 (10)	0.0012 (5)	-0.0018 (6)	-0.0020 (6)
C2	0.0398 (7)	0.0360 (7)	0.0506 (8)	-0.0017 (6)	-0.0087 (6)	-0.0006 (6)
C3	0.0306 (6)	0.0341 (7)	0.0485 (8)	-0.0021 (5)	0.0063 (6)	-0.0064 (6)
C4	0.0370 (7)	0.0429 (8)	0.0363 (7)	-0.0010 (6)	0.0031 (5)	-0.0087 (6)

C5	0.0341 (6)	0.0335 (7)	0.0324 (6)	-0.0010 (5)	0.0019 (5)	-0.0022 (5)
C6	0.0268 (6)	0.0379 (7)	0.0453 (8)	0.0013 (5)	0.0021 (5)	-0.0061 (6)
C7	0.0329 (6)	0.0390 (7)	0.0408 (7)	-0.0060 (6)	0.0087 (5)	-0.0057 (6)
C8	0.0298 (6)	0.0342 (7)	0.0361 (7)	-0.0005 (5)	0.0003 (5)	-0.0046 (5)
C9	0.0303 (6)	0.0383 (7)	0.0364 (7)	0.0015 (5)	0.0009 (5)	-0.0041 (6)
C10	0.0275 (6)	0.0322 (6)	0.0401 (7)	-0.0028 (5)	-0.0008 (5)	-0.0046 (5)
C11	0.0288 (6)	0.0327 (6)	0.0401 (7)	-0.0067 (5)	0.0013 (5)	-0.0045 (5)
C12	0.0274 (6)	0.0345 (7)	0.0331 (6)	-0.0046 (5)	-0.0013 (5)	-0.0036 (5)
C13	0.0373 (7)	0.0374 (7)	0.0393 (7)	-0.0058 (6)	-0.0032 (6)	0.0012 (6)
C14	0.0448 (8)	0.0500 (9)	0.0410 (8)	-0.0150 (7)	0.0034 (6)	0.0065 (7)
C15	0.0398 (8)	0.0572 (9)	0.0453 (8)	-0.0138 (7)	0.0129 (6)	-0.0087 (7)
C16	0.0338 (7)	0.0408 (8)	0.0524 (9)	-0.0051 (6)	0.0086 (6)	-0.0122 (6)
C21	0.0343 (7)	0.0443 (8)	0.0395 (7)	-0.0034 (6)	-0.0021 (6)	0.0082 (6)
C22	0.0248 (6)	0.0350 (7)	0.0451 (7)	0.0041 (5)	0.0016 (5)	0.0059 (6)
C23	0.0299 (7)	0.0387 (8)	0.0661 (10)	0.0003 (6)	0.0020 (6)	0.0107 (7)
C24	0.0336 (7)	0.0656 (11)	0.0672 (11)	-0.0028 (7)	0.0011 (7)	0.0377 (9)
C25	0.0500 (10)	0.1086 (16)	0.0438 (9)	-0.0169 (10)	-0.0062 (7)	0.0300 (10)
C26	0.0565 (10)	0.0842 (13)	0.0384 (8)	-0.0195 (9)	-0.0065 (7)	0.0085 (8)
C31	0.0329 (6)	0.0339 (6)	0.0314 (6)	-0.0037 (5)	0.0021 (5)	0.0016 (5)
C32	0.0317 (6)	0.0343 (7)	0.0326 (6)	-0.0045 (5)	-0.0002 (5)	0.0012 (5)
C33	0.0315 (6)	0.0422 (8)	0.0426 (8)	-0.0012 (6)	-0.0016 (5)	0.0035 (6)
C34	0.0383 (7)	0.0506 (9)	0.0405 (8)	-0.0078 (6)	-0.0084 (6)	0.0112 (7)
C35	0.0506 (8)	0.0539 (9)	0.0304 (7)	-0.0085 (7)	-0.0025 (6)	0.0022 (6)
C36	0.0441 (8)	0.0448 (8)	0.0337 (7)	-0.0022 (6)	0.0038 (6)	-0.0022 (6)
C41	0.0344 (6)	0.0337 (6)	0.0315 (6)	0.0026 (5)	0.0029 (5)	-0.0031 (5)
C42	0.0332 (6)	0.0338 (7)	0.0340 (7)	0.0021 (5)	0.0033 (5)	-0.0037 (5)
C43	0.0355 (7)	0.0469 (8)	0.0442 (8)	-0.0007 (6)	-0.0036 (6)	-0.0021 (6)
C44	0.0521 (9)	0.0539 (9)	0.0411 (8)	0.0043 (8)	-0.0105 (7)	0.0036 (7)
C45	0.0602 (10)	0.0506 (9)	0.0383 (8)	0.0006 (8)	0.0010 (7)	0.0085 (7)
C46	0.0438 (8)	0.0429 (8)	0.0392 (7)	-0.0038 (6)	0.0055 (6)	0.0021 (6)

Geometric parameters (Å, °)

F1—C14	1.3628 (18)	C12—C13	1.3966 (19)
F2—C24	1.3632 (18)	C13—C14	1.367 (2)
O1—C11	1.3581 (16)	C13—H13	0.9500
O1—C1	1.4262 (16)	C14—C15	1.370 (2)
O2—C21	1.3643 (17)	C15—C16	1.379 (2)
O2—C2	1.4230 (17)	C15—H15	0.9500
O3—C3	1.2183 (16)	C16—H16	0.9500
O4—C32	1.3651 (16)	C21—C26	1.388 (2)
O4—C6	1.4247 (16)	C21—C22	1.390 (2)
O5—C41	1.3695 (15)	C22—C23	1.387 (2)
O5—C7	1.4231 (15)	C23—C24	1.373 (2)
O6—C10	1.2261 (16)	C23—H23	0.9500
C1—C2	1.493 (2)	C24—C25	1.355 (3)
C1—H1A	0.9900	C25—C26	1.382 (3)
C1—H1B	0.9900	C25—H25	0.9500

C2—H2A	0.9900	C26—H26	0.9500
C2—H2B	0.9900	C31—C36	1.3962 (19)
C3—C4	1.460 (2)	C31—C32	1.4035 (19)
C3—C22	1.505 (2)	C32—C33	1.3918 (18)
C4—C5	1.3338 (19)	C33—C34	1.386 (2)
C4—H4	0.9500	C33—H33	0.9500
C5—C31	1.4608 (18)	C34—C35	1.374 (2)
C5—H5	0.9500	C34—H34	0.9500
C6—C7	1.4931 (19)	C35—C36	1.385 (2)
C6—H6A	0.9900	C35—H35	0.9500
C6—H6B	0.9900	C36—H36	0.9500
C7—H7A	0.9900	C41—C46	1.3867 (19)
C7—H7B	0.9900	C41—C42	1.4041 (18)
C8—C9	1.3317 (18)	C42—C43	1.4006 (19)
C8—C42	1.4652 (19)	C43—C44	1.380 (2)
C8—H8	0.9500	C43—H43	0.9500
C9—C10	1.471 (2)	C44—C45	1.379 (2)
C9—H9	0.9500	C44—H44	0.9500
C10—C12	1.5022 (18)	C45—C46	1.385 (2)
C11—C16	1.3955 (19)	C45—H45	0.9500
C11—C12	1.4017 (19)	C46—H46	0.9500
C11—O1—C1	119.15 (11)	C14—C15—H15	120.6
C21—O2—C2	117.92 (11)	C16—C15—H15	120.6
C32—O4—C6	118.67 (10)	C15—C16—C11	120.02 (14)
C41—O5—C7	117.40 (10)	C15—C16—H16	120.0
O1—C1—C2	106.30 (11)	C11—C16—H16	120.0
O1—C1—H1A	110.5	O2—C21—C26	124.24 (14)
C2—C1—H1A	110.5	O2—C21—C22	115.48 (12)
O1—C1—H1B	110.5	C26—C21—C22	120.25 (14)
C2—C1—H1B	110.5	C23—C22—C21	119.38 (14)
H1A—C1—H1B	108.7	C23—C22—C3	119.20 (13)
O2—C2—C1	106.68 (12)	C21—C22—C3	121.41 (12)
O2—C2—H2A	110.4	C24—C23—C22	118.46 (15)
C1—C2—H2A	110.4	C24—C23—H23	120.8
O2—C2—H2B	110.4	C22—C23—H23	120.8
C1—C2—H2B	110.4	C25—C24—F2	118.96 (17)
H2A—C2—H2B	108.6	C25—C24—C23	123.32 (15)
O3—C3—C4	121.47 (14)	F2—C24—C23	117.70 (17)
O3—C3—C22	120.06 (13)	C24—C25—C26	118.48 (17)
C4—C3—C22	118.44 (12)	C24—C25—H25	120.8
C5—C4—C3	123.76 (13)	C26—C25—H25	120.8
C5—C4—H4	118.1	C25—C26—C21	120.06 (17)
C3—C4—H4	118.1	C25—C26—H26	120.0
C4—C5—C31	126.44 (13)	C21—C26—H26	120.0
C4—C5—H5	116.8	C36—C31—C32	117.91 (12)
C31—C5—H5	116.8	C36—C31—C5	122.77 (13)
O4—C6—C7	106.72 (10)	C32—C31—C5	119.30 (12)

O4—C6—H6A	110.4	O4—C32—C33	123.95 (12)
C7—C6—H6A	110.4	O4—C32—C31	115.31 (11)
O4—C6—H6B	110.4	C33—C32—C31	120.74 (12)
C7—C6—H6B	110.4	C34—C33—C32	119.55 (14)
H6A—C6—H6B	108.6	C34—C33—H33	120.2
O5—C7—C6	108.18 (11)	C32—C33—H33	120.2
O5—C7—H7A	110.1	C35—C34—C33	120.69 (13)
C6—C7—H7A	110.1	C35—C34—H34	119.7
O5—C7—H7B	110.1	C33—C34—H34	119.7
C6—C7—H7B	110.1	C34—C35—C36	119.78 (14)
H7A—C7—H7B	108.4	C34—C35—H35	120.1
C9—C8—C42	124.89 (12)	C36—C35—H35	120.1
C9—C8—H8	117.6	C35—C36—C31	121.33 (14)
C42—C8—H8	117.6	C35—C36—H36	119.3
C8—C9—C10	122.56 (12)	C31—C36—H36	119.3
C8—C9—H9	118.7	O5—C41—C46	123.62 (12)
C10—C9—H9	118.7	O5—C41—C42	115.56 (11)
O6—C10—C9	121.58 (12)	C46—C41—C42	120.82 (12)
O6—C10—C12	118.40 (12)	C43—C42—C41	117.39 (13)
C9—C10—C12	120.02 (11)	C43—C42—C8	121.89 (12)
O1—C11—C16	122.60 (13)	C41—C42—C8	120.70 (12)
O1—C11—C12	116.94 (11)	C44—C43—C42	121.96 (14)
C16—C11—C12	120.39 (13)	C44—C43—H43	119.0
C13—C12—C11	118.54 (12)	C42—C43—H43	119.0
C13—C12—C10	117.03 (12)	C45—C44—C43	119.33 (14)
C11—C12—C10	124.34 (12)	C45—C44—H44	120.3
C14—C13—C12	119.41 (14)	C43—C44—H44	120.3
C14—C13—H13	120.3	C44—C45—C46	120.58 (15)
C12—C13—H13	120.3	C44—C45—H45	119.7
F1—C14—C13	118.59 (15)	C46—C45—H45	119.7
F1—C14—C15	118.66 (14)	C45—C46—C41	119.89 (14)
C13—C14—C15	122.75 (14)	C45—C46—H46	120.1
C14—C15—C16	118.84 (14)	C41—C46—H46	120.1
C11—O1—C1—C2	-172.73 (12)	C4—C3—C22—C21	109.61 (15)
C21—O2—C2—C1	-178.45 (12)	C21—C22—C23—C24	-0.8 (2)
O1—C1—C2—O2	-75.81 (14)	C3—C22—C23—C24	-179.61 (13)
O3—C3—C4—C5	172.72 (15)	C22—C23—C24—C25	-1.2 (2)
C22—C3—C4—C5	-8.9 (2)	C22—C23—C24—F2	-179.71 (13)
C3—C4—C5—C31	-179.75 (13)	F2—C24—C25—C26	-179.42 (17)
C32—O4—C6—C7	-175.40 (11)	C23—C24—C25—C26	2.1 (3)
C41—O5—C7—C6	164.74 (11)	C24—C25—C26—C21	-0.9 (3)
O4—C6—C7—O5	-71.01 (14)	O2—C21—C26—C25	-179.07 (16)
C42—C8—C9—C10	178.86 (12)	C22—C21—C26—C25	-1.0 (3)
C8—C9—C10—O6	-25.3 (2)	C4—C5—C31—C36	-16.4 (2)
C8—C9—C10—C12	155.24 (13)	C4—C5—C31—C32	165.00 (14)
C1—O1—C11—C16	-16.33 (19)	C6—O4—C32—C33	-1.32 (19)
C1—O1—C11—C12	166.75 (12)	C6—O4—C32—C31	178.13 (12)

O1—C11—C12—C13	178.80 (11)	C36—C31—C32—O4	-179.10 (12)
C16—C11—C12—C13	1.80 (19)	C5—C31—C32—O4	-0.44 (18)
O1—C11—C12—C10	2.40 (19)	C36—C31—C32—C33	0.4 (2)
C16—C11—C12—C10	-174.60 (12)	C5—C31—C32—C33	179.02 (12)
O6—C10—C12—C13	-23.85 (18)	O4—C32—C33—C34	178.22 (13)
C9—C10—C12—C13	155.65 (12)	C31—C32—C33—C34	-1.2 (2)
O6—C10—C12—C11	152.60 (13)	C32—C33—C34—C35	0.9 (2)
C9—C10—C12—C11	-27.90 (18)	C33—C34—C35—C36	0.2 (2)
C11—C12—C13—C14	-0.82 (19)	C34—C35—C36—C31	-1.0 (2)
C10—C12—C13—C14	175.85 (12)	C32—C31—C36—C35	0.8 (2)
C12—C13—C14—F1	179.36 (13)	C5—C31—C36—C35	-177.86 (14)
C12—C13—C14—C15	-1.3 (2)	C7—O5—C41—C46	6.93 (19)
F1—C14—C15—C16	-178.23 (13)	C7—O5—C41—C42	-172.61 (11)
C13—C14—C15—C16	2.5 (2)	O5—C41—C42—C43	177.69 (12)
C14—C15—C16—C11	-1.4 (2)	C46—C41—C42—C43	-1.9 (2)
O1—C11—C16—C15	-177.51 (13)	O5—C41—C42—C8	-4.12 (18)
C12—C11—C16—C15	-0.7 (2)	C46—C41—C42—C8	176.32 (13)
C2—O2—C21—C26	-20.2 (2)	C9—C8—C42—C43	-26.3 (2)
C2—O2—C21—C22	161.70 (13)	C9—C8—C42—C41	155.60 (13)
O2—C21—C22—C23	-179.87 (12)	C41—C42—C43—C44	1.4 (2)
C26—C21—C22—C23	1.9 (2)	C8—C42—C43—C44	-176.79 (14)
O2—C21—C22—C3	-1.14 (19)	C42—C43—C44—C45	0.1 (2)
C26—C21—C22—C3	-179.34 (15)	C43—C44—C45—C46	-1.0 (3)
O3—C3—C22—C23	106.71 (16)	C44—C45—C46—C41	0.6 (2)
C4—C3—C22—C23	-71.66 (17)	O5—C41—C46—C45	-178.58 (14)
O3—C3—C22—C21	-72.03 (19)	C42—C41—C46—C45	0.9 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg3 is the centroid of the C31–C36 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5 \cdots O4	0.95	2.35	2.7023 (16)	101
C9—H9 \cdots O1	0.95	2.40	2.7281 (16)	100
C4—H4 \cdots F2 ⁱ	0.95	2.37	3.1387 (17)	138
C15—H15 \cdots O3 ⁱⁱ	0.95	2.51	3.3211 (19)	143
C33—H33 \cdots F2 ⁱⁱⁱ	0.95	2.53	3.483 (2)	176
C34—H34 \cdots O6 ^{iv}	0.95	2.51	3.3649 (17)	150
C36—H36 \cdots F2 ⁱ	0.95	2.43	3.3380 (19)	161
C44—H44 \cdots O1 ^v	0.95	2.58	3.4986 (18)	163
C46—H46 \cdots Cg3 ⁱⁱⁱ	0.95	2.84	3.6829 (16)	149

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $-x, y-1/2, -z+3/2$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x+1, y+1/2, -z+3/2$; (v) $x, -y+1/2, z-1/2$.