



Received 23 November 2016
Accepted 25 December 2016

Edited by D.-J. Xu, Zhejiang University (Yuquan Campus), China

Keywords: crystal structure; macroheterocycles; crown ethers; stacking interactions.

CCDC reference: 1524447

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of 22,24,25-trimethyl-8,11,14-trioxa-25-azatetracyclo[19.3.1.0^{2,7}.0^{15,20}]pentacosa-2,4,6,15(20),16,18-hexaen-23-one

Van Tuyen Nguyen,^{a,b*} Hong Hieu Truong,^c Tuan Anh Le,^d Anatoly T. Soldatenkov,^e Tuyet Anh Dang Thi,^a Thi Thanh Van Tran,^d Natalia Ya. Esina^f and Victor N. Khrustalev^{f,g}


^aInstitute of Chemistry, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam, ^bGraduate University of Science and Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam, ^cDepartment of Biotechnology, Vietnam–Russia Tropical Centre, 58 Nguyen Van Huyen, Hanoi, Vietnam, ^dFaculty of Chemistry, VNU University of Science, Vietnam National University, 19 Le Thanh Tong, Hoan Kiem, Hanoi, Vietnam, ^eOrganic Chemistry Department, Peoples' Friendship University of Russia, 6 Miklukho-Maklaya St., Moscow 117198, Russian Federation, ^fInorganic Chemistry Department, Peoples' Friendship University of Russia, 6 Miklukho-Maklay St., Moscow 117198, Russian Federation, and ^gX-Ray Structural Centre, A.N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 28 Vavilov St., B-334, Moscow 119991, Russian Federation. *Correspondence e-mail: ngvtuyen@hotmail.com

The title compound, C₂₄H₂₉NO₄, is the product of a Petrenko–Kritchenko condensation of 1,5-bis(2-formylphenoxy)-3-oxapentane, pentan-3-one and methylammonium acetate in ethanol. The molecule has mirror symmetry. The aza-14-crown-3 ether ring adopts a *bowl* conformation stabilized by a weak intramolecular C–H···O hydrogen bond. The conformation of the C–O–C–C–O–C–O–C polyether chain is t–g⁺–t–t–g[–]–t (t = *trans*, 180°; g = *gauche*, ±60°). The dihedral angle between the benzene rings fused to the aza-14-crown-4-ether moiety is 72.68 (4)°. The piperidinone ring adopts a *chair* conformation. The nitrogen atom has a trigonal–pyramidal geometry, the sum of the bond angles being 335.9°. In the crystal, the molecules are linked by weak C–H···O interactions, forming zigzag chains propagating along the [100] direction.

1. Chemical context


Macrocyclic heterocycles containing both crown ether and azaheterocyclic moieties are prospective compounds not only as metal-ion receptors (Pedersen, 1988), but also as membrane ion-transporting vehicles (Gökel & Murillo, 1996), as active components of environmental chemistry (Bradshaw & Izatt, 1997), for the design of organic sensors (Costero *et al.*, 2005), in nanosized on-off switches and other molecular electronic devices (Natali & Giordani, 2012). Moreover, they can possess antibacterial (An *et al.*, 1998) and anticancer properties (Artiemenko *et al.*, 2002; Le *et al.*, 2014, 2015), and other useful biological activity (Anh & Soldatenkov, 2016; Tran *et al.*, 2016).

Recently, we have developed effective methods for the synthesis of azacrown ethers containing γ-piperidone (Levov *et al.*, 2006, 2008; Anh *et al.*, 2008, 2012a,b,c; Hieu *et al.* (2011, 2012a,b) or γ-arylpyridine (Anh & Soldatenkov, 2016; Tran *et al.*, 2016) subunits. This chemistry allowed us to make systematic studies of the fine structural features of a novel series of azacrown macrocycles using X-ray diffraction. Such data should be of use for the subsequent design of more certain drug-like macroheterocyclic molecules bearing new would-be pharmacophore groups.



OPEN ACCESS

In attempts to apply this chemistry for obtaining azacrown ethers which contain a 1,3,5-trimethyl-substituted γ -piperidinone moiety, we studied the condensation of diethylketone and 1,5-bis(2-formylphenoxy)-3-oxapentane in the presence of methylammonium acetate taken both as the nitrogen source and as the template in ethanol/acetic acid solution. The reaction proceeded smoothly to give the expected azacrown title molecule, (I), with 32% yield.



2. Structural commentary

The title compound (Fig. 1), the product of a Petrenko-Kritchenko condensation of 1,5-bis(2-formylphenoxy)-3-oxapentane, pentan-3-one and methylammonium acetate in ethanol, crystallizes in the orthorhombic space group *Pnma* and, in the crystal, occupies a special position on a mirror plane. The aza-14-crown-3 ether ring adopts a *bowl* conformation stabilized by the weak intramolecular C16–H16B···O11 hydrogen bond (Table 1, Fig. 1). The distances from the center of the macrocycle cavity (defined as the centroid of atoms O8/O11/O8A/N14) to the O8, O11 and N14 atoms are 2.265 (2), 1.880 (2) and 2.393 (2) Å, respectively [atoms with the suffix *A* are related by the symmetry operation




Figure 1

The molecular structure of (I). Displacement ellipsoids are shown at the 50% probability level. H atoms are presented as small spheres of arbitrary radius. The dashed line indicates the intramolecular C–H···O hydrogen bond. Atoms with the suffix *A* are related by the symmetry operation $x, \frac{1}{2} - y, z$.

Table 1
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C10–H10B···O12 ⁱ	0.99	2.58	3.449 (2)	147
C16–H16B···O11	0.96	2.57	3.530 (3)	180

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, -z - \frac{1}{2}$.

$x, \frac{1}{2} - y, z$.]. The conformation of the C7–O8–C9–C10–O11–C10A–C9A–O8A–C7A polyether chain is t–g⁺–t–t–g[–]–t (*t* = *trans*, 180°; *g* = *gauche*, $\pm 60^\circ$). The dihedral angle between the planes of the benzene rings fused to the aza-14-crown-4-ether moiety is 72.68 (4)°. The piperidinone ring adopts a *chair* conformation. The nitrogen N14 atom has a trigonal–pyramidal geometry (sum of the bond angles is 335.9°).

The molecule of (I) possesses four asymmetric centers at the C1, C13, C13A and C1A carbon atoms and can have potentially numerous diastereomers. The crystal of (I) is racemic and consists of enantiomeric pairs with the following relative configuration of the centers: *rac*–1*R**–1*S**–1*3A**R**–1*AS**

3. Supramolecular features

In the crystal, molecules of (I) form zigzag chains along [100] via weak C–H···O interactions (Table 1, Figs. 2 and 3). π – π stacking is observed in the crystal, the distance between

Table 2
Experimental details.

Crystal data	$C_{24}H_{29}NO_4$
Chemical formula	395.48
M_r	Orthorhombic, <i>Pnma</i>
Crystal system, space group	120
Temperature (K)	8.1468 (5), 20.3402 (11), 12.0155 (7)
a, b, c (Å)	1991.1 (2)
V (Å ³)	4
Z	Mo $K\alpha$
Radiation type	0.09
μ (mm ^{−1})	0.20 × 0.15 × 0.15
Crystal size (mm)	
Data collection	Bruker APEXII CCD
Diffractometer	Multi-scan (SADABS; Bruker, 2003)
Absorption correction	0.970, 0.980
T_{\min}, T_{\max}	25079, 3128, 2469
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	0.065
R_{int}	($\sin \theta/\lambda$) _{max} (Å ^{−1})
	0.715
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.056, 0.130, 1.02
No. of reflections	3128
No. of parameters	140
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ^{−3})	0.34, −0.23

Computer programs: APEX2 (Bruker, 2001) and SAINT (Bruker, 2005), SHELXT (Sheldrick, 2015a), SHELXL2015 (Sheldrick, 2015b) and SHELXTL (Sheldrick, 2008).




Figure 2

Crystal packing of (I) along the *b* axis demonstrating the zigzag hydrogen-bonded chains along [100]. Dashed lines indicate the intra- and intermolecular C—H···O hydrogen bonds.




Figure 3

A portion of the crystal packing of (I) indicating the intermolecular π — π stacking interactions. Dashed lines indicate the intra- and intermolecular C—H···O hydrogen bonds.

parallel benzene rings is 3.446 (3) Å and the shortest intermolecular C5···C7ⁱ distance is 3.495 (2) Å [symmetry code: (i) 1 - x , 1 - y , - z].

4. Synthesis and crystallization

Methylammonium acetate (3.85 g, 50 mmol) was added to a solution of 1,5-bis(2-formylphenoxy)-3-oxapentane (3.14 g, 10.0 mmol) and diethylketone (1.41 g, 10.0 mmol) in ethanol/acetic acid (40 mL/1 mL) mixture. The reaction mixture was stirred at 293 K for three days (monitored by TLC until the disappearance of the starting heterocyclic ketone spot). At the end of the reaction, the formed precipitate was filtered off, washed with ethanol and recrystallized from ethanol solution to give 2.1 g of crystals of (I) (yield 32% m.p. 485–487 K). IR (KBr), ν/cm^{-1} : 1702. ¹H NMR (CDCl_3 , 400 MHz, 300 K): δ = 0.89 (d, 6H, CH_3 , J = 6.7 Hz), 1.86 (c, 3H, NCH_3), 3.19 (d, 2H, H1, H21, J = 10.8 Hz), 3.76–4.16 (m, 10H, H_{ether} and H22, H24), 6.78–7.26 (m, 8H, H_{arom}). Analysis calculated for $\text{C}_{24}\text{H}_{29}\text{NO}_4$: C, 67.72; H, 7.31; N, 5.64. Found: C, 67.54; H, 7.42; N, 5.41.

5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All hydrogen atoms were placed in calculated positions with C—H = 0.95 Å (aryl-H), 0.96 Å (methyl-H), 0.98 Å (methylene-H) or 1.00 Å (methine-H) and refined using a riding model with fixed isotropic displacement parameters [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for the methyl groups and $1.2U_{\text{eq}}(\text{C})$ for all other atoms].

Funding information

Funding for this research was provided by: National Foundation for Science and Technology Development (award No. 104.01-2014.39); Ministry of Education and Science of the Russian Federation (award No. 02.a03.21.0008 of June 24, 2016).

References

- An, H., Wang, T., Mohan, V., Griffey, R. H. & Cook, P. D. (1998). *Tetrahedron*, **54**, 3999–4012.
- Anh, L. T., Hieu, T. H., Soldatenkov, A. T., Kolyadina, N. M. & Khrustalev, V. N. (2012b). *Acta Cryst. E***68**, o1588–o1589.
- Anh, L. T., Hieu, T. H., Soldatenkov, A. T., Kolyadina, N. M. & Khrustalev, V. N. (2012c). *Acta Cryst. E***68**, o2165–o2166.
- Anh, L. T., Hieu, T. H., Soldatenkov, A. T., Soldatova, S. A. & Khrustalev, V. N. (2012a). *Acta Cryst. E***68**, o1386–o1387.
- Anh, L. T., Levov, A. N., Soldatenkov, A. T., Gruzdev, R. D. & Hieu, T. H. (2008). *Russ. J. Org. Chem.* **44**, 463–465.
- Artiemenko, A. G., Kovdiienko, N. A., Kuz'min, V. E., Kamalov, G. L., Lozitskaya, R. N., Fedchuk, A. S., Lozitsky, V. P., Dyachenko, N. S. & Nosach, L. N. (2002). *Exp. Oncol.* **24**, 123–127.
- Bradshaw, J. S. & Izatt, R. M. (1997). *Acc. Chem. Res.* **30**, 338–345.
- Bruker (2001). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2003). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
- Costero, A. M., Bañuls, M. J., Aurell, M. J., Ochando, L. E. & Doménech, A. J. (2005). *Tetrahedron*, **61**, 10309–10320.
- Gökel, G. W. & Murillo, O. (1996). *Acc. Chem. Res.* **29**, 425–432.
- Hieu, T. H., Anh, L. T., Soldatenkov, A. T., Golovtsov, N. I. & Soldatova, S. A. (2011). *Chem. Heterocycl. Compd.* **47**, 1307–1308.
- Hieu, T. H., Anh, L. T., Soldatenkov, A. T., Kolyadina, N. M. & Khrustalev, V. N. (2012a). *Acta Cryst. E***68**, o2431–o2432.
- Hieu, T. H., Anh, L. T., Soldatenkov, A. T., Kurilkin, V. V. & Khrustalev, V. N. (2012b). *Acta Cryst. E***68**, o2848–o2849.
- Le, A. T. & Soldatenkov, A. T. (2016). *Chem. Heterocycl. Compd.* **52**, 152–154.
- Le, T. A., Truong, H. H., Nguyen, P. T. T., Pham, H. T., Kotsuba, V. E., Soldatenkov, A. T., Khrustalev, V. N. & Wodajo, A. T. (2014). *Macroheterocycles*, **7**, 386–390.
- Le, A. T., Truong, H. H., Thi, T. P. N., Thi, N. D., To, H. T., Thi, H. P. & Soldatenkov, A. T. (2015). *Mendeleev Commun.* **25**, 224–225.
- Levov, A. N., Anh, L. T., Komarova, A. I., Strokina, V. M., Soldatenkov, A. T. & Khrustalev, V. N. (2008). *Russ. J. Org. Chem.* **44**, 457–462.
- Levov, A. N., Strokina, V. M., Anh, L. T., Komarova, A. I., Soldatenkov, A. T. & Khrustalev, V. N. (2006). *Mendeleev Commun.* **16**, 35–36.
- Natali, M. & Giordani, S. (2012). *Chem. Soc. Rev.* **41**, 4010–4029.
- Pedersen, C. J. (1988). *Angew. Chem.* **100**, 1053–1059.
- Sheldrick, G. M. (2008). *Acta Cryst. A***64**, 112–122.
- Sheldrick, G. M. (2015a). *Acta Cryst. A***71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst. C***71**, 3–8.
- Tran, T. T. V., Anh, L. T., Nguyen, H. H., Truong, H. H. & Soldatenkov, A. T. (2016). *Acta Cryst. E***72**, 663–666.

supporting information

Acta Cryst. (2017). E73, 118-121 [https://doi.org/10.1107/S2056989016020508]

Crystal structure of 22,24,25-trimethyl-8,11,14-trioxa-25-azatetracyclo-[19.3.1.0^{2,7}.0^{15,20}]pentacosa-2,4,6,15(20),16,18-hexaen-23-one

Van Tuyen Nguyen, Hong Hieu Truong, Tuan Anh Le, Anatoly T. Soldatenkov, Tuyet Anh Dang Thi, Thi Thanh Van Tran, Natalia Ya. Esina and Victor N. Khrustalev

Computing details

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT* (Bruker, 2001); program(s) used to solve structure: *SHELXT* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2015* (Sheldrick, 2015b); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

22,24,25-Trimethyl-8,11,14-trioxa-25-azatetracyclo[19.3.1.0^{2,7}.0^{15,20}]pentacosa-2,4,6,15 (20),16,18-hexaen-23-one

Crystal data

C₂₄H₂₉NO₄

M_r = 395.48

Orthorhombic, *Pnma*

a = 8.1468 (5) Å

b = 20.3402 (11) Å

c = 12.0155 (7) Å

V = 1991.1 (2) Å³

Z = 4

F(000) = 848

D_x = 1.319 Mg m⁻³

Mo Kα radiation, λ = 0.71073 Å

Cell parameters from 4090 reflections

θ = 3.0–29.2°

μ = 0.09 mm⁻¹

T = 120 K

Prism, colourless

0.20 × 0.15 × 0.15 mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2003)

T_{min} = 0.970, T_{max} = 0.980

25079 measured reflections

3128 independent reflections

2469 reflections with I > 2σ(I)

R_{int} = 0.065

θ_{max} = 30.6°, θ_{min} = 2.0°

h = -11→11

k = -29→29

l = -17→16

Refinement

Refinement on F²

Least-squares matrix: full

R[F² > 2σ(F²)] = 0.056

wR(F²) = 0.130

S = 1.02

3128 reflections

140 parameters

0 restraints

Primary atom site location: difference Fourier

map

Secondary atom site location: difference Fourier

map

Hydrogen site location: mixed

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.046P)^2 + 1.66P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.58338 (18)	0.30912 (7)	-0.07447 (12)	0.0151 (3)
H1	0.4639	0.3029	-0.0903	0.018*
C2	0.60218 (18)	0.37109 (7)	-0.00494 (12)	0.0156 (3)
C3	0.75023 (19)	0.40480 (7)	0.00526 (13)	0.0190 (3)
H3	0.8432	0.3900	-0.0354	0.023*
C4	0.7652 (2)	0.45976 (7)	0.07366 (14)	0.0223 (3)
H4	0.8670	0.4822	0.0793	0.027*
C5	0.6301 (2)	0.48134 (7)	0.13342 (13)	0.0223 (3)
H5	0.6398	0.5188	0.1803	0.027*
C6	0.4808 (2)	0.44885 (7)	0.12557 (13)	0.0195 (3)
H6	0.3888	0.4637	0.1672	0.023*
C7	0.46690 (18)	0.39412 (7)	0.05590 (12)	0.0162 (3)
O8	0.32384 (13)	0.35995 (5)	0.04244 (9)	0.0194 (2)
C9	0.19912 (19)	0.36553 (8)	0.12588 (13)	0.0201 (3)
H9A	0.1387	0.4074	0.1172	0.024*
H9B	0.2488	0.3645	0.2011	0.024*
C10	0.08483 (19)	0.30816 (7)	0.11056 (13)	0.0197 (3)
H10A	-0.0119	0.3129	0.1597	0.024*
H10B	0.0462	0.3062	0.0325	0.024*
O11	0.17272 (19)	0.2500	0.13785 (13)	0.0192 (3)
C12	0.6367 (3)	0.2500	-0.24937 (18)	0.0157 (4)
O12	0.5743 (2)	0.2500	-0.34102 (13)	0.0219 (3)
C13	0.67442 (18)	0.31314 (7)	-0.18726 (12)	0.0160 (3)
H13	0.7951	0.3147	-0.1722	0.019*
N14	0.6428 (2)	0.2500	-0.01435 (14)	0.0148 (3)
C15	0.6282 (2)	0.37347 (7)	-0.25565 (13)	0.0192 (3)
H15A	0.6833	0.3715	-0.3281	0.029*
H15B	0.6626	0.4133	-0.2161	0.029*
H15C	0.5091	0.3744	-0.2667	0.029*
C16	0.6033 (3)	0.2500	0.10474 (17)	0.0186 (4)
H16A	0.6489	0.2885	0.1390	0.028*
H16B	0.4864	0.2500	0.1141	0.028*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0192 (7)	0.0111 (6)	0.0151 (6)	0.0006 (5)	0.0007 (5)	-0.0001 (5)
C2	0.0197 (7)	0.0118 (6)	0.0154 (6)	0.0002 (5)	-0.0012 (5)	0.0008 (5)
C3	0.0200 (7)	0.0152 (6)	0.0217 (7)	0.0009 (5)	-0.0016 (6)	0.0010 (5)
C4	0.0240 (8)	0.0169 (7)	0.0259 (8)	-0.0028 (6)	-0.0061 (6)	-0.0005 (6)
C5	0.0317 (8)	0.0133 (6)	0.0219 (7)	0.0005 (6)	-0.0075 (6)	-0.0042 (6)
C6	0.0248 (7)	0.0161 (7)	0.0177 (7)	0.0031 (6)	-0.0015 (6)	-0.0026 (5)
C7	0.0202 (7)	0.0124 (6)	0.0158 (7)	0.0003 (5)	-0.0018 (5)	0.0004 (5)
O8	0.0198 (5)	0.0201 (5)	0.0183 (5)	-0.0026 (4)	0.0035 (4)	-0.0054 (4)
C9	0.0208 (7)	0.0203 (7)	0.0193 (7)	0.0025 (6)	0.0038 (6)	-0.0033 (6)
C10	0.0176 (7)	0.0217 (7)	0.0199 (7)	0.0033 (6)	0.0015 (6)	-0.0008 (6)
O11	0.0177 (7)	0.0183 (7)	0.0216 (8)	0.000	-0.0020 (6)	0.000
C12	0.0152 (9)	0.0157 (9)	0.0162 (9)	0.000	0.0047 (7)	0.000
O12	0.0306 (9)	0.0190 (7)	0.0160 (7)	0.000	-0.0006 (6)	0.000
C13	0.0161 (6)	0.0149 (6)	0.0169 (7)	-0.0019 (5)	0.0014 (5)	-0.0001 (5)
N14	0.0193 (8)	0.0104 (7)	0.0147 (8)	0.000	-0.0017 (6)	0.000
C15	0.0245 (8)	0.0144 (6)	0.0187 (7)	-0.0008 (6)	-0.0002 (6)	0.0028 (5)
C16	0.0271 (11)	0.0154 (9)	0.0134 (9)	0.000	-0.0003 (8)	0.000

Geometric parameters (\AA , $^\circ$)

C1—N14	1.4840 (17)	C9—H9B	0.9900
C1—C2	1.5197 (19)	C10—O11	1.4211 (17)
C1—C13	1.547 (2)	C10—H10A	0.9900
C1—H1	1.0000	C10—H10B	0.9900
C2—C3	1.393 (2)	O11—C10 ⁱ	1.4211 (17)
C2—C7	1.403 (2)	C12—O12	1.213 (3)
C3—C4	1.393 (2)	C12—C13 ⁱ	1.5168 (18)
C3—H3	0.9500	C12—C13	1.5168 (18)
C4—C5	1.385 (2)	C13—C15	1.524 (2)
C4—H4	0.9500	C13—H13	1.0000
C5—C6	1.388 (2)	N14—C16	1.467 (3)
C5—H5	0.9500	N14—C1 ⁱ	1.4840 (17)
C6—C7	1.397 (2)	C15—H15A	0.9800
C6—H6	0.9500	C15—H15B	0.9800
C7—O8	1.3666 (18)	C15—H15C	0.9800
O8—C9	1.4319 (18)	C16—H16A	0.9595
C9—C10	1.504 (2)	C16—H16B	0.9593
C9—H9A	0.9900		
N14—C1—C2	111.82 (12)	H9A—C9—H9B	108.6
N14—C1—C13	108.23 (12)	O11—C10—C9	107.80 (13)
C2—C1—C13	112.92 (12)	O11—C10—H10A	110.1
N14—C1—H1	107.9	C9—C10—H10A	110.1
C2—C1—H1	107.9	O11—C10—H10B	110.1
C13—C1—H1	107.9	C9—C10—H10B	110.1

C3—C2—C7	118.03 (13)	H10A—C10—H10B	108.5
C3—C2—C1	122.95 (13)	C10—O11—C10 ⁱ	112.69 (16)
C7—C2—C1	118.96 (13)	O12—C12—C13 ⁱ	122.11 (9)
C2—C3—C4	121.51 (15)	O12—C12—C13	122.11 (9)
C2—C3—H3	119.2	C13 ⁱ —C12—C13	115.71 (18)
C4—C3—H3	119.2	C12—C13—C15	111.50 (13)
C5—C4—C3	119.40 (15)	C12—C13—C1	106.80 (12)
C5—C4—H4	120.3	C15—C13—C1	113.36 (12)
C3—C4—H4	120.3	C12—C13—H13	108.3
C4—C5—C6	120.68 (14)	C15—C13—H13	108.3
C4—C5—H5	119.7	C1—C13—H13	108.3
C6—C5—H5	119.7	C16—N14—C1 ⁱ	113.80 (10)
C5—C6—C7	119.40 (14)	C16—N14—C1	113.80 (10)
C5—C6—H6	120.3	C1 ⁱ —N14—C1	108.27 (15)
C7—C6—H6	120.3	C13—C15—H15A	109.5
O8—C7—C6	123.03 (14)	C13—C15—H15B	109.5
O8—C7—C2	116.00 (12)	H15A—C15—H15B	109.5
C6—C7—C2	120.97 (14)	C13—C15—H15C	109.5
C7—O8—C9	118.82 (11)	H15A—C15—H15C	109.5
O8—C9—C10	106.99 (12)	H15B—C15—H15C	109.5
O8—C9—H9A	110.3	N14—C16—H16A	109.5
C10—C9—H9A	110.3	N14—C16—H16B	109.4
O8—C9—H9B	110.3	H16A—C16—H16B	109.5
C10—C9—H9B	110.3		
N14—C1—C2—C3	80.51 (18)	C2—C7—O8—C9	159.44 (13)
C13—C1—C2—C3	−41.84 (19)	C7—O8—C9—C10	−161.82 (12)
N14—C1—C2—C7	−96.59 (16)	O8—C9—C10—O11	67.31 (16)
C13—C1—C2—C7	141.06 (13)	C9—C10—O11—C10 ⁱ	−171.15 (10)
C7—C2—C3—C4	0.1 (2)	O12—C12—C13—C15	−1.4 (2)
C1—C2—C3—C4	−177.05 (14)	C13 ⁱ —C12—C13—C15	−178.58 (11)
C2—C3—C4—C5	0.3 (2)	O12—C12—C13—C1	122.9 (2)
C3—C4—C5—C6	−0.1 (2)	C13 ⁱ —C12—C13—C1	−54.2 (2)
C4—C5—C6—C7	−0.5 (2)	N14—C1—C13—C12	58.80 (16)
C5—C6—C7—O8	−179.15 (14)	C2—C1—C13—C12	−176.87 (13)
C5—C6—C7—C2	0.9 (2)	N14—C1—C13—C15	−178.02 (12)
C3—C2—C7—O8	179.34 (13)	C2—C1—C13—C15	−53.68 (16)
C1—C2—C7—O8	−3.41 (19)	C2—C1—N14—C16	38.57 (19)
C3—C2—C7—C6	−0.7 (2)	C13—C1—N14—C16	163.56 (14)
C1—C2—C7—C6	176.59 (13)	C2—C1—N14—C1 ⁱ	166.15 (9)
C6—C7—O8—C9	−20.6 (2)	C13—C1—N14—C1 ⁱ	−68.85 (18)

Symmetry code: (i) $x, -y+1/2, z$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C10—H10B ⁱⁱ —O12 ⁱⁱ	0.99	2.58	3.449 (2)	147

supporting information

C16—H16B···O11	0.96	2.57	3.530 (3)	180
----------------	------	------	-----------	-----

Symmetry code: (ii) $x-1/2, -y+1/2, -z-1/2$.