

Expected and unexpected products of reactions of 2-hydrazinylbenzothiazole with 3-nitrobenzenesulfonyl chloride in different solvents

Alexandra Morscher,^a Marcus V. N. de Souza,^b James L. Wardell^{b,a} and William T. A. Harrison^{a*}

Received 6 April 2018

Accepted 10 April 2018

Edited by J. Simpson, University of Otago, New Zealand

Keywords: crystal structure; hydrazinylbenzothiazole; hydrogen bonds; molecular salt.

CCDC references: 1835988; 1835987; 1835986

Supporting information: this article has supporting information at journals.iucr.org/e


^aDepartment of Chemistry, University of Aberdeen, Meston Walk, Aberdeen AB24 3UE, Scotland, and ^bInstituto de Tecnologia em Fármacos – Farmanguinhos, Fiocruz. R. Sizenando, Nabuco, 100, Mangunhos, 21041-250, Rio de Janeiro, RJ, Brazil. *Correspondence e-mail: w.harrison@abdn.ac.uk


The syntheses and crystal structures of 2-[2-(propan-2-ylidene)hydrazinyl]-1,3-benzothiazol-3-ium 3-nitrobenzenesulfonate ($C_{10}H_{12}N_2S^+ \cdot C_6H_4NO_5S^-$), (I), 2-[2-(3-nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazole ($C_{13}H_{10}N_4O_4S_2$), (II) and 2-[2-(3-nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazol-3-ium 3-nitrobenzenesulfonate ($C_{13}H_{11}N_4O_4S_2^+ \cdot C_6H_4NO_5S^-$), (III) are reported. Salt (I) arose from an unexpected reaction of 2-hydrazinylbenzothiazole with the acetone solvent in the presence of 3-nitrobenzenesulfonyl chloride, whereas (II) and (III) were recovered from the equivalent reaction carried out in methanol. The crystal of (I) features ion pairs linked by pairs of $N-H \cdots O_s$ ($s = \text{sulfonate}$) hydrogen bonds; adjacent cations interact by way of short $\pi-\pi$ stacking interactions between the thiazole rings [centroid-centroid separation = 3.4274 (18) Å]. In (II), which crystallizes with two neutral molecules in the asymmetric unit, the molecules are linked by $N-H \cdots N$ and $N-H \cdots O_n$ ($n = \text{nitro}$) hydrogen bonds to generate $[\bar{1}1\bar{1}]$ chains, which are cross-linked by $C-H \cdots O$ and $\pi-\pi$ stacking interactions. The crystal of (III) features centrosymmetric tetramers (two cations and two anions) linked by cooperative $N-H \cdots O$ hydrogen bonds; $C-H \cdots O$ and $\pi-\pi$ interactions occur between tetramers.

1. Chemical context

Heteroaromatic benzothiazole derivatives are well-studied compounds, due in the main to their various and useful biological activities (for a review, see Gulati *et al.*, 2017), but also to their fluorescent and optical properties (*e.g.* Liu *et al.*, 2018). Hydrazonyl derivatives, 2-Ar-CH=N-NH-benzothiazoles, formed from 2-hydrazinylbenzothiazole and ArCHO have attracted attention: for example, Katava *et al.* (2017) have reported antitumor activities and Behera & Manivannan (2017) studied their use as sensors. Less attention has been paid generally to 2-(ArSO₂NHNH)-benzothiazoles, although antimicrobial activities have been briefly reported (Rao *et al.*, 2005; Hipparagi *et al.*, 2007).

We have initiated a study of the syntheses, structures and biological activities of 2-(ArSO₂NHNH)-benzothiazoles and we now describe the structures of three products of the reactions of 2-hydrazinylbenzothiazole with 3-nitrobenzenesulfonyl chloride in different solvents, *viz.* 2-[2-(propan-2-ylidene)hydrazinyl]-1,3-benzothiazol-3-ium 3-nitrobenzenesulfonate (I), 2-[2-(3-nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazole (II) and 2-[2-(3-nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazol-3-ium 3-nitrobenzenesulfonate (III).





2. Structural commentary

Compound (I) crystallizes in space group $P\bar{1}$ with one $C_{10}H_{12}N_3S^+$ cation (protonated at N1) and one $C_6H_4NO_5S^-$ sulfonate anion in the asymmetric unit (Fig. 1). Evidently, the starting hydrazone has reacted with the acetone solvent (Day & Whiting, 1970) to generate an *N*-propylidene group; at the same time, the sulfonyl chloride has been hydrolysed to sulfonic acid and a molecular salt has crystallized after proton transfer from the sulfonic acid to the N atom of the thiazole ring. The cation is close to planar; the dihedral angle between the benzothiazole ring system (r.m.s. deviation = 0.005 Å) and the N2/N3/C8/C9/C10 grouping (r.m.s. deviation = 0.004 Å) is 7.89 (10)°; the C7–N2–N3–C8 torsion angle is –172.8 (2)°.




Figure 1
The asymmetric unit of (I) showing 50% displacement ellipsoids. Hydrogen bonds are indicated by double-dashed lines.




Figure 2
The asymmetric unit of (II) showing 50% displacement ellipsoids. Hydrogen bonds are indicated by double-dashed lines.

The C8–N3 bond length of 1.278 (4) Å is fully consistent with double-bond character. In the anion, the nitro group is twisted by 26.7 (4)° with respect to the benzene ring. As expected, the S–O bond lengths of the sulfonate group are almost the same, indicating the usual delocalization of the negative charge and the same situation is found in compound (III) described below.

Compound (II) represents the expected condensation product of the starting hydrazone and sulfonyl chloride and crystallizes with two neutral $C_{13}H_{10}N_4O_4S$ molecules in the asymmetric unit (Fig. 2) in space group $P\bar{1}$. In the first (S1) molecule, the dihedral angle between the benzothiazole ring system (r.m.s. deviation = 0.013 Å) and the C8 benzene ring is 32.59 (4)°; the nitro group is twisted by 0.68 (7)° from the C8 benzene ring. The C7–N2–N3–S2 torsion angle is –99.88 (12) and the H2–N2–N3–H3 torsion angle is –54 (2)°. The bond-angle sum at N2 is 359.9°, indicative of sp^2 hybridization, whereas the corresponding value for N3 of 341.1° points towards substantial sp^3 hybrid character. The C7–N2 bond length of 1.3529 (16) Å is short for a nominal single bond, presumably indicative of conjugation of the N2 nominal lone pair of electrons with the adjacent ring system. In the second (S3) molecule, the corresponding geometrical




Figure 3
The asymmetric unit of (III) showing 50% displacement ellipsoids. Hydrogen bonds are indicated by double-dashed lines.

data are 0.008 Å (r.m.s. deviation for S3 ring system), 30.01 (3)° (S3/C21 rings), 3.46 (13)° (nitro group and C21 ring), -103.53 (12)° (C20–N6–N7–S4), -50.3 (18)° (H6–N6–N7–H7), 359.9 ° (bond-angle sum at N6), 341.7 ° (bond-angle sum at N7) and 1.3549 (16) Å (C20–N6 bond length). All-in-all, the S1 and S3 molecules have similar conformations as indicated by the r.m.s. overlay fit of 0.221 Å for their non-hydrogen atoms.

Compound (III), which was recovered from the same reaction as (II), represents the same condensation product, which has gone on to further react with a hydrolysed sulfonyl chloride species to form a molecular salt (proton transfer to N1). Once again, the space group is $P\bar{1}$ and one cation and one anion (Fig. 3) make up the asymmetric unit. The benzothiazole ring system (r.m.s. deviation = 0.005 Å) subtends a dihedral angle of 57.54 (3)° with the C8 benzene ring and the nitro group is twisted from its attached ring by 4.8 (3)°. The C7–N2–N3–S2 and H2–N2–N3–H3 torsion angles are -110.54 (12) and -48.5 (19)°, respectively. The bond-angle sums at N2 and N3 are 359.0 and 339.1°, respectively, and the same conclusions *re* hybridization states for these atoms as in (II) may be drawn. This is backed up by the shortened C7–N2 bond length of 1.3317 (17) Å in (III) compared to (II), presumably because resonance is enhanced by the positive charge on N1. In the anion, the nitro group is twisted from its attached ring by 17.7 (2)°.

3. Supramolecular features

In the crystal of (I), the cation and the anion are linked by a pair of N–H···O hydrogen bonds (Table 1), which generate an $R_2^2(8)$ loop. The ion pairs are connected by various weak C–H···O interactions, with the acceptor O atoms being parts of the sulfonate and nitro groups. No C–H··· π interactions could be identified in the crystal of (I) but aromatic π – π stacking interactions are seen, with the shortest centroid–centroid separation of 3.4274 (18) Å (slippage = 0.729 Å) occurring between inversion-related pairs of thiazole rings (Fig. 4); atom N2 of the hydrazone group lies above the benzene ring ($Cg\cdots N2 = 3.385$ Å) and possibly provides some additional stabilization. Taken together, the directional inter-




Figure 4
Detail of the extended structure of (I) showing π – π stacking between inversion-related thiazole rings and possible secondary N··· π interactions. Symmetry code: (i) $-x, 1 - y, 1 - z$.

Table 1
Hydrogen-bond geometry (Å, °) for (I).

D–H···A	D–H	H···A	D···A	D–H···A
N1–H1···O1	0.86 (4)	1.87 (4)	2.721 (3)	167 (3)
N2–H2···O2	0.82 (4)	1.96 (4)	2.773 (3)	169 (3)
C3–H3···O2 ⁱ	0.95	2.51	3.415 (4)	160
C4–H4···O4 ⁱⁱ	0.95	2.55	3.292 (4)	135
C10–H10B···O4 ⁱⁱⁱ	0.98	2.54	3.297 (5)	134
C15–H15···O3 ^{iv}	0.95	2.50	3.315 (4)	144

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, -y + 1, -z$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x + 1, y, z$.

Table 2
Hydrogen-bond geometry (Å, °) for (II).

D–H···A	D–H	H···A	D···A	D–H···A
N2–H2···N5	0.816 (18)	2.033 (18)	2.8447 (15)	172.7 (16)
N3–H3···O3 ⁱ	0.848 (17)	2.129 (18)	2.9427 (15)	160.8 (15)
N6–H6···N1	0.820 (18)	2.050 (18)	2.8601 (15)	169.2 (17)
N7–H7···O7 ⁱⁱ	0.871 (17)	2.123 (18)	2.9472 (15)	157.6 (15)
C15–H15···O3 ⁱ	0.95	2.65	3.4888 (17)	147
C26–H26···O2	0.95	2.44	3.1774 (16)	134
C5–H5···O1 ⁱⁱⁱ	0.95	2.66	3.3218 (16)	127
C13–H13···O6	0.95	2.56	3.2731 (16)	133

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x + 1, -y, -z + 1$; (iii) $-x, -y + 1, -z + 1$.

Table 3
Hydrogen-bond geometry (Å, °) for (III).


D–H···A	D–H	H···A	D···A	D–H···A
N1–H1···O6	0.841 (19)	1.888 (19)	2.7267 (15)	175.2 (17)
N2–H2···O5	0.826 (19)	1.92 (2)	2.7489 (16)	175.4 (18)
N3–H3···O7 ⁱ	0.869 (18)	1.968 (19)	2.8058 (16)	161.6 (16)
C2–H2A···O7 ⁱⁱ	0.95	2.55	3.2510 (18)	130
C9–H9···O8 ⁱⁱⁱ	0.95	2.58	3.487 (2)	161

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $x + 1, y, z$; (iii) $-x + 1, -y, -z + 1$.

molecular interactions in (I) generate a three-dimensional network.

The dominant intermolecular interactions in (II) are N–H···N and N–H···O hydrogen bonds (Table 2); the first of these (N2–H2···N5 and N6–H6···N1) occur in the arbitrarily chosen asymmetric unit to link the molecules into dimers that 'slot together': the dihedral angle between the benzothiazole planes in the two molecules is 36.06 (4)° and the pendant benzene sulfonyl groups project to the same side of the ensemble. The N–H···O_n ($n = \text{nitro}$) links connect the dimers into infinite $[\bar{1}1\bar{1}]$ chains (Fig. 5). A number of weak C–H···O interactions are also observed, which serve to cross-link the chains. Several π – π stacking contacts occur in the crystal of (II), with the shortest [centroid–centroid separation = 3.5186 (7) Å] occurring between the C8–C13 and C14–C19 rings. Finally, a short N8–O7··· π ($\pi = \text{centroid of the C21–C26 benzene ring}$) contact is observed with N··· $\pi = 3.2497$ (12) Å and N–O··· $\pi = 86.24$ (8)°.

The packing in (III) features a pair of cation-to-anion N–H···O links from N1 and N2 (Table 3), which is essentially the same motif as seen in (I). The N3–H3 grouping links to a symmetry-generated sulfonate O atom and a centrosymmetric


Figure 5

Fragment of a $[1\bar{1}\bar{1}]$ hydrogen-bonded chain in (II) with the N—H...N and N—H...O bonds shown as blue and red double-dashed lines, respectively. All C-bonded hydrogen atoms have been omitted for clarity. Symmetry codes: (i) $-x, 1 - y, -z$; (ii) $1 - x, -y, 1 - z$.

tetramer (two cations and two anions) results (Fig. 6). A pair of weak C—H...O interactions helps to provide cohesion between tetramers in the crystal and π – π stacking is also observed, with the shortest centroid–centroid separation being 3.6743 (8) Å between the thiazole and C1–C6 rings.

4. Database survey


A survey of the Cambridge Structural Database (Groom *et al.*, 2016; updated to March 2018) for benzothiazole hydrazones revealed the prototype compound, benzothiazol-2-ylhydrazine (refcode: NAPKAR; Rajnikant *et al.*, 2005) as well

as three derivatives with various substituents attached to the benzene ring, *viz.* EVARID (Liu *et al.*, 2011), LAPCAI (Fun *et al.*, 2012*a*) and LEFTEX (Fun *et al.*, 2012*b*). No hits for benzenesulfonylhydrazino-benzothiazoles were recorded.

5. Synthesis and crystallization

To prepare (I), a mixture of 2-hydrazinylbenzothiazole (1.00 mmol) and 3-nitrobenzenesulfonyl chloride (1.00 mmol) in acetone (15 ml) was gently heated at 313–323 K for 30 minutes, then rotary evaporated and the residue was recrystallized by slow evaporation from methanol solution at room temperature; m.p. 444–445 K. ESI–HRMS ($M + H$). Calculated: 206.0752 for $C_{10}H_{12}N_3S$, found: 206.0755. IR: 2930(*br*), 1621, 1530, 1350, 1241, 1151, 1028 cm^{-1}

Compounds (II) and (III) arose from the same reaction: a solution of 2-hydrazinylbenzothiazole (1.00 mmol) and 3-nitrobenzenesulfonyl chloride (1.00 mmol) in methanol (15 ml) was gently heated at 313–323 K for 30 minutes, then rotary evaporated and the residue was recrystallized by slow evaporation from methanol solution at room temperature. A mixture of two distinct crystalline products, one yellow [compound (II)] and the other colourless [compound (III)], was isolated. These were separated by eye, and each product was further recrystallized from methanol solution. (II); m.p. 442–444 K. ESI–HRMS ($M - H$). Calculated: 349.0222 for $C_{13}H_9N_4O_2S_2$; found: 351.0220 ESI–HRMS ($M + H$). Calculated: 351.0065 for $C_{13}H_{11}N_4O_2S_2$; found: 349.0062 IR; 2989 (*br*), 1531, 1457, 1341, 1306, 1167 cm^{-1} . (III); m.p. 463–466 K. ESI–HRMS ($M + H$). Calculated: 351.0065 for $C_{13}H_{11}N_4O_2S_2$; found: 349.0065 IR: 2870 (*br*), 1553, 1436, 1363, 1241, 1127, 1065 cm^{-1} .


Figure 6

An inversion-generated tetramer in the crystal of (III). Symmetry code: (i) $1 - x, 1 - y, -z$.

Table 4
Experimental details.

	(I)	(II)	(III)
Crystal data			
Chemical formula	C ₁₀ H ₁₂ N ₃ S ⁺ ·C ₆ H ₄ NO ₅ S ⁻	C ₁₃ H ₁₀ N ₄ O ₄ S ₂	C ₁₃ H ₁₁ N ₄ O ₄ S ₂ ⁺ ·C ₆ H ₄ NO ₅ S ⁻
<i>M_r</i>	408.45	350.37	553.54
Crystal system, space group	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$	Triclinic, <i>P</i> $\bar{1}$
Temperature (K)	100	100	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	7.5308 (4), 10.9167 (7), 12.4438 (8)	6.83537 (15), 13.5788 (3), 15.6907 (4)	10.0399 (5), 10.7585 (4), 11.3372 (6)
α , β , γ (°)	66.058 (6), 79.034 (5), 72.156 (5)	99.382 (2), 98.2324 (19), 91.9841 (19)	85.607 (4), 71.369 (5), 77.115 (4)
<i>V</i> (Å ³)	887.47 (10)	1419.55 (6)	1131.16 (10)
<i>Z</i>	2	4	2
Radiation type	Mo <i>K</i> α	Mo <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.34	0.40	0.39
Crystal size (mm)	0.06 × 0.05 × 0.01	0.15 × 0.10 × 0.03	0.23 × 0.18 × 0.04
Data collection			
Diffractometer	Rigaku Mercury CCD	Rigaku Mercury CCD	Rigaku Mercury CCD
Absorption correction	Multi-scan (<i>FS_ABSCOR</i> ; Rigaku, 2013)	Multi-scan (<i>FS_ABSCOR</i> ; Rigaku, 2013)	Multi-scan (<i>FS_ABSCOR</i> ; Rigaku, 2013)
<i>T</i> _{min} , <i>T</i> _{max}	0.614, 1.000	0.861, 1.000	0.879, 1.000
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	11341, 3490, 3249	24032, 6465, 6033	19781, 5159, 4880
<i>R</i> _{int}	0.036	0.018	0.017
(sin θ / λ) _{max} (Å ⁻¹)	0.617	0.649	0.651
Refinement			
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.066, 0.184, 1.11	0.027, 0.072, 1.04	0.029, 0.076, 1.02
No. of reflections	3490	6465	5159
No. of parameters	250	427	334
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	1.32, -0.63	0.40, -0.38	0.39, -0.42

Computer programs: *CrystalClear* (Rigaku, 2012), *SHELXS97* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015), *ORTEP-3 for Windows* (Farrugia, 2012) and *pubCIF* (Westrip, 2010).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 4. The N-bound hydrogen atoms were located in difference maps and their positions freely refined. The C-bound hydrogen atoms were geometrically placed (C–H = 0.95–0.98 Å) and refined as riding atoms. The constraint $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$ or $1.5U_{\text{eq}}(\text{methyl carrier})$ was applied in all cases. The methyl groups in (I) were allowed to rotate, but not to tip, to best fit the electron density.

Acknowledgements

We thank the EPSRC National Crystallography Service (University of Southampton) for the X-ray data collections.

Funding information

MVNdS and JLW thank CNPq (Brazil) for financial support.

References

Behera, N. & Manivannan, V. (2017). *ChemistrySelect*, **2**, 11048–11054.
Day, A. C. & Whiting, M. C. (1970). *Org. Synth.* **50**, 3–5.

Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
Fun, H.-K., Ooi, C. W., Sarojini, B. K., Mohan, B. J. & Narayana, B. (2012a). *Acta Cryst.* **E68**, o691–o692.
Fun, H.-K., Quah, C. K., Sarojini, B. K., Mohan, B. J. & Narayana, B. (2012b). *Acta Cryst.* **E68**, o2459.
Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
Gulati, S., Wakode, S., Kaur, A. & Anand, K. (2017). *World J. Pharm. Pharm. Sci.* **6**, 1842–1869.
Hipparagi, S. M., Majumder, U. K., Parikshit, B., Geethavani, M. & Kumar, V. S. (2007). *Orient. J. Chem.* **23**, 65–66.
Katava, R., Pavelić, S. K., Harej, A., Hrenar, T. & Pavlović, G. (2017). *Struct. Chem.* **28**, 709–721.
Liu, C. T., Wang, F., Xiao, T., Chi, B., Wu, Y. H., Zhu, D.-R. & Chen, X. Q. (2018). *Sens. Actuators B Chem.* **256**, 55–62.
Liu, X.-F., Yu, X.-Y. & Jiang, S.-L. (2011). *Acta Cryst.* **E67**, o1641.
Rajnikant, Dinesh, Kamni, Deshmukh, M. B., Jagpat, S. S. & Desai, S. R. (2005). *J. Chem. Cryst.* **35**, 293–296.
Rao, D. S., Jayachandran, E., Sreenivasa, G. M. & Shivakumar, B. (2005). *Orient. J. Chem.* **21**, 113–116.
Rigaku (2012). *CrysAlis PRO*. Rigaku Corporation, Tokyo, Japan.
Rigaku (2013). *FS_ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2018). E74, 673-677 [https://doi.org/10.1107/S2056989018005595]

Expected and unexpected products of reactions of 2-hydrazinylbenzothiazole with 3-nitrobenzenesulfonyl chloride in different solvents

Alexandra Morscher, Marcus V. N. de Souza, James L. Wardell and William T. A. Harrison

Computing details

For all structures, data collection: *CrystalClear* (Rigaku, 2012); cell refinement: *CrystalClear* (Rigaku, 2012); data reduction: *CrystalClear* (Rigaku, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *publCIF* (Westrip, 2010).

2-[2-(Propan-2-ylidene)hydrazinyl]-1,3-benzothiazol-3-ium 3-nitrobenzenesulfonate (I)

Crystal data

$C_{10}H_{12}N_3S^+ \cdot C_6H_4NO_3S^-$

$M_r = 408.45$

Triclinic, $P\bar{1}$

$a = 7.5308$ (4) Å

$b = 10.9167$ (7) Å

$c = 12.4438$ (8) Å

$\alpha = 66.058$ (6)°

$\beta = 79.034$ (5)°

$\gamma = 72.156$ (5)°

$V = 887.47$ (10) Å³

$Z = 2$

$F(000) = 424$

$D_x = 1.528$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71075$ Å

Cell parameters from 6372 reflections

$\theta = 2.8\text{--}30.4^\circ$

$\mu = 0.34$ mm⁻¹

$T = 100$ K

Block, colourless

$0.06 \times 0.05 \times 0.01$ mm

Data collection

Rigaku Mercury CCD
diffractometer

ω scans

Absorption correction: multi-scan
(*FS_ABSCOR*; Rigaku, 2013)

$T_{\min} = 0.614$, $T_{\max} = 1.000$

11341 measured reflections

3490 independent reflections

3249 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -9 \rightarrow 7$

$k = -13 \rightarrow 13$

$l = -15 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.066$

$wR(F^2) = 0.184$

$S = 1.11$

3490 reflections

250 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.1206P)^2 + 0.8477P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.32$ e Å⁻³

$\Delta\rho_{\min} = -0.63$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2281 (4)	0.3162 (3)	0.5033 (3)	0.0203 (6)
C2	0.2122 (4)	0.1899 (3)	0.5087 (3)	0.0228 (6)
H2A	0.2284	0.1105	0.5798	0.027*
C3	0.1720 (4)	0.1837 (3)	0.4070 (3)	0.0262 (7)
H3	0.1611	0.0986	0.4083	0.031*
C4	0.1472 (4)	0.3006 (3)	0.3030 (3)	0.0248 (6)
H4	0.1204	0.2937	0.2344	0.030*
C5	0.1609 (4)	0.4267 (3)	0.2978 (3)	0.0231 (6)
H5	0.1419	0.5065	0.2270	0.028*
C6	0.2032 (4)	0.4330 (3)	0.3988 (3)	0.0191 (6)
C7	0.2606 (4)	0.5238 (3)	0.5207 (3)	0.0187 (6)
C8	0.3338 (4)	0.6563 (3)	0.7133 (3)	0.0228 (6)
C9	0.2989 (5)	0.8099 (3)	0.6481 (3)	0.0276 (7)
H9A	0.3122	0.8531	0.7004	0.041*
H9B	0.3897	0.8293	0.5795	0.041*
H9C	0.1719	0.8476	0.6217	0.041*
C10	0.3777 (5)	0.6021 (4)	0.8395 (3)	0.0298 (7)
H10A	0.3780	0.6793	0.8606	0.045*
H10B	0.2827	0.5556	0.8906	0.045*
H10C	0.5011	0.5359	0.8499	0.045*
N1	0.2209 (3)	0.5486 (3)	0.4133 (2)	0.0190 (5)
H1	0.188 (5)	0.633 (4)	0.364 (3)	0.023*
N2	0.2853 (3)	0.6164 (3)	0.5555 (2)	0.0197 (5)
H2	0.272 (5)	0.697 (4)	0.509 (3)	0.024*
N3	0.3282 (3)	0.5676 (3)	0.6723 (2)	0.0221 (5)
S1	0.27708 (10)	0.35384 (7)	0.61732 (6)	0.0196 (2)
C11	0.4451 (4)	0.9109 (3)	0.1794 (2)	0.0183 (6)
C12	0.5278 (4)	0.8235 (3)	0.1184 (2)	0.0195 (6)
H12	0.4602	0.7699	0.1067	0.023*
C13	0.7139 (4)	0.8170 (3)	0.0747 (3)	0.0223 (6)
C14	0.8145 (4)	0.8949 (3)	0.0890 (3)	0.0262 (7)
H14	0.9422	0.8861	0.0602	0.031*
C15	0.7274 (4)	0.9858 (3)	0.1455 (3)	0.0259 (7)
H15	0.7938	1.0430	0.1529	0.031*
C16	0.5418 (4)	0.9941 (3)	0.1918 (3)	0.0221 (6)
H16	0.4820	1.0560	0.2315	0.027*
S2	0.21789 (9)	0.91074 (7)	0.25157 (6)	0.0184 (2)
N4	0.8044 (4)	0.7214 (3)	0.0138 (2)	0.0302 (6)
O1	0.1647 (3)	0.8018 (2)	0.23498 (18)	0.0222 (5)

O2	0.2432 (3)	0.8753 (2)	0.37542 (18)	0.0238 (5)
O3	0.1002 (3)	1.0479 (2)	0.19792 (19)	0.0255 (5)
O4	0.7054 (4)	0.6912 (3)	-0.0342 (2)	0.0360 (6)
O5	0.9750 (3)	0.6763 (3)	0.0148 (2)	0.0432 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0162 (13)	0.0222 (14)	0.0203 (14)	-0.0039 (11)	0.0017 (11)	-0.0077 (11)
C2	0.0213 (14)	0.0197 (14)	0.0270 (15)	-0.0063 (11)	0.0015 (12)	-0.0089 (12)
C3	0.0203 (14)	0.0263 (15)	0.0358 (17)	-0.0088 (12)	0.0037 (13)	-0.0158 (14)
C4	0.0183 (14)	0.0342 (16)	0.0259 (15)	-0.0076 (12)	0.0028 (12)	-0.0164 (13)
C5	0.0171 (13)	0.0290 (15)	0.0204 (14)	-0.0085 (12)	0.0039 (11)	-0.0069 (12)
C6	0.0122 (12)	0.0201 (13)	0.0227 (14)	-0.0056 (10)	0.0048 (11)	-0.0073 (11)
C7	0.0099 (12)	0.0218 (14)	0.0226 (14)	-0.0058 (10)	0.0047 (10)	-0.0078 (11)
C8	0.0135 (13)	0.0302 (16)	0.0254 (15)	-0.0100 (12)	0.0056 (11)	-0.0107 (13)
C9	0.0294 (16)	0.0281 (16)	0.0300 (16)	-0.0103 (13)	-0.0003 (13)	-0.0140 (13)
C10	0.0295 (16)	0.0390 (18)	0.0248 (16)	-0.0160 (14)	0.0026 (13)	-0.0125 (14)
N1	0.0162 (11)	0.0178 (12)	0.0196 (12)	-0.0062 (9)	0.0028 (9)	-0.0042 (10)
N2	0.0188 (12)	0.0178 (12)	0.0214 (12)	-0.0070 (10)	0.0009 (10)	-0.0056 (10)
N3	0.0178 (12)	0.0264 (13)	0.0220 (12)	-0.0088 (10)	0.0023 (10)	-0.0084 (10)
S1	0.0197 (4)	0.0179 (4)	0.0193 (4)	-0.0062 (3)	-0.0008 (3)	-0.0044 (3)
C11	0.0153 (13)	0.0199 (13)	0.0170 (13)	-0.0062 (11)	-0.0021 (10)	-0.0026 (11)
C12	0.0202 (14)	0.0197 (13)	0.0168 (13)	-0.0071 (11)	-0.0021 (11)	-0.0034 (11)
C13	0.0201 (14)	0.0241 (14)	0.0174 (13)	-0.0032 (12)	-0.0003 (11)	-0.0045 (11)
C14	0.0156 (13)	0.0353 (17)	0.0207 (14)	-0.0068 (12)	-0.0018 (11)	-0.0029 (12)
C15	0.0177 (14)	0.0363 (17)	0.0250 (15)	-0.0141 (13)	-0.0015 (12)	-0.0077 (13)
C16	0.0217 (14)	0.0231 (14)	0.0213 (14)	-0.0066 (12)	-0.0024 (11)	-0.0071 (12)
S2	0.0147 (4)	0.0195 (4)	0.0200 (4)	-0.0077 (3)	0.0015 (3)	-0.0051 (3)
N4	0.0333 (15)	0.0265 (14)	0.0196 (13)	-0.0047 (12)	0.0065 (11)	-0.0033 (11)
O1	0.0214 (10)	0.0224 (10)	0.0223 (10)	-0.0098 (8)	-0.0013 (8)	-0.0049 (8)
O2	0.0250 (11)	0.0247 (11)	0.0221 (11)	-0.0119 (9)	0.0029 (8)	-0.0071 (8)
O3	0.0174 (10)	0.0240 (11)	0.0305 (12)	-0.0078 (9)	0.0016 (9)	-0.0052 (9)
O4	0.0485 (15)	0.0343 (13)	0.0270 (12)	-0.0169 (11)	0.0107 (11)	-0.0144 (10)
O5	0.0280 (13)	0.0419 (14)	0.0406 (15)	0.0041 (11)	0.0079 (11)	-0.0105 (12)

Geometric parameters (Å, °)

C1—C2	1.393 (4)	C10—H10B	0.9800
C1—C6	1.397 (4)	C10—H10C	0.9800
C1—S1	1.757 (3)	N1—H1	0.86 (4)
C2—C3	1.388 (4)	N2—N3	1.395 (3)
C2—H2A	0.9500	N2—H2	0.82 (4)
C3—C4	1.394 (5)	C11—C12	1.386 (4)
C3—H3	0.9500	C11—C16	1.393 (4)
C4—C5	1.387 (4)	C11—S2	1.775 (3)
C4—H4	0.9500	C12—C13	1.395 (4)
C5—C6	1.386 (4)	C12—H12	0.9500

C5—H5	0.9500	C13—C14	1.377 (5)
C6—N1	1.393 (4)	C13—N4	1.462 (4)
C7—N2	1.321 (4)	C14—C15	1.379 (5)
C7—N1	1.327 (4)	C14—H14	0.9500
C7—S1	1.734 (3)	C15—C16	1.398 (4)
C8—N3	1.278 (4)	C15—H15	0.9500
C8—C9	1.499 (4)	C16—H16	0.9500
C8—C10	1.500 (4)	S2—O3	1.440 (2)
C9—H9A	0.9800	S2—O1	1.463 (2)
C9—H9B	0.9800	S2—O2	1.464 (2)
C9—H9C	0.9800	N4—O4	1.226 (4)
C10—H10A	0.9800	N4—O5	1.227 (4)
C2—C1—C6	120.9 (3)	C7—N1—C6	114.0 (2)
C2—C1—S1	127.7 (2)	C7—N1—H1	120 (2)
C6—C1—S1	111.4 (2)	C6—N1—H1	125 (2)
C3—C2—C1	117.9 (3)	C7—N2—N3	115.8 (2)
C3—C2—H2A	121.0	C7—N2—H2	119 (2)
C1—C2—H2A	121.0	N3—N2—H2	125 (2)
C2—C3—C4	120.9 (3)	C8—N3—N2	117.8 (3)
C2—C3—H3	119.6	C7—S1—C1	89.17 (14)
C4—C3—H3	119.6	C12—C11—C16	121.1 (3)
C5—C4—C3	121.3 (3)	C12—C11—S2	120.3 (2)
C5—C4—H4	119.4	C16—C11—S2	118.5 (2)
C3—C4—H4	119.4	C11—C12—C13	117.6 (3)
C6—C5—C4	118.0 (3)	C11—C12—H12	121.2
C6—C5—H5	121.0	C13—C12—H12	121.2
C4—C5—H5	121.0	C14—C13—C12	122.4 (3)
C5—C6—N1	127.4 (3)	C14—C13—N4	119.5 (3)
C5—C6—C1	120.9 (3)	C12—C13—N4	118.0 (3)
N1—C6—C1	111.6 (3)	C13—C14—C15	119.1 (3)
N2—C7—N1	125.2 (3)	C13—C14—H14	120.4
N2—C7—S1	121.0 (2)	C15—C14—H14	120.4
N1—C7—S1	113.8 (2)	C14—C15—C16	120.1 (3)
N3—C8—C9	126.5 (3)	C14—C15—H15	119.9
N3—C8—C10	117.1 (3)	C16—C15—H15	119.9
C9—C8—C10	116.5 (3)	C11—C16—C15	119.5 (3)
C8—C9—H9A	109.5	C11—C16—H16	120.3
C8—C9—H9B	109.5	C15—C16—H16	120.3
H9A—C9—H9B	109.5	O3—S2—O1	114.29 (12)
C8—C9—H9C	109.5	O3—S2—O2	113.10 (13)
H9A—C9—H9C	109.5	O1—S2—O2	111.73 (12)
H9B—C9—H9C	109.5	O3—S2—C11	106.88 (13)
C8—C10—H10A	109.5	O1—S2—C11	105.44 (13)
C8—C10—H10B	109.5	O2—S2—C11	104.46 (12)
H10A—C10—H10B	109.5	O4—N4—O5	124.4 (3)
C8—C10—H10C	109.5	O4—N4—C13	118.1 (3)
H10A—C10—H10C	109.5	O5—N4—C13	117.5 (3)

H10B—C10—H10C	109.5		
C6—C1—C2—C3	0.4 (4)	C2—C1—S1—C7	-179.2 (3)
S1—C1—C2—C3	179.9 (2)	C6—C1—S1—C7	0.4 (2)
C1—C2—C3—C4	-0.4 (4)	C16—C11—C12—C13	3.0 (4)
C2—C3—C4—C5	-0.3 (4)	S2—C11—C12—C13	-173.0 (2)
C3—C4—C5—C6	1.0 (4)	C11—C12—C13—C14	-0.9 (4)
C4—C5—C6—N1	-179.0 (3)	C11—C12—C13—N4	177.9 (2)
C4—C5—C6—C1	-1.0 (4)	C12—C13—C14—C15	-1.9 (4)
C2—C1—C6—C5	0.4 (4)	N4—C13—C14—C15	179.3 (3)
S1—C1—C6—C5	-179.2 (2)	C13—C14—C15—C16	2.6 (4)
C2—C1—C6—N1	178.6 (2)	C12—C11—C16—C15	-2.2 (4)
S1—C1—C6—N1	-0.9 (3)	S2—C11—C16—C15	173.8 (2)
N2—C7—N1—C6	179.5 (2)	C14—C15—C16—C11	-0.6 (4)
S1—C7—N1—C6	-0.9 (3)	C12—C11—S2—O3	-118.7 (2)
C5—C6—N1—C7	179.3 (3)	C16—C11—S2—O3	65.2 (2)
C1—C6—N1—C7	1.2 (3)	C12—C11—S2—O1	3.3 (3)
N1—C7—N2—N3	-179.3 (2)	C16—C11—S2—O1	-172.8 (2)
S1—C7—N2—N3	1.2 (3)	C12—C11—S2—O2	121.2 (2)
C9—C8—N3—N2	-0.1 (4)	C16—C11—S2—O2	-54.9 (2)
C10—C8—N3—N2	179.2 (2)	C14—C13—N4—O4	-154.8 (3)
C7—N2—N3—C8	-172.8 (2)	C12—C13—N4—O4	26.4 (4)
N2—C7—S1—C1	179.9 (2)	C14—C13—N4—O5	25.7 (4)
N1—C7—S1—C1	0.3 (2)	C12—C13—N4—O5	-153.2 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O1	0.86 (4)	1.87 (4)	2.721 (3)	167 (3)
N2—H2 \cdots O2	0.82 (4)	1.96 (4)	2.773 (3)	169 (3)
C3—H3 \cdots O2 ⁱ	0.95	2.51	3.415 (4)	160
C4—H4 \cdots O4 ⁱⁱ	0.95	2.55	3.292 (4)	135
C10—H10B \cdots O4 ⁱⁱⁱ	0.98	2.54	3.297 (5)	134
C15—H15 \cdots O3 ^{iv}	0.95	2.50	3.315 (4)	144

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y+1, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x+1, y, z$.

2-[2-(3-Nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazole (II)

Crystal data

 $C_{13}H_{10}N_4O_4S_2$ $M_r = 350.37$ Triclinic, $P1$ $a = 6.83537$ (15) \AA $b = 13.5788$ (3) \AA $c = 15.6907$ (4) \AA $\alpha = 99.382$ (2) $^\circ$ $\beta = 98.2324$ (19) $^\circ$ $\gamma = 91.9841$ (19) $^\circ$ $V = 1419.55$ (6) \AA^3 $Z = 4$ $F(000) = 720$ $D_x = 1.639$ Mg m^{-3} Mo $K\alpha$ radiation, $\lambda = 0.71073$ \AA

Cell parameters from 18994 reflections

 $\theta = 2.7\text{--}27.5^\circ$ $\mu = 0.40$ mm^{-1} $T = 100$ K

Plate, yellow

 $0.15 \times 0.10 \times 0.03$ mm

*Data collection*Rigaku Mercury CCD
diffractometer ω scansAbsorption correction: multi-scan
(*FS_ABSCOR*; Rigaku, 2013) $T_{\min} = 0.861$, $T_{\max} = 1.000$

24032 measured reflections

6465 independent reflections

6033 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.018$ $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.2^\circ$ $h = -8 \rightarrow 7$ $k = -17 \rightarrow 17$ $l = -20 \rightarrow 19$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.027$ $wR(F^2) = 0.072$ $S = 1.04$

6465 reflections

427 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0369P)^2 + 0.771P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.38 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.43539 (18)	0.33184 (9)	0.47102 (8)	0.0141 (2)
C2	0.55062 (19)	0.27184 (10)	0.51978 (8)	0.0169 (2)
H3A	0.6190	0.2193	0.4915	0.020*
C3	0.5642 (2)	0.28988 (10)	0.61023 (9)	0.0202 (3)
H3B	0.6435	0.2497	0.6439	0.024*
C4	0.4627 (2)	0.36625 (11)	0.65225 (9)	0.0210 (3)
H4	0.4751	0.3779	0.7142	0.025*
C5	0.3439 (2)	0.42538 (10)	0.60476 (8)	0.0185 (3)
H5	0.2733	0.4767	0.6333	0.022*
C6	0.33125 (18)	0.40728 (9)	0.51429 (8)	0.0149 (2)
C7	0.28467 (18)	0.38771 (9)	0.35588 (8)	0.0139 (2)
C8	-0.12349 (18)	0.31751 (9)	0.11243 (8)	0.0140 (2)
C9	-0.13679 (18)	0.36196 (9)	0.03836 (8)	0.0147 (2)
H9	-0.1521	0.4317	0.0415	0.018*
C10	-0.12680 (18)	0.30065 (9)	-0.04025 (8)	0.0151 (2)
C11	-0.10555 (19)	0.19861 (10)	-0.04774 (9)	0.0177 (3)
H11	-0.1002	0.1586	-0.1029	0.021*
C12	-0.09235 (19)	0.15654 (10)	0.02733 (9)	0.0177 (3)
H12	-0.0772	0.0868	0.0238	0.021*
C13	-0.10106 (18)	0.21541 (9)	0.10793 (8)	0.0161 (2)
H13	-0.0918	0.1862	0.1593	0.019*

S1	0.19123 (5)	0.46708 (2)	0.43815 (2)	0.01478 (7)
S2	-0.14385 (4)	0.39379 (2)	0.21312 (2)	0.01401 (7)
N1	0.40625 (15)	0.32218 (8)	0.38016 (7)	0.0139 (2)
N2	0.23064 (17)	0.39617 (9)	0.27112 (7)	0.0172 (2)
H2	0.284 (3)	0.3656 (13)	0.2323 (11)	0.021*
N3	0.07391 (16)	0.45532 (8)	0.25127 (7)	0.0156 (2)
H3	0.101 (2)	0.4983 (13)	0.2204 (11)	0.019*
N4	-0.14030 (17)	0.34643 (8)	-0.11915 (7)	0.0180 (2)
O1	-0.27389 (14)	0.47084 (7)	0.19593 (6)	0.01904 (19)
O2	-0.17746 (14)	0.32943 (7)	0.27375 (6)	0.01830 (19)
O3	-0.15792 (16)	0.43763 (7)	-0.11068 (6)	0.0248 (2)
O4	-0.13227 (18)	0.29368 (8)	-0.18884 (6)	0.0306 (2)
C14	0.38706 (17)	0.27186 (9)	0.04845 (8)	0.0135 (2)
C15	0.37430 (18)	0.35578 (10)	0.00746 (8)	0.0162 (2)
H15	0.3734	0.4208	0.0407	0.019*
C16	0.36290 (19)	0.34242 (10)	-0.08280 (9)	0.0188 (3)
H16	0.3550	0.3990	-0.1114	0.023*
C17	0.3629 (2)	0.24711 (10)	-0.13217 (9)	0.0203 (3)
H17	0.3535	0.2398	-0.1939	0.024*
C18	0.3765 (2)	0.16275 (10)	-0.09252 (8)	0.0183 (3)
H18	0.3766	0.0979	-0.1261	0.022*
C19	0.38981 (18)	0.17643 (9)	-0.00197 (8)	0.0146 (2)
C20	0.40680 (18)	0.18334 (9)	0.15515 (8)	0.0139 (2)
C21	0.15315 (19)	0.07929 (9)	0.36629 (8)	0.0163 (2)
C22	0.22077 (19)	0.04559 (10)	0.44344 (8)	0.0165 (2)
H22	0.2967	-0.0117	0.4438	0.020*
C23	0.17295 (19)	0.09898 (10)	0.51990 (8)	0.0175 (3)
C24	0.0642 (2)	0.18347 (10)	0.52236 (9)	0.0205 (3)
H24	0.0348	0.2185	0.5761	0.025*
C25	-0.0005 (2)	0.21522 (10)	0.44401 (10)	0.0211 (3)
H25	-0.0747	0.2731	0.4439	0.025*
C26	0.0421 (2)	0.16336 (10)	0.36575 (9)	0.0190 (3)
H26	-0.0042	0.1850	0.3122	0.023*
S3	0.40564 (5)	0.08651 (2)	0.06631 (2)	0.01512 (7)
S4	0.21030 (5)	0.01172 (2)	0.26734 (2)	0.01740 (8)
N5	0.39616 (15)	0.27357 (8)	0.13792 (7)	0.0136 (2)
N6	0.41743 (18)	0.16301 (8)	0.23735 (7)	0.0184 (2)
H6	0.430 (2)	0.2085 (13)	0.2795 (12)	0.022*
N7	0.42029 (17)	0.06345 (8)	0.24827 (7)	0.0170 (2)
H7	0.520 (3)	0.0496 (12)	0.2847 (11)	0.020*
N8	0.23973 (17)	0.06263 (9)	0.60183 (7)	0.0205 (2)
O5	0.26067 (17)	-0.08604 (7)	0.28157 (6)	0.0248 (2)
O6	0.05911 (15)	0.02700 (8)	0.19880 (6)	0.0242 (2)
O7	0.32765 (15)	-0.01577 (8)	0.59683 (7)	0.0260 (2)
O8	0.20339 (17)	0.10970 (9)	0.67020 (7)	0.0312 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0141 (6)	0.0152 (6)	0.0123 (6)	-0.0026 (4)	0.0022 (4)	0.0005 (4)
C2	0.0156 (6)	0.0167 (6)	0.0180 (6)	0.0007 (5)	0.0022 (5)	0.0023 (5)
C3	0.0207 (6)	0.0218 (7)	0.0186 (6)	0.0001 (5)	0.0000 (5)	0.0076 (5)
C4	0.0248 (7)	0.0256 (7)	0.0121 (6)	-0.0011 (5)	0.0017 (5)	0.0035 (5)
C5	0.0211 (6)	0.0208 (6)	0.0131 (6)	0.0011 (5)	0.0044 (5)	0.0002 (5)
C6	0.0143 (6)	0.0166 (6)	0.0132 (6)	-0.0002 (4)	0.0011 (4)	0.0019 (5)
C7	0.0145 (6)	0.0142 (6)	0.0124 (6)	-0.0014 (4)	0.0039 (4)	-0.0006 (4)
C8	0.0135 (5)	0.0154 (6)	0.0120 (6)	0.0018 (4)	0.0007 (4)	0.0005 (4)
C9	0.0150 (6)	0.0146 (6)	0.0139 (6)	0.0004 (4)	0.0014 (4)	0.0016 (5)
C10	0.0142 (6)	0.0177 (6)	0.0130 (6)	-0.0006 (4)	0.0008 (4)	0.0025 (5)
C11	0.0160 (6)	0.0173 (6)	0.0175 (6)	-0.0001 (5)	0.0012 (5)	-0.0028 (5)
C12	0.0177 (6)	0.0131 (6)	0.0211 (6)	0.0012 (5)	0.0006 (5)	0.0011 (5)
C13	0.0152 (6)	0.0164 (6)	0.0166 (6)	0.0012 (5)	0.0001 (5)	0.0043 (5)
S1	0.01749 (15)	0.01571 (15)	0.01076 (14)	0.00315 (11)	0.00279 (11)	0.00015 (11)
S2	0.01745 (15)	0.01462 (14)	0.01033 (14)	0.00364 (11)	0.00230 (11)	0.00244 (11)
N1	0.0147 (5)	0.0151 (5)	0.0114 (5)	-0.0002 (4)	0.0026 (4)	0.0005 (4)
N2	0.0186 (5)	0.0224 (6)	0.0106 (5)	0.0069 (4)	0.0035 (4)	0.0004 (4)
N3	0.0193 (5)	0.0144 (5)	0.0130 (5)	0.0019 (4)	0.0015 (4)	0.0030 (4)
N4	0.0196 (5)	0.0212 (6)	0.0126 (5)	-0.0015 (4)	0.0015 (4)	0.0017 (4)
O1	0.0238 (5)	0.0196 (5)	0.0145 (4)	0.0093 (4)	0.0029 (4)	0.0032 (4)
O2	0.0222 (5)	0.0197 (5)	0.0144 (4)	0.0023 (4)	0.0039 (4)	0.0057 (4)
O3	0.0390 (6)	0.0192 (5)	0.0181 (5)	0.0015 (4)	0.0069 (4)	0.0060 (4)
O4	0.0487 (7)	0.0290 (6)	0.0116 (5)	0.0004 (5)	0.0035 (4)	-0.0022 (4)
C14	0.0113 (5)	0.0162 (6)	0.0131 (6)	0.0016 (4)	0.0024 (4)	0.0018 (5)
C15	0.0151 (6)	0.0158 (6)	0.0176 (6)	0.0016 (4)	0.0020 (5)	0.0033 (5)
C16	0.0187 (6)	0.0206 (6)	0.0186 (6)	0.0025 (5)	0.0018 (5)	0.0087 (5)
C17	0.0223 (7)	0.0265 (7)	0.0126 (6)	0.0026 (5)	0.0014 (5)	0.0050 (5)
C18	0.0226 (6)	0.0187 (6)	0.0136 (6)	0.0038 (5)	0.0031 (5)	0.0015 (5)
C19	0.0152 (6)	0.0151 (6)	0.0138 (6)	0.0025 (4)	0.0019 (4)	0.0036 (5)
C20	0.0145 (6)	0.0146 (6)	0.0121 (6)	0.0007 (4)	0.0030 (4)	0.0003 (4)
C21	0.0192 (6)	0.0151 (6)	0.0147 (6)	-0.0028 (5)	0.0037 (5)	0.0025 (5)
C22	0.0169 (6)	0.0154 (6)	0.0174 (6)	-0.0023 (5)	0.0026 (5)	0.0040 (5)
C23	0.0175 (6)	0.0196 (6)	0.0156 (6)	-0.0048 (5)	0.0026 (5)	0.0042 (5)
C24	0.0202 (6)	0.0195 (6)	0.0218 (7)	-0.0041 (5)	0.0082 (5)	0.0000 (5)
C25	0.0202 (6)	0.0161 (6)	0.0290 (7)	0.0001 (5)	0.0080 (5)	0.0058 (5)
C26	0.0197 (6)	0.0175 (6)	0.0212 (7)	-0.0008 (5)	0.0035 (5)	0.0078 (5)
S3	0.02245 (16)	0.01240 (14)	0.01072 (14)	0.00307 (11)	0.00390 (11)	0.00095 (11)
S4	0.02582 (17)	0.01316 (15)	0.01286 (15)	-0.00091 (12)	0.00193 (12)	0.00245 (11)
N5	0.0158 (5)	0.0131 (5)	0.0118 (5)	0.0008 (4)	0.0029 (4)	0.0009 (4)
N6	0.0332 (6)	0.0112 (5)	0.0108 (5)	-0.0002 (4)	0.0048 (4)	0.0009 (4)
N7	0.0238 (6)	0.0146 (5)	0.0135 (5)	0.0035 (4)	0.0034 (4)	0.0039 (4)
N8	0.0198 (5)	0.0257 (6)	0.0152 (5)	-0.0052 (5)	0.0021 (4)	0.0030 (4)
O5	0.0423 (6)	0.0128 (4)	0.0197 (5)	0.0008 (4)	0.0056 (4)	0.0035 (4)
O6	0.0283 (5)	0.0252 (5)	0.0171 (5)	-0.0024 (4)	-0.0020 (4)	0.0038 (4)
O7	0.0274 (5)	0.0324 (6)	0.0192 (5)	0.0041 (4)	0.0015 (4)	0.0090 (4)

O8	0.0406 (6)	0.0368 (6)	0.0148 (5)	-0.0036 (5)	0.0069 (4)	-0.0009 (4)
----	------------	------------	------------	-------------	------------	-------------

Geometric parameters (Å, °)

C1—C2	1.3944 (18)	C14—N5	1.3927 (16)
C1—N1	1.3947 (16)	C14—C15	1.3963 (17)
C1—C6	1.4068 (17)	C14—C19	1.4061 (17)
C2—C3	1.3894 (19)	C15—C16	1.3887 (18)
C2—H3A	0.9500	C15—H15	0.9500
C3—C4	1.396 (2)	C16—C17	1.3958 (19)
C3—H3B	0.9500	C16—H16	0.9500
C4—C5	1.3877 (19)	C17—C18	1.3895 (19)
C4—H4	0.9500	C17—H17	0.9500
C5—C6	1.3902 (17)	C18—C19	1.3918 (18)
C5—H5	0.9500	C18—H18	0.9500
C6—S1	1.7450 (13)	C19—S3	1.7473 (13)
C7—N1	1.3033 (16)	C20—N5	1.2992 (16)
C7—N2	1.3529 (16)	C20—N6	1.3549 (16)
C7—S1	1.7511 (12)	C20—S3	1.7522 (12)
C8—C9	1.3876 (17)	C21—C22	1.3869 (18)
C8—C13	1.3917 (17)	C21—C26	1.3934 (18)
C8—S2	1.7685 (12)	C21—S4	1.7717 (13)
C9—C10	1.3829 (17)	C22—C23	1.3846 (18)
C9—H9	0.9500	C22—H22	0.9500
C10—C11	1.3858 (18)	C23—C24	1.3868 (19)
C10—N4	1.4661 (16)	C23—N8	1.4693 (17)
C11—C12	1.3837 (19)	C24—C25	1.387 (2)
C11—H11	0.9500	C24—H24	0.9500
C12—C13	1.3918 (18)	C25—C26	1.387 (2)
C12—H12	0.9500	C25—H25	0.9500
C13—H13	0.9500	C26—H26	0.9500
S2—O2	1.4287 (9)	S4—O5	1.4271 (10)
S2—O1	1.4301 (9)	S4—O6	1.4272 (10)
S2—N3	1.6636 (11)	S4—N7	1.6628 (12)
N2—N3	1.3912 (15)	N6—N7	1.3906 (15)
N2—H2	0.816 (18)	N6—H6	0.820 (18)
N3—H3	0.848 (17)	N7—H7	0.871 (17)
N4—O4	1.2150 (15)	N8—O8	1.2186 (16)
N4—O3	1.2358 (15)	N8—O7	1.2374 (16)
C2—C1—N1	125.63 (11)	N5—C14—C15	125.14 (11)
C2—C1—C6	119.30 (11)	N5—C14—C19	115.21 (11)
N1—C1—C6	115.02 (11)	C15—C14—C19	119.65 (11)
C3—C2—C1	119.12 (12)	C16—C15—C14	118.75 (12)
C3—C2—H3A	120.4	C16—C15—H15	120.6
C1—C2—H3A	120.4	C14—C15—H15	120.6
C2—C3—C4	120.83 (12)	C15—C16—C17	120.99 (12)
C2—C3—H3B	119.6	C15—C16—H16	119.5

C4—C3—H3B	119.6	C17—C16—H16	119.5
C5—C4—C3	120.91 (12)	C18—C17—C16	121.09 (12)
C5—C4—H4	119.5	C18—C17—H17	119.5
C3—C4—H4	119.5	C16—C17—H17	119.5
C4—C5—C6	118.08 (12)	C17—C18—C19	117.81 (12)
C4—C5—H5	121.0	C17—C18—H18	121.1
C6—C5—H5	121.0	C19—C18—H18	121.1
C5—C6—C1	121.74 (12)	C18—C19—C14	121.70 (12)
C5—C6—S1	128.32 (10)	C18—C19—S3	128.77 (10)
C1—C6—S1	109.93 (9)	C14—C19—S3	109.51 (9)
N1—C7—N2	122.68 (11)	N5—C20—N6	122.64 (11)
N1—C7—S1	117.31 (9)	N5—C20—S3	116.93 (9)
N2—C7—S1	120.01 (10)	N6—C20—S3	120.43 (9)
C9—C8—C13	121.44 (11)	C22—C21—C26	121.43 (12)
C9—C8—S2	118.05 (9)	C22—C21—S4	118.12 (10)
C13—C8—S2	120.47 (10)	C26—C21—S4	120.45 (10)
C10—C9—C8	117.29 (11)	C23—C22—C21	117.16 (12)
C10—C9—H9	121.4	C23—C22—H22	121.4
C8—C9—H9	121.4	C21—C22—H22	121.4
C9—C10—C11	123.15 (12)	C22—C23—C24	123.27 (12)
C9—C10—N4	118.01 (11)	C22—C23—N8	117.66 (12)
C11—C10—N4	118.84 (11)	C24—C23—N8	119.07 (12)
C12—C11—C10	118.18 (12)	C25—C24—C23	118.01 (13)
C12—C11—H11	120.9	C25—C24—H24	121.0
C10—C11—H11	120.9	C23—C24—H24	121.0
C11—C12—C13	120.65 (12)	C26—C25—C24	120.68 (13)
C11—C12—H12	119.7	C26—C25—H25	119.7
C13—C12—H12	119.7	C24—C25—H25	119.7
C8—C13—C12	119.28 (12)	C25—C26—C21	119.45 (12)
C8—C13—H13	120.4	C25—C26—H26	120.3
C12—C13—H13	120.4	C21—C26—H26	120.3
C6—S1—C7	88.04 (6)	C19—S3—C20	88.30 (6)
O2—S2—O1	121.70 (6)	O5—S4—O6	121.88 (6)
O2—S2—N3	106.58 (6)	O5—S4—N7	103.87 (6)
O1—S2—N3	103.87 (6)	O6—S4—N7	106.93 (6)
O2—S2—C8	107.73 (6)	O5—S4—C21	108.35 (6)
O1—S2—C8	108.20 (6)	O6—S4—C21	107.54 (6)
N3—S2—C8	108.09 (6)	N7—S4—C21	107.49 (6)
C7—N1—C1	109.69 (10)	C20—N5—C14	110.05 (10)
C7—N2—N3	117.74 (11)	C20—N6—N7	118.05 (11)
C7—N2—H2	121.7 (12)	C20—N6—H6	120.6 (12)
N3—N2—H2	120.5 (12)	N7—N6—H6	121.2 (12)
N2—N3—S2	115.55 (9)	N6—N7—S4	116.35 (9)
N2—N3—H3	113.4 (11)	N6—N7—H7	114.9 (11)
S2—N3—H3	112.1 (11)	S4—N7—H7	110.4 (11)
O4—N4—O3	123.45 (12)	O8—N8—O7	123.72 (12)
O4—N4—C10	118.93 (11)	O8—N8—C23	118.84 (12)
O3—N4—C10	117.61 (10)	O7—N8—C23	117.43 (11)

N1—C1—C2—C3	-179.34 (12)	N5—C14—C15—C16	-178.92 (12)
C6—C1—C2—C3	-1.78 (18)	C19—C14—C15—C16	0.54 (18)
C1—C2—C3—C4	0.6 (2)	C14—C15—C16—C17	0.38 (19)
C2—C3—C4—C5	0.7 (2)	C15—C16—C17—C18	-0.7 (2)
C3—C4—C5—C6	-0.9 (2)	C16—C17—C18—C19	0.1 (2)
C4—C5—C6—C1	-0.28 (19)	C17—C18—C19—C14	0.88 (19)
C4—C5—C6—S1	178.53 (10)	C17—C18—C19—S3	179.20 (10)
C2—C1—C6—C5	1.64 (19)	N5—C14—C19—C18	178.32 (11)
N1—C1—C6—C5	179.45 (11)	C15—C14—C19—C18	-1.20 (19)
C2—C1—C6—S1	-177.37 (10)	N5—C14—C19—S3	-0.30 (14)
N1—C1—C6—S1	0.44 (14)	C15—C14—C19—S3	-179.81 (9)
C13—C8—C9—C10	-0.02 (18)	C26—C21—C22—C23	0.08 (19)
S2—C8—C9—C10	-178.02 (9)	S4—C21—C22—C23	-179.45 (9)
C8—C9—C10—C11	0.36 (19)	C21—C22—C23—C24	-0.62 (19)
C8—C9—C10—N4	-179.90 (11)	C21—C22—C23—N8	178.59 (11)
C9—C10—C11—C12	-0.48 (19)	C22—C23—C24—C25	0.4 (2)
N4—C10—C11—C12	179.79 (11)	N8—C23—C24—C25	-178.77 (11)
C10—C11—C12—C13	0.25 (19)	C23—C24—C25—C26	0.3 (2)
C9—C8—C13—C12	-0.19 (19)	C24—C25—C26—C21	-0.8 (2)
S2—C8—C13—C12	177.77 (10)	C22—C21—C26—C25	0.63 (19)
C11—C12—C13—C8	0.07 (19)	S4—C21—C26—C25	-179.85 (10)
C5—C6—S1—C7	-179.42 (13)	C18—C19—S3—C20	-178.41 (13)
C1—C6—S1—C7	-0.50 (9)	C14—C19—S3—C20	0.08 (9)
N1—C7—S1—C6	0.52 (10)	N5—C20—S3—C19	0.17 (10)
N2—C7—S1—C6	-179.24 (11)	N6—C20—S3—C19	179.86 (11)
C9—C8—S2—O2	165.20 (10)	C22—C21—S4—O5	19.55 (12)
C13—C8—S2—O2	-12.82 (12)	C26—C21—S4—O5	-159.98 (11)
C9—C8—S2—O1	31.90 (12)	C22—C21—S4—O6	153.05 (10)
C13—C8—S2—O1	-146.12 (10)	C26—C21—S4—O6	-26.48 (12)
C9—C8—S2—N3	-79.99 (11)	C22—C21—S4—N7	-92.13 (11)
C13—C8—S2—N3	101.99 (11)	C26—C21—S4—N7	88.34 (11)
N2—C7—N1—C1	179.39 (11)	N6—C20—N5—C14	179.96 (12)
S1—C7—N1—C1	-0.35 (13)	S3—C20—N5—C14	-0.36 (14)
C2—C1—N1—C7	177.58 (12)	C15—C14—N5—C20	179.91 (12)
C6—C1—N1—C7	-0.07 (15)	C19—C14—N5—C20	0.42 (15)
N1—C7—N2—N3	168.50 (11)	N5—C20—N6—N7	177.46 (11)
S1—C7—N2—N3	-11.75 (16)	S3—C20—N6—N7	-2.20 (17)
C7—N2—N3—S2	-99.88 (12)	C20—N6—N7—S4	-103.52 (12)
O2—S2—N3—N2	52.10 (10)	O5—S4—N7—N6	-176.11 (9)
O1—S2—N3—N2	-178.24 (9)	O6—S4—N7—N6	53.82 (10)
C8—S2—N3—N2	-63.47 (10)	C21—S4—N7—N6	-61.41 (10)
C9—C10—N4—O4	-179.83 (12)	C22—C23—N8—O8	178.31 (12)
C11—C10—N4—O4	-0.08 (18)	C24—C23—N8—O8	-2.44 (18)
C9—C10—N4—O3	0.70 (17)	C22—C23—N8—O7	-2.85 (17)
C11—C10—N4—O3	-179.55 (12)	C24—C23—N8—O7	176.39 (12)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2...N5	0.816 (18)	2.033 (18)	2.8447 (15)	172.7 (16)
N3—H3...O3 ⁱ	0.848 (17)	2.129 (18)	2.9427 (15)	160.8 (15)
N6—H6...N1	0.820 (18)	2.050 (18)	2.8601 (15)	169.2 (17)
N7—H7...O7 ⁱⁱ	0.871 (17)	2.123 (18)	2.9472 (15)	157.6 (15)
C15—H15...O3 ⁱ	0.95	2.65	3.4888 (17)	147
C26—H26...O2	0.95	2.44	3.1774 (16)	134
C5—H5...O1 ⁱⁱⁱ	0.95	2.66	3.3218 (16)	127
C13—H13...O6	0.95	2.56	3.2731 (16)	133

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+1, -y, -z+1$; (iii) $-x, -y+1, -z+1$.

2-[2-(3-Nitrobenzenesulfonyl)hydrazinyl]-1,3-benzothiazol-3-ium; 3-nitrobenzenesulfonate (III)

Crystal data

$C_{13}H_{11}N_4O_4S_2^+ \cdot C_6H_4NO_5S^-$

$M_r = 553.54$

Triclinic, $P\bar{1}$

$a = 10.0399$ (5) Å

$b = 10.7585$ (4) Å

$c = 11.3372$ (6) Å

$\alpha = 85.607$ (4)°

$\beta = 71.369$ (5)°

$\gamma = 77.115$ (4)°

$V = 1131.16$ (10) Å³

$Z = 2$

$F(000) = 568$

$D_x = 1.625$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 14254 reflections

$\theta = 2.0$ – 27.5 °

$\mu = 0.39$ mm⁻¹

$T = 100$ K

Plate, colourless

$0.23 \times 0.18 \times 0.04$ mm

Data collection

Rigaku Mercury CCD
diffractometer

ω scans

Absorption correction: multi-scan

(*FS_ABSCOR*; Rigaku, 2013)

$T_{\min} = 0.879$, $T_{\max} = 1.000$

19781 measured reflections

5159 independent reflections

4880 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 27.6$ °, $\theta_{\min} = 2.2$ °

$h = -13 \rightarrow 11$

$k = -14 \rightarrow 13$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.029$

$wR(F^2) = 0.076$

$S = 1.02$

5159 reflections

334 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0353P)^2 + 0.8265P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.39$ e Å⁻³

$\Delta\rho_{\min} = -0.42$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.07390 (14)	0.16015 (12)	−0.12063 (12)	0.0159 (3)
C2	1.21341 (15)	0.10073 (13)	−0.18674 (14)	0.0198 (3)
H2A	1.2930	0.1380	−0.1940	0.024*
C3	1.23134 (15)	−0.01528 (13)	−0.24169 (14)	0.0211 (3)
H3A	1.3252	−0.0581	−0.2878	0.025*
C4	1.11437 (15)	−0.07065 (13)	−0.23068 (13)	0.0197 (3)
H4	1.1304	−0.1505	−0.2689	0.024*
C5	0.97546 (15)	−0.01107 (13)	−0.16489 (13)	0.0177 (3)
H5	0.8958	−0.0484	−0.1574	0.021*
C6	0.95732 (14)	0.10512 (12)	−0.11042 (12)	0.0151 (2)
C7	0.84167 (14)	0.28675 (12)	−0.00034 (12)	0.0151 (2)
C8	0.55707 (14)	0.47110 (13)	0.32289 (12)	0.0167 (3)
C9	0.53199 (16)	0.36315 (14)	0.39425 (13)	0.0211 (3)
H9	0.6099	0.3004	0.4059	0.025*
C10	0.39098 (18)	0.34797 (15)	0.44861 (14)	0.0269 (3)
H10	0.3725	0.2748	0.4983	0.032*
C11	0.27802 (17)	0.43861 (17)	0.43063 (14)	0.0285 (3)
H11	0.1816	0.4289	0.4679	0.034*
C12	0.30799 (16)	0.54390 (16)	0.35725 (13)	0.0255 (3)
C13	0.44587 (15)	0.56403 (14)	0.30272 (13)	0.0209 (3)
H13	0.4637	0.6378	0.2538	0.025*
S1	1.01665 (3)	0.30546 (3)	−0.04133 (3)	0.01753 (8)
S2	0.73513 (3)	0.49153 (3)	0.25131 (3)	0.01507 (8)
N1	0.82742 (12)	0.17982 (11)	−0.04048 (10)	0.0154 (2)
H1	0.746 (2)	0.1638 (16)	−0.0270 (16)	0.018*
N2	0.73035 (13)	0.37297 (11)	0.06510 (11)	0.0176 (2)
H2	0.646 (2)	0.3709 (17)	0.0737 (16)	0.021*
N3	0.76169 (12)	0.48236 (11)	0.09996 (11)	0.0166 (2)
H3	0.7208 (19)	0.5525 (17)	0.0710 (16)	0.020*
N4	0.18823 (15)	0.63839 (18)	0.33454 (13)	0.0396 (4)
O1	0.82921 (11)	0.38488 (9)	0.28530 (10)	0.0220 (2)
O2	0.74518 (11)	0.61876 (9)	0.26804 (9)	0.0195 (2)
O3	0.21622 (15)	0.73477 (17)	0.27552 (14)	0.0584 (5)
O4	0.06728 (14)	0.61654 (19)	0.37558 (15)	0.0589 (5)
C14	0.31611 (14)	0.18627 (13)	0.18875 (13)	0.0164 (3)
C15	0.36568 (15)	0.07109 (13)	0.23976 (13)	0.0197 (3)
H15	0.4570	0.0192	0.1998	0.024*
C16	0.27603 (16)	0.03482 (14)	0.35187 (14)	0.0218 (3)
C17	0.14253 (15)	0.10749 (14)	0.41347 (13)	0.0219 (3)

H17	0.0849	0.0799	0.4907	0.026*
C18	0.09530 (15)	0.22141 (14)	0.35952 (14)	0.0212 (3)
H18	0.0036	0.2727	0.3995	0.025*
C19	0.18114 (15)	0.26111 (13)	0.24729 (13)	0.0194 (3)
H19	0.1480	0.3392	0.2103	0.023*
S3	0.42650 (3)	0.24195 (3)	0.04827 (3)	0.01634 (8)
N5	0.32639 (15)	-0.08700 (13)	0.40760 (13)	0.0315 (3)
O5	0.45472 (11)	0.36044 (9)	0.08076 (10)	0.0225 (2)
O6	0.55538 (10)	0.14188 (9)	0.00769 (9)	0.0200 (2)
O7	0.34166 (11)	0.26546 (10)	-0.03675 (10)	0.0224 (2)
O8	0.23692 (14)	-0.13450 (12)	0.48774 (11)	0.0393 (3)
O9	0.45499 (14)	-0.13511 (13)	0.37160 (14)	0.0493 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0170 (6)	0.0135 (6)	0.0176 (6)	-0.0045 (5)	-0.0045 (5)	-0.0013 (5)
C2	0.0157 (6)	0.0183 (6)	0.0247 (7)	-0.0044 (5)	-0.0044 (5)	-0.0016 (5)
C3	0.0179 (7)	0.0176 (6)	0.0247 (7)	-0.0003 (5)	-0.0041 (5)	-0.0034 (5)
C4	0.0241 (7)	0.0141 (6)	0.0214 (7)	-0.0027 (5)	-0.0081 (5)	-0.0031 (5)
C5	0.0202 (6)	0.0163 (6)	0.0195 (6)	-0.0065 (5)	-0.0082 (5)	-0.0002 (5)
C6	0.0149 (6)	0.0156 (6)	0.0149 (6)	-0.0037 (5)	-0.0048 (5)	0.0008 (5)
C7	0.0146 (6)	0.0172 (6)	0.0142 (6)	-0.0054 (5)	-0.0040 (5)	0.0006 (5)
C8	0.0162 (6)	0.0195 (6)	0.0141 (6)	-0.0047 (5)	-0.0032 (5)	-0.0019 (5)
C9	0.0269 (7)	0.0191 (7)	0.0171 (6)	-0.0079 (5)	-0.0041 (5)	-0.0012 (5)
C10	0.0321 (8)	0.0294 (8)	0.0193 (7)	-0.0163 (6)	-0.0011 (6)	-0.0025 (6)
C11	0.0225 (7)	0.0459 (9)	0.0186 (7)	-0.0158 (7)	-0.0005 (6)	-0.0091 (6)
C12	0.0178 (7)	0.0423 (9)	0.0155 (6)	-0.0017 (6)	-0.0056 (5)	-0.0059 (6)
C13	0.0193 (7)	0.0270 (7)	0.0143 (6)	-0.0020 (5)	-0.0042 (5)	-0.0004 (5)
S1	0.01279 (15)	0.01565 (15)	0.02306 (17)	-0.00532 (12)	-0.00137 (12)	-0.00507 (12)
S2	0.01461 (15)	0.01342 (15)	0.01680 (16)	-0.00269 (11)	-0.00443 (12)	-0.00064 (11)
N1	0.0128 (5)	0.0175 (5)	0.0167 (5)	-0.0055 (4)	-0.0039 (4)	-0.0018 (4)
N2	0.0124 (5)	0.0190 (6)	0.0214 (6)	-0.0053 (4)	-0.0025 (4)	-0.0063 (4)
N3	0.0174 (5)	0.0146 (5)	0.0174 (5)	-0.0046 (4)	-0.0036 (4)	-0.0021 (4)
N4	0.0208 (7)	0.0723 (12)	0.0192 (7)	0.0057 (7)	-0.0070 (5)	-0.0037 (7)
O1	0.0195 (5)	0.0190 (5)	0.0278 (5)	-0.0004 (4)	-0.0105 (4)	0.0006 (4)
O2	0.0224 (5)	0.0160 (5)	0.0196 (5)	-0.0059 (4)	-0.0044 (4)	-0.0021 (4)
O3	0.0337 (7)	0.0828 (12)	0.0411 (8)	0.0156 (7)	-0.0114 (6)	0.0212 (8)
O4	0.0166 (6)	0.1043 (14)	0.0500 (9)	-0.0015 (7)	-0.0084 (6)	-0.0081 (9)
C14	0.0150 (6)	0.0186 (6)	0.0185 (6)	-0.0065 (5)	-0.0071 (5)	0.0009 (5)
C15	0.0154 (6)	0.0201 (7)	0.0225 (7)	-0.0030 (5)	-0.0053 (5)	0.0020 (5)
C16	0.0209 (7)	0.0207 (7)	0.0222 (7)	-0.0030 (5)	-0.0066 (5)	0.0050 (5)
C17	0.0199 (7)	0.0261 (7)	0.0187 (7)	-0.0061 (6)	-0.0037 (5)	-0.0005 (5)
C18	0.0161 (6)	0.0239 (7)	0.0228 (7)	-0.0015 (5)	-0.0056 (5)	-0.0054 (5)
C19	0.0197 (7)	0.0174 (6)	0.0233 (7)	-0.0034 (5)	-0.0099 (5)	-0.0010 (5)
S3	0.01426 (15)	0.01613 (16)	0.02054 (17)	-0.00592 (12)	-0.00693 (12)	0.00324 (12)
N5	0.0284 (7)	0.0281 (7)	0.0269 (7)	0.0010 (5)	-0.0006 (5)	0.0102 (5)
O5	0.0195 (5)	0.0176 (5)	0.0322 (6)	-0.0076 (4)	-0.0083 (4)	0.0000 (4)

O6	0.0152 (5)	0.0189 (5)	0.0253 (5)	-0.0058 (4)	-0.0043 (4)	0.0014 (4)
O7	0.0230 (5)	0.0253 (5)	0.0236 (5)	-0.0090 (4)	-0.0126 (4)	0.0066 (4)
O8	0.0388 (7)	0.0325 (6)	0.0298 (6)	-0.0029 (5)	0.0058 (5)	0.0143 (5)
O9	0.0287 (7)	0.0444 (8)	0.0511 (8)	0.0100 (6)	0.0022 (6)	0.0252 (6)

Geometric parameters (Å, °)

C1—C2	1.3913 (19)	S2—O2	1.4260 (10)
C1—C6	1.3950 (18)	S2—O1	1.4279 (10)
C1—S1	1.7513 (13)	S2—N3	1.6574 (12)
C2—C3	1.3876 (19)	N1—H1	0.841 (19)
C2—H2A	0.9500	N2—N3	1.3985 (16)
C3—C4	1.399 (2)	N2—H2	0.826 (19)
C3—H3A	0.9500	N3—H3	0.869 (18)
C4—C5	1.386 (2)	N4—O4	1.223 (2)
C4—H4	0.9500	N4—O3	1.227 (2)
C5—C6	1.3863 (18)	C14—C15	1.3844 (19)
C5—H5	0.9500	C14—C19	1.3949 (19)
C6—N1	1.3946 (17)	C14—S3	1.7736 (14)
C7—N1	1.3215 (17)	C15—C16	1.3884 (19)
C7—N2	1.3317 (17)	C15—H15	0.9500
C7—S1	1.7218 (13)	C16—C17	1.383 (2)
C8—C9	1.3867 (19)	C16—N5	1.4677 (18)
C8—C13	1.3887 (19)	C17—C18	1.383 (2)
C8—S2	1.7674 (14)	C17—H17	0.9500
C9—C10	1.393 (2)	C18—C19	1.387 (2)
C9—H9	0.9500	C18—H18	0.9500
C10—C11	1.379 (2)	C19—H19	0.9500
C10—H10	0.9500	S3—O7	1.4526 (10)
C11—C12	1.383 (2)	S3—O6	1.4558 (10)
C11—H11	0.9500	S3—O5	1.4630 (10)
C12—C13	1.383 (2)	N5—O9	1.2244 (18)
C12—N4	1.469 (2)	N5—O8	1.2270 (17)
C13—H13	0.9500		
C2—C1—C6	121.20 (12)	O2—S2—C8	110.40 (6)
C2—C1—S1	128.05 (10)	O1—S2—C8	107.66 (6)
C6—C1—S1	110.75 (10)	N3—S2—C8	104.82 (6)
C3—C2—C1	117.18 (13)	C7—N1—C6	113.48 (11)
C3—C2—H2A	121.4	C7—N1—H1	120.7 (12)
C1—C2—H2A	121.4	C6—N1—H1	125.7 (12)
C2—C3—C4	121.52 (13)	C7—N2—N3	116.58 (11)
C2—C3—H3A	119.2	C7—N2—H2	123.0 (12)
C4—C3—H3A	119.2	N3—N2—H2	119.4 (12)
C5—C4—C3	121.15 (13)	N2—N3—S2	114.02 (9)
C5—C4—H4	119.4	N2—N3—H3	113.4 (12)
C3—C4—H4	119.4	S2—N3—H3	111.7 (12)
C6—C5—C4	117.39 (12)	O4—N4—O3	124.11 (16)

C6—C5—H5	121.3	O4—N4—C12	118.27 (17)
C4—C5—H5	121.3	O3—N4—C12	117.62 (15)
C5—C6—N1	126.36 (12)	C15—C14—C19	121.07 (13)
C5—C6—C1	121.56 (12)	C15—C14—S3	119.86 (11)
N1—C6—C1	112.08 (12)	C19—C14—S3	119.06 (10)
N1—C7—N2	122.94 (12)	C14—C15—C16	117.05 (13)
N1—C7—S1	114.10 (10)	C14—C15—H15	121.5
N2—C7—S1	122.96 (10)	C16—C15—H15	121.5
C9—C8—C13	121.98 (13)	C17—C16—C15	123.45 (13)
C9—C8—S2	119.96 (11)	C17—C16—N5	118.58 (13)
C13—C8—S2	118.04 (11)	C15—C16—N5	117.97 (13)
C8—C9—C10	119.11 (14)	C18—C17—C16	118.15 (13)
C8—C9—H9	120.4	C18—C17—H17	120.9
C10—C9—H9	120.4	C16—C17—H17	120.9
C11—C10—C9	120.37 (14)	C17—C18—C19	120.32 (13)
C11—C10—H10	119.8	C17—C18—H18	119.8
C9—C10—H10	119.8	C19—C18—H18	119.8
C10—C11—C12	118.62 (14)	C18—C19—C14	119.95 (13)
C10—C11—H11	120.7	C18—C19—H19	120.0
C12—C11—H11	120.7	C14—C19—H19	120.0
C13—C12—C11	123.14 (15)	O7—S3—O6	113.88 (6)
C13—C12—N4	118.14 (15)	O7—S3—O5	111.37 (6)
C11—C12—N4	118.71 (14)	O6—S3—O5	113.23 (6)
C12—C13—C8	116.76 (14)	O7—S3—C14	105.96 (6)
C12—C13—H13	121.6	O6—S3—C14	106.14 (6)
C8—C13—H13	121.6	O5—S3—C14	105.46 (6)
C7—S1—C1	89.58 (6)	O9—N5—O8	123.89 (14)
O2—S2—O1	120.92 (6)	O9—N5—C16	118.16 (13)
O2—S2—N3	104.08 (6)	O8—N5—C16	117.95 (13)
O1—S2—N3	107.77 (6)		
C6—C1—C2—C3	-0.1 (2)	C5—C6—N1—C7	-179.60 (13)
S1—C1—C2—C3	179.86 (11)	C1—C6—N1—C7	0.83 (16)
C1—C2—C3—C4	-0.3 (2)	N1—C7—N2—N3	-179.22 (12)
C2—C3—C4—C5	0.4 (2)	S1—C7—N2—N3	0.16 (17)
C3—C4—C5—C6	-0.1 (2)	C7—N2—N3—S2	-110.54 (12)
C4—C5—C6—N1	-179.80 (13)	O2—S2—N3—N2	-168.76 (9)
C4—C5—C6—C1	-0.3 (2)	O1—S2—N3—N2	61.70 (11)
C2—C1—C6—C5	0.4 (2)	C8—S2—N3—N2	-52.78 (11)
S1—C1—C6—C5	-179.57 (10)	C13—C12—N4—O4	175.11 (15)
C2—C1—C6—N1	179.98 (12)	C11—C12—N4—O4	-4.1 (2)
S1—C1—C6—N1	0.03 (14)	C13—C12—N4—O3	-5.0 (2)
C13—C8—C9—C10	-0.8 (2)	C11—C12—N4—O3	175.70 (16)
S2—C8—C9—C10	-179.19 (11)	C19—C14—C15—C16	-0.8 (2)
C8—C9—C10—C11	0.6 (2)	S3—C14—C15—C16	177.97 (11)
C9—C10—C11—C12	0.3 (2)	C14—C15—C16—C17	-0.2 (2)
C10—C11—C12—C13	-1.2 (2)	C14—C15—C16—N5	179.96 (13)
C10—C11—C12—N4	177.97 (14)	C15—C16—C17—C18	0.9 (2)

C11—C12—C13—C8	1.1 (2)	N5—C16—C17—C18	-179.26 (14)
N4—C12—C13—C8	-178.12 (13)	C16—C17—C18—C19	-0.6 (2)
C9—C8—C13—C12	-0.1 (2)	C17—C18—C19—C14	-0.4 (2)
S2—C8—C13—C12	178.39 (11)	C15—C14—C19—C18	1.1 (2)
N1—C7—S1—C1	1.13 (11)	S3—C14—C19—C18	-177.70 (11)
N2—C7—S1—C1	-178.31 (12)	C15—C14—S3—O7	125.09 (11)
C2—C1—S1—C7	179.43 (14)	C19—C14—S3—O7	-56.11 (12)
C6—C1—S1—C7	-0.62 (10)	C15—C14—S3—O6	3.69 (13)
C9—C8—S2—O2	-135.32 (11)	C19—C14—S3—O6	-177.51 (10)
C13—C8—S2—O2	46.19 (13)	C15—C14—S3—O5	-116.72 (12)
C9—C8—S2—O1	-1.39 (13)	C19—C14—S3—O5	62.08 (12)
C13—C8—S2—O1	-179.89 (11)	C17—C16—N5—O9	-162.03 (16)
C9—C8—S2—N3	113.16 (12)	C15—C16—N5—O9	17.8 (2)
C13—C8—S2—N3	-65.33 (12)	C17—C16—N5—O8	17.9 (2)
N2—C7—N1—C6	178.09 (12)	C15—C16—N5—O8	-162.28 (15)
S1—C7—N1—C6	-1.34 (15)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O6	0.841 (19)	1.888 (19)	2.7267 (15)	175.2 (17)
N2—H2...O5	0.826 (19)	1.92 (2)	2.7489 (16)	175.4 (18)
N3—H3...O7 ⁱ	0.869 (18)	1.968 (19)	2.8058 (16)	161.6 (16)
C2—H2 <i>A</i> ...O7 ⁱⁱ	0.95	2.55	3.2510 (18)	130
C9—H9...O8 ⁱⁱⁱ	0.95	2.58	3.487 (2)	161

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $x+1, y, z$; (iii) $-x+1, -y, -z+1$.