

Crystal structures and Hirshfeld surfaces of four methoxybenzaldehyde oxime derivatives, 2-MeO- $\text{XC}_6\text{H}_3\text{C}=\text{NOH}$ ($\text{X} = \text{H}$ and 2-, 3- and 4-MeO): different conformations and hydrogen-bonding patterns

Ligia R. Gomes,^{a,b} Marcus V. N. de Souza,^c Cristiane F. Da Costa,^c James L. Wardell^{c,d} and John Nicolson Low^{d,*}

Received 1 October 2018

Accepted 3 October 2018

Edited by P. McArdle, National University of Ireland, Ireland

Keywords: crystal structure; oxime derivative; hydrogen bonding.

CCDC references: 1871165; 1871164; 1871163; 1871162

Supporting information: this article has supporting information at journals.iucr.org/e


^aREQUIMTE, Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre, 687, P-4169-007, Porto, Portugal, ^bFP-ENAS-Faculdade de Ciências de Saúde, Escola Superior de Saúde da UFP, Universidade Fernando Pessoa, Rua Carlos da Maia, 296, P-4200-150 Porto, Portugal, ^cInstituto de Tecnologia em Fármacos e Farmanguinhos, Fundação Oswaldo Cruz, 21041-250 Rio de Janeiro, RJ, Brazil, and ^dDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen, AB24 3UE, Scotland. *Correspondence e-mail: jnlow111@gmail.com

The crystal structures of four (*E*)-methoxybenzaldehyde oxime derivatives, namely (2-methoxybenzaldehyde oxime, **1**, 2,3-dimethoxybenzaldehyde oxime, **2**, 4-dimethoxybenzaldehyde oxime, **3**, and 2,5-dimethoxybenzaldehyde oxime, **4**), are discussed. The arrangements of the 2-methoxy group and the H atom of the oxime unit are *s-cis* in compounds **1–3**, but in both independent molecules of compound **4**, the arrangements are *s-trans*. There is also a difference in the conformation of the two molecules in **4**, involving the orientations of the 2- and 5-methoxy groups. The primary intermolecular O—H(oxime)···O(hydroxy) hydrogen bonds generate *C*(3) chains in **1** and **2**. In contrast, in compound **3**, the O—H(oxime)···O(hydroxy) hydrogen bonds generate symmetric *R*₂²(6) dimers. A more complex dimer is generated in **4** from the O—H(oxime)···O(hydroxy) and C—H(2-methoxy)···O(hydroxy) hydrogen bonds. In all cases, further interactions, C—H···O and C—H··· π or π — π , generate three-dimensional arrays. Hirshfeld surface and fingerprint analyses are discussed.

1. Chemical context

In the plant kingdom, oximes play a vital role in metabolism (Sørensen *et al.*, 2018). Aldoximes, $\text{RCH}=\text{NOH}$, are found in many biologically active compounds (Abele *et al.*, 2008; Nikitjuka & Jirgensons, 2014), having a diverse range of uses including as anti-tumour agents (Martínez-Pascual *et al.*, 2017; Qin *et al.*, 2017; Canario *et al.*, 2018; Huang *et al.*, 2018), acaricidal and insecticidal agents (Dai *et al.*, 2017), thymidine phosphorylase inhibitors (Zhao *et al.*, 2018), anti-microbial agents (Yadav *et al.*, 2017), bacteriocides (Kozłowska *et al.*, 2017), anti-inflammatory agents (Mohassab *et al.* 2017), and in the treatment of nerve-gas poisoning (Lorke *et al.*, 2008; Voicu *et al.*, 2010; Katalinić *et al.*, 2017; Radić *et al.*, 2013).

Benzaldehyde oximes, $\text{ArCH}=\text{NOH}$, with their $-\text{CH}=\text{N}-\text{OH}$ functional group are ideally arranged for classical O—H···O and/or O—H···N hydrogen bonding. The last survey of the classical hydrogen-bonding patterns in benzaldehyde oximes reported in 2010 (Low *et al.*, 2010) confirmed that the most frequently found arrangements, with the exception of salicylaloximes, are *R*₂²(6) dimers and *C*(3) chains, Fig. 1.







Figure 1
Illustrations of the $C(3)$ chains and $R_2^2(6)$ dimers formed by oximes

Aakeröy *et al.* (2013) reported the percentages of $R_2^2(6)$ dimers and $C(3)$ chains found in non-salicylaloxime to be *ca* 72 and 24%, respectively – similar percentages can be derived from a recent survey of the Cambridge Structural Database (CSD Version 5.39, August 2018 update; Groom *et al.*, 2016). Hydrogen bonds are considered to be the strongest and most directional of intermolecular interactions in molecules (Etter, 1990) and thus play the major roles in determining the overall supramolecular structures. However, the involvement of weaker intermolecular interactions, such as $C-H \cdots O$ hydrogen bonds, $\pi-\pi$ interactions and interactions involving the substituents, can have a significant influence on the supramolecular arrays generated. In a continuation of recent studies on aldoximes (Low *et al.* 2018; Gomes *et al.*, 2018), we have determined the crystal structures of four methoxybenzaldehyde derivatives, namely 2-MeO- X - $C_6H_3CH=NOH$ where $X = H$ in **1**, $X = 3$ -MeO in **2**, $X = 4$ -MeO in **3** and $X = 5$ -MeO in **4**. The aim of the study was to further investigate the occurrence of $R_2^2(6)$ dimers and $C(3)$ chains in a series of related compounds.



2. Structural commentary

There are no unusual features in the molecular structures. Compound **1** crystallizes in the *orthorhombic* space group $Pna2_1$ with one molecule in the asymmetric unit (Fig. 2), compound **2** crystallizes in the *orthorhombic* space group $P2_12_12_1$ with one molecule in the asymmetric unit (Fig. 3), compound **3** crystallizes in the *triclinic* space group $P\bar{1}$ with one molecule in the asymmetric unit (Fig. 4), and compound **4** crystallizes in the *monoclinic* space group, $P2_1/c$ with two independent molecules, Mol A and Mol B, in the asymmetric unit (Fig. 5). The geometry about the oxime moiety in all molecules is (*E*). In compounds **1–3**, the 2-methoxy group and the hydrogen of the oxime moiety have an *s-cis* arrangement.




Figure 2
Atom arrangements and numbering scheme for compound **1**. Displacement ellipsoids are drawn at the 50% probability level.




Figure 3
Atom arrangements and numbering system for compound **2**. Displacement ellipsoids are drawn at the 50% probability level.




Figure 4
Atom arrangements and numbering system for compound **3**. Displacement ellipsoids are drawn at the 50% probability level.




Figure 5 Mol A Mol B
Atom arrangements and numbering system for the two independent molecules, Mol A and Mol B, of compound **4**. Displacement ellipsoids are drawn at the 50% probability level.




Figure 6
Compound **1**. Part of a $C(3)$ chain formed by $O13-H13 \cdots N12$ hydrogen bonds (dashed lines; see Table 2).

In contrast, in both molecules of compound **4**, the 2-methoxy group and the hydrogen atom of the oxime moiety have an *s-trans* arrangement. The *s-trans* arrangement of the 2-alkoxy group and hydrogen atom of the oxime units in compound **4** is very much rarer than the *s-cis* arrangement found in compounds **1–3** and other non-salicylaldoximes. A search of the Cambridge Structural Database (CSD Version 5.39, August 2018 update; Groom *et al.*, 2016) revealed that only salicylaldoximes and 2-alkoxybenzaldehyde oxime (*E*)-2-([2-[(*E*)-(hydroxyimino)methyl]phenoxy)methyl]-3-*p*-tolylacrylonitrile (LAQRIG; Suresh *et al.* 2012) had this *s-trans* arrangement. In contrast, the isomer 2-([2-[(hydroxyimino)methyl]phenoxy)methyl]-3-(2-methylphenyl)acrylonitrile (GARNEU; Govindan *et al.*, 2012a) and some similar compounds such as (*E*)-2-([2-[(*E*)-(hydroxyimino)methyl]phenoxy)methyl]-3-phenylacrylonitrile (LAQRUS; Govindan *et al.*, 2012b) had the *s-cis* arrangement.

There is a conformational difference between the two independent molecules Mol A and Mol B of compound **4**. This

Table 1
Distances (Å) of OMe C atoms and oxime N and O atoms from benzene ring mean plane in compounds **1–4**.

Atom	1	2	3	4 Mol A	4 Mol B
C21	0.086 (3)	−1.140 (4)	0.195 (1)	0.121 (1)	0.059 (1)
C31	–	−0.011 (4)	–	–	–
C41	–	–	0.081 (1)	–	–
C51	–	–	–	0.033 (1)	0.061 (1)
N12	0.061 (2)	0.259 (3)	−0.177 (1)	0.264 (1)	−0.020 (1)
O13	−0.009 (2)	−0.027 (3)	0.051 (1)	0.242 (1)	0.010 (1)

Table 2
Hydrogen-bond geometry (Å, °) for **1**.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O13-H13 \cdots N12^i$	0.84	1.93	2.764 (2)	170
$C3-H3 \cdots O13^{ii}$	0.95	2.50	3.442 (2)	174
$C21-H21C \cdots O13^{iii}$	0.98	2.57	3.506 (3)	160

Symmetry codes: (i) $-x+1, -y+1, z-\frac{1}{2}$; (ii) $x+\frac{1}{2}, -y+\frac{1}{2}, z+1$; (iii) $x+\frac{1}{2}, -y+\frac{1}{2}, z$.

difference is in the orientation of the two methoxy groups, see Fig. 5: in Mol A the orientation is *s-trans* and in Mol B, it is *s-cis*. As expected for a 1,2,3-trisubstituted benzene derivative, compound **4** is the least planar of the four oxime derivatives, with the 2-methoxy substituent furthest out of the plane of the attached phenyl group, see Table 1.

3. Supramolecular features

3.1. Hydrogen bonding

In the crystal of **1**, molecules are primarily linked by strong $O13-H13 \cdots N12^i$ hydrogen bonds (Table 2), forming $C(3)$ chains, illustrated in Fig. 6. Also present in compound **1** are two weaker hydrogen bonds, namely, $C3-H3 \cdots O13^{ii}$ and $C21-H21C \cdots O13^{iii}$, as well as a weak $\pi-\pi$ stacking interaction [$Cg \cdots Cg^{iv} = 4.025 (2)$ Å; slippage 2.105 Å; symmetry code; $x, y, z - 1$]. These three interactions generate the molecular arrangement shown in Fig. 7. The $C3-H3 \cdots O13^{ii}$ hydrogen bonds generate $C(7)$ chains in the c -axis direction, while the $C21-H21C \cdots O13^{iii}$ hydrogen bonds form $C(8)$




Figure 7
Compound **1**. Part of the arrangement generated from the combination of hydrogen bonds and $\pi-\pi$ interactions (dashed lines; see Table 2).




Figure 8
Compound **2**. Part of the arrangement generated from C21–H21B...O31, C31–H31B...O13 and π – π interactions (dashed lines; see Table 3).

spiral chains along the *a*-axis direction: together these hydrogen bonds form $R_4^4(22)$ rings. The tilted π – π stacks propagate in the *c*-axis direction. The involvement of the weaker C3–H3...O13ⁱⁱ, C21–H21C...O13ⁱⁱⁱ and π – π interactions, along with the stronger O13–H13...N12ⁱ hydrogen bonds, creates the three-dimensional structure for **1**.

As in **1**, molecules of **2** are primarily linked by strong O13–H13...N12ⁱ hydrogen bonds (Table 3), forming $C(3)$ chains: as such chains are very similar to those in compound **1**, see Fig. 6, an illustration has not been provided for the $C(3)$ chain in compound **2**. Other intermolecular interactions in **2** are the weaker C21–H21B...O31ⁱⁱⁱ and C31–H31B...O13^{iv} hydrogen bonds and a C31–H31C...Cg1^v interaction involving the C1–C6 ring. These three interactions combine to form the arrangement illustrated in Fig. 8. The C21–H21B...O31ⁱⁱⁱ hydrogen bonds on their own generate $C(6)$ chains, which

Table 3
Hydrogen-bond geometry (\AA , $^\circ$) for **2**.

Cg1 is the centroid of the C1–C6 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
O13–H13...N12 ⁱ	0.97 (4)	1.87 (5)	2.805 (4)	161 (4)
C4–H4...O21 ⁱⁱ	0.95	2.63	3.284 (4)	126
C21–H21B...O31 ⁱⁱⁱ	0.98	2.54	3.323 (5)	136
C31–H31B...O13 ^{iv}	0.98	2.51	3.448 (5)	161
C31–H31C...Cg1 ^v	0.98	2.73	3.599 (5)	148

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - 1, y, z$; (iv) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (v) $x + 1, y, z$.

Table 4
Hydrogen-bond geometry (\AA , $^\circ$) for **3**.

Cg1 is the centroid of the C1–C6 ring.

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
O13–H13...N12 ⁱ	0.893 (18)	1.995 (19)	2.8124 (13)	151.5 (15)
C41–H41A...O13 ⁱⁱ	0.98	2.63	3.0680 (15)	107
C41–H41C...Cg1 ⁱⁱⁱ	0.98	2.60	3.4479 (13)	144

Symmetry codes: (i) $-x, -y, -z$; (ii) $x + 2, y + 1, z$; (iii) $x + 1, y, z$.

propagate in the *a*-axis direction while the C31–H31B...O13^{iv} hydrogen bonds generate spiral $C(9)$ chains in the *b*-axis direction. Together these hydrogen bonds generate a network of $R_4^4(26)$ rings. The C31–H31C...Cg1^v interactions lead to chains along the *a*-axis direction. The involvement of the weaker C21–H21B...O31ⁱⁱⁱ, C31–H31B...O13^{iv} C and C–H... π interactions, along with the stronger O13–H13...N12ⁱ hydrogen bonds, creates a three-dimensional structure for **2**. C4–H4...O12ⁱⁱ hydrogen bonds also occur.

In compound **3**, $R_2^2(6)$ dimers are generated from strong O13–H13...N12ⁱ hydrogen bonds (Table 4), as illustrated in Fig. 9. Linkages of these $R_2^2(6)$ dimers by weaker C41–H41A(methoxy)...O13ⁱⁱ hydrogen bonds provide a two-molecule-wide ribbon. Within the ribbons are $R_4^4(22)$ rings as well as the $R_2^2(6)$ rings. An additional interaction in **3** is the C41–H41C...Cg1ⁱⁱⁱ interaction, which generates a tilted ladder assembly, propagating in the *a*-axis direction, with the $R_2^2(6)$




Figure 9
Compound **3**. A two-molecule-wide ribbon generated from linking the $R_2^2(6)$ dimers, formed by pairs of strong O13–H13–N12 hydrogen bonds and by weaker C41–H41A...O13 hydrogen bonds (dashed lines; see Table 4).

Table 5
Hydrogen-bond geometry (Å, °) for **4**.

Cg_1 and Cg_2 are the centroids of the C11–C16 and C21–C26 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O113—H113 \cdots O121 ⁱ	0.875 (16)	2.247 (15)	2.8944 (9)	130.7 (12)
O113—H113 \cdots N112 ⁱ	0.875 (16)	1.965 (16)	2.7567 (10)	149.9 (13)
O213—H213 \cdots O221 ⁱⁱ	0.877 (15)	2.204 (15)	2.8758 (9)	133.1 (12)
O213—H213 \cdots N212 ⁱⁱ	0.877 (15)	2.034 (15)	2.8160 (10)	147.9 (13)
C111—H111 \cdots O251	0.95	2.46	3.2458 (11)	140
C121—H12C \cdots N212 ⁱⁱⁱ	0.98	2.53	3.4400 (13)	155
C151—H15A \cdots O113 ^{iv}	0.98	2.50	3.3947 (11)	152
C14—H14 \cdots Cg_2 ⁱⁱⁱ	0.95	2.98	3.6656 (9)	130
C151—H15B \cdots Cg_2	0.98	2.72	3.5973 (10)	149
C24—H24 \cdots Cg_1 ^v	0.95	2.67	3.4281 (10)	137
C211—H211 \cdots Cg_1 ^{vi}	0.95	2.78	3.6272 (9)	149

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$; (iii) $x+1, -y+\frac{1}{2}, z-\frac{1}{2}$; (iv) $x, -y+\frac{1}{2}, z-\frac{1}{2}$; (v) $x-1, y, z$; (vi) $x, -y+\frac{1}{2}, z+\frac{1}{2}$.

rings acting as the rungs and the C41—H41C \cdots Cg_1 ⁱⁱⁱ interactions as the supports.

In compound **4**, each of the two independent molecules forms symmetric dimers, see Fig. 10. These are generated from combinations of O113—H113 \cdots N112ⁱ and O113—H113 \cdots O121ⁱ hydrogen bonds (Table 5) for Mol A and O213—H213 \cdots N212ⁱⁱ and O213—H213 \cdots O221ⁱⁱ hydrogen bonds for Mol B. In each case, the dimers contain three rings, two $R_2^2(6)$ and one $R_2^2(6)$. There are short N \cdots N distances across the $R_2^2(6)$ dimer rings, 2.8595 (12) Å for MolA and 2.8956 (12) Å for Mol B, each being less than the sum of the van der Waals radius (3.10 Å) for two N atoms.

The links between the two different dimers of **4** are provided by a number of C—H \cdots O and C—H \cdots π interactions, listed in Table 5. Fig. 11 restricts the contacts to just the C—H \cdots O hydrogen bonds, namely C121—H12C \cdots N212ⁱⁱⁱ, C111—H111 \cdots O251 and C151—H15A \cdots O113^{iv}. To




Figure 10
Compound **4**. Symmetric dimers of (a) Mol A and (b) Mol B. Hydrogen bonds (see Table 5) are shown as dashed lines.




Figure 11
Compound **4**. Symmetric dimers of Mol A (green) and Mol B (blue). Intermolecular interactions (see Table 5) are shown as dashed lines.

facilitate the viewing of the connection in Fig. 11, the two different dimers are drawn in different colours.

3.2. Hirshfeld surface analysis

Hirshfeld surfaces (Spackman & Jayatilaka, 2009) and two-dimensional fingerprint (FP) plots (Spackman & McKinnon, 2002), provide complementary information concerning the intermolecular interactions discussed above. The analyses were generated using *Crystal Explorer3.1* (Wolff *et al.*, 2012). The Hirshfeld surfaces mapped over d_{norm} for **1–4** are illustrated in Fig. 12. The red areas on the surfaces correspond to close contacts. The fingerprint plots are shown in Fig. 13. In all of the FP plots, the pair of spikes pointing south-west relate to the N—H contacts, which in compounds **1** and **2** are involved in the C(3) chains, while in compounds **3** and **4**, they are responsible for the creation of the dimers. In compound **3**, the fins ending at $d_e, d_i = 1.9, 1.1$ Å are due to C(π) \cdots H/C(π) \cdots H contacts. The FP plots for Mol A and Mol B of compound **4** are asymmetric because of the different interactions of each molecule. The double wings in the FP plot for Mol A in the second quadrant are complementary to those displayed in the fourth quadrant by MolB and relate to C \cdots H close contacts connecting the two molecules. The spike ending at $d_i, d_e = 1.1$ Å in Mol A is due to H \cdots H contacts.

The percentages of the various atom–atom contacts, derived from the fingerprint plots, for the four compounds are shown in Table 6. The fact that compound **1** has only one methoxy group while the isomers, **2–4**, have two is reflected in the greater percentages of contacts involving the oxygen close contacts. The C(3)-chain-forming compounds **1** and **2** show higher percentages of H \cdots H and C \cdots C contacts, but a lower percentage of H \cdots C/C \cdots H contacts, than the dimer-forming compounds **3** and **4**.

Table 6

Percentages of atom–atom contacts for compounds **1**, **2**, **3** and **4** (Mol A and Mol B).

Compound	1	2	3	4 Mol A	4 Mol B
H···H	52.7	49.1	43.7	41.5	38.6
H···O/O···H	16.2	22.5	23.4	24.9	26.3
H···C/C···H	11.3	14.5	20.4	22.7	25.9
H···N/N···H	8.1	6.6	8.4	9.0	8.1
C···C	7.9	3.5	1.3	0.1	0.1
O···C/C···O	2.1	2.0	2.6	1.5	0.8
N···O/O···N	–	–	–	–	–
N···C/C···N	1.6	1.8	–	–	–
O···O	–	–	–	0.4	0.2

4. Database survey

A search of the Cambridge Structural Database survey (CSD Version 5.39, August 2018 update; Groom *et al.*, 2016) revealed compounds similar to **2** and **3**. The classical hydrogen bonds in 3,5-dimethoxybenzene oxime generate $C(3)$ chains (VUZJAC; Dong *et al.*, 2010). No benzene oxime derivative with only methoxy substituents has been reported in the database to form an $R_2^2(6)$ or related dimer. The structure has been reported of 3,4,5-trimethoxybenzene oxime (MEQDAO; Chang, 2006) in which classical hydrogen bonds, formed between the oxime unit and the 4- and 5-methoxy moieties, result in the formation of a tetramer. The water molecule in 3,4,5-trimethoxybenzene monohydrate (HESWUY; Priya *et al.*, 2006) is strongly involved in the hydrogen-bonding arrangements.

There are 376 structures, (411 fragments) in the CSD database with oxime $R_2^2(6)$ dimers in which the N···N distance across the ring is less than or equal to 3.10 Å, the sum of two




Figure 13
Fingerprint plots for compounds **1–4**.

N-atom van der Waals radii. The H···O hydrogen-bond distance range was restricted to 1.739–2.285 Å to exclude improbable O···H distances based on a statistical analysis in *Mercury* (Macrae *et al.*, 2006). The N···N distances range from 2.727 to 3.097 Å with a mean value of 2.987 Å. There are 27 structures within the range 2.838 to 2.909 Å in which our values of 2.8595 (12) Å for MolA and 2.8596 (12) Å for MolB of compound **4** lie. Only single-crystal organic compounds were searched for with no limit on the R factor.

5. Synthesis and crystallization

The title compounds were prepared from hydroxyamine and the corresponding benzaldehyde in methanol in the presence




Figure 12
Hirshfeld surfaces for compounds **1–4**. In each case, the interactions related to the red areas are designated.

Table 7
Experimental details.

	1	2	3	4
Crystal data				
Chemical formula	C ₈ H ₉ NO ₂	C ₉ H ₁₁ NO ₃	C ₉ H ₁₁ NO ₃	C ₉ H ₁₁ NO ₃
<i>M_r</i>	151.16	181.19	181.19	181.19
Crystal system, space group	Orthorhombic, <i>Pna</i> 2 ₁	Orthorhombic, <i>P2</i> ₁ 2 ₁ 2 ₁	Triclinic, <i>P</i> $\bar{1}$	Monoclinic, <i>P2</i> ₁ / <i>c</i>
Temperature (K)	100	100	100	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	11.1719 (2), 16.4260 (3), 4.0249 (1)	4.6775 (2), 13.0996 (5), 14.1984 (5)	4.9441 (2), 8.2188 (4), 12.1308 (3)	7.6480 (1), 21.3380 (4), 10.9421 (2)
α , β , γ (°)	90, 90, 90	90, 90, 90	108.849 (3), 92.288 (3), 106.273 (4)	90, 90.555 (2), 90
<i>V</i> (Å ³)	738.61 (3)	869.98 (6)	443.17 (3)	1785.59 (5)
<i>Z</i>	4	4	2	8
Radiation type	Cu <i>K</i> α	Cu <i>K</i> α	Cu <i>K</i> α	Mo <i>K</i> α
μ (mm ⁻¹)	0.82	0.87	0.86	0.10
Crystal size (mm)	0.05 × 0.05 × 0.03	0.30 × 0.05 × 0.02	0.20 × 0.10 × 0.05	0.20 × 0.15 × 0.13
Data collection				
Diffractometer	Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector	Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector	Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector	Rigaku FRE+ equipped with VHF Varimax confocal mirrors and an AFC12 goniometer and HyPix 6000 detector
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku OD, 2017)	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku OD, 2017)	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku OD, 2017)	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku OD, 2017)
<i>T</i> _{min} , <i>T</i> _{max}	0.848, 1.000	0.507, 1.000	0.802, 1.000	0.935, 1.000
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	12857, 1345, 1325	7835, 1596, 1371	7618, 1594, 1462	38753, 4082, 3761
<i>R</i> _{int}	0.038	0.095	0.033	0.020
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.602	0.602	0.602	0.649
Refinement				
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.033, 0.088, 1.08	0.058, 0.151, 1.04	0.035, 0.101, 0.88	0.031, 0.086, 1.06
No. of reflections	1345	1596	1594	4082
No. of parameters	102	124	124	247
No. of restraints	1	0	0	0
H-atom treatment	H-atom parameters constrained	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.17, -0.16	0.35, -0.20	0.20, -0.19	0.32, -0.19
Absolute structure	Refined as a perfect inversion twin.	Flack <i>x</i> determined using 474 quotients [(<i>I</i> ⁺) - (<i>I</i> ⁻)] / [(<i>I</i> ⁺) + (<i>I</i> ⁻)] (Parsons <i>et al.</i> , 2013)	-	-
Absolute structure parameter	0.5	0.2 (3)	-	-

Computer programs: *CrysAlis PRO* (Rigaku OD, 2017), *OSCAIL* (McArdle *et al.*, 2004), *SHELXT* (Sheldrick, 2015a), *ShelXle* (Hübschle *et al.*, 2011), *SHELXL* (Sheldrick, 2015b), *Mercury* (Macrae *et al.*, 2006) and *PLATON* (Spek, 2009).

of potassium carbonate and were recrystallized from methanol solutions, m.p. = 364–365 K for compound **1**, 371–373 K for **2**, 378–380 K for **3** and 370–371 K for **4**.

6. Refinement details

Crystal data, data collection and structure refinement details are summarized in Table 7. All hydroxy hydrogen atoms were refined isotropically. C-bound H atoms were refined as riding with C–H = 0.95–0.98 Å and *U*_{iso}(H) = 1.2–1.5 *U*_{eq}(C).

Acknowledgements

The authors thank the staff at the National Crystallographic Service, University of Southampton, for the data collection, help and advice (Coles & Gale, 2012).

References

- Aakeröy, C. B., Sinha, A. S., Epa, K. N., Chopade, P. D., Smith, M. M. & Desper, J. (2013). *Cryst. Growth Des.* **13**, 2687–2695.
- Abele, E., Abele, R. & Lukevics, E. (2008). *Chem. Heterocycl. Cmpds*, **44**, 769–792.
- Canario, C., Silvestre, S., Falcao, A. & Alves, G. (2018). *Curr. Med. Chem.* **25**, 660–686.
- Chang, X.-H. (2006). *Acta Cryst.* **E62**, o5699–o5700.
- Coles, S. J. & Gale, P. A. (2012). *Chem. Sci.* **3**, 683–689.
- Dai, H., Chen, J., Li, G., Ge, S. S., Shi, Y. J., Fang, Y. & Ling, Y. (2017). *Bioorg. Med. Chem. Lett.* **27**, 950–953.
- Dong, B., Zhang, Y. & Chen, J.-Z. (2010). *Acta Cryst.* **E66**, o2719.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Gomes, L. R., de Souza, M. V. N., Da Costa, C. F., Wardell, J. L. & Low, J. N. (2018). *Acta Cryst.* **E74**, 1480–1485.
- Govindan, E., Srinivasan, J., Bakthadoss, M. & SubbiahPandi, A. (2012a). *Acta Cryst.* **E68**, o484.
- Govindan, S., Vijayakumar, S., Jayakumar, S., Mannickam, B. & Sanmargam, A. (2012b). *Acta Cryst.* **E68**, o596.

- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Huang, G., Zhao, H. R., Meng, Q. Q., Zhang, Q. J., Dong, J. Y., Zhu, B. Q. & Li, S. S. (2018). *Eur. J. Med. Chem.* **143**, 166–181.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). *J. Appl. Cryst.* **44**, 1281–1284.
- Katalinić, M., Zandona, A., Ramić, A., Zorbaz, T., Primožič, I. & Kovarik, Z. (2017). *Molecules*, **22**, 1234.
- Kozłowska, J., Potaniec, B., Żarowska, B. & Anioł, M. (2017). *Molecules*, **22**, 1485.
- Lorke, D. E., Kalasz, H., Petroianu, G. A. & Tekes, K. (2008). *Curr. Med. Chem.* **15**, 743–753.
- Low, J. N., Santos, L. M. N. B. F., Lima, C. F. R. A. C., Brandão, P. & Gomes, L. R. (2010). *Eur. J. Chem.* **1**, 61–66.
- Low, J. N., Wardell, J. L., Da Costa, C. F., Souza, M. V. N. & Gomes, L. R. (2018). *Eur. J. Chem.* **9**, 151–160.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Martínez-Pascual, R., Meza-Reyes, S., Vega-Baez, J. L., Merino-Montiel, P., Padrón, J. M., Mendoza, Á. & Montiel-Smith, S. (2017). *Steroids*, **122**, 24–33.
- McArdle, P., Gilligan, K., Cunningham, D., Dark, R. & Mahon, M. (2004). *CrystEngComm*, **6**, 303–309.
- Mohassab, M., Hassan, H. A., Abdelhamid, D., Abdel-Aziz, M., Dalby, K. N. & Kaoud, T. S. (2017). *Bioorg. Chem.* **75**, 242–259.
- Nikitjuka, A. & Jirgensons, A. (2014). *Chem. Heterocycl. Compd.* **49**, 1544–1559.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Priya, B. S., Basappa, S. N., Anandalwar, S. M., Prasad, J. S. & Rangappa, K. S. (2006). *Anal. Sci.:X-Ray Structures Online*, **22**, x161–x162.
- Qin, H. L., Leng, J., Youssif, B. G. M., Amjad, M. W., Raja, M. A. G., Hussain, M. A., Hussain, Z., Kazmi, S. N. & Bukhari, S. N. A. (2017). *Chem. Biol. Drug Des.* **90**, 443–449.
- Radić, Z., Dale, T., Kovarik, Z., Berend, S., Garcia, E., Zhang, L., Amitai, G., Green, C., Radić, B., Duggan, B. M., Ajami, D., Rebek, J. Jr & Taylor, P. (2013). *Biochem. J.* **450**, 231–242.
- Rigaku OD (2017). *CrysAlis PRO*. Rigaku Oxford Diffraction, Yarnton, England.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Sørensen, M., Neilson, E. H. J. & Møller, B. L. (2018). *Mol. Plant.* **11**, 95–117.
- Spackman, M. A. & Jayatilaka, D. (2009). *CrystEngComm*, **11**, 19–32.
- Spackman, M. A. & McKinnon, J. J. (2002). *CrystEngComm*, **4**, 378–392.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Suresh, G., Sabari, V., Srinivasan, J., Mannickam, B. & Aravindhan, S. (2012). *Acta Cryst.* **E68**, o570.
- Voicu, V. A., Thiermann, H., Rădulescu, F. Ş., Mircioiu, C. & Miron, D. S. (2010). *Basic Clin. Pharmacol. Toxicol.* **106**, 73–85.
- Wolff, S. K., Grimwood, D. I., McKinnon, J. J., Turner, M. J., Jayatilaka, D. & Spackman, M. A. (2012). *Crystal Explorer*. The University of Western Australia.
- Yadav, P., Lal, K., Rani, P., Mor, S., Kumar, A. & Kumar, A. (2017). *Med. Chem. Res.* **26**, 1469–1480.
- Zhao, S. Y., Li, K., Jin, Y. & Lin, J. (2018). *Eur. J. Med. Chem.* **144**, 41–51.

supporting information

Acta Cryst. (2018). E74, 1553-1560 [https://doi.org/10.1107/S2056989018014020]

Crystal structures and Hirshfeld surfaces of four methoxybenzaldehyde oxime derivatives, 2-MeO- $\text{XC}_6\text{H}_3\text{C}\&z\text{-dbnd;NOH}$ ($X = \text{H}$ and 2-, 3- and 4-MeO): different conformations and hydrogen-bonding patterns

Ligia R. Gomes, Marcus V. N. de Souza, Cristiane F. Da Costa, James L. Wardell and John Nicolson Low

Computing details

For all structures, data collection: *CrysAlis PRO* (Rigaku OD, 2017); cell refinement: *CrysAlis PRO* (Rigaku OD, 2017); data reduction: *CrysAlis PRO* (Rigaku OD, 2017); program(s) used to solve structure: *OSCAIL* (McArdle *et al.*, 2004) and *SHELXT* (Sheldrick, 2015a); program(s) used to refine structure: *OSCAIL* (McArdle *et al.*, 2004), *ShelXle* (Hübschle *et al.*, 2011) and *SHELXL* (Sheldrick, 2015b); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *OSCAIL* (McArdle *et al.*, 2004), *SHELXL* (Sheldrick, 2015b) and *PLATON* (Spek, 2009).

2-Methoxy-benzaldehyde oxime (1)

Crystal data

$\text{C}_8\text{H}_9\text{NO}_2$

$M_r = 151.16$

Orthorhombic, *Pna*2₁

$a = 11.1719$ (2) Å

$b = 16.4260$ (3) Å

$c = 4.0249$ (1) Å

$V = 738.61$ (3) Å³

$Z = 4$

$F(000) = 320$

$D_x = 1.359$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 7376 reflections

$\theta = 2.7\text{--}70.0^\circ$

$\mu = 0.82$ mm⁻¹

$T = 100$ K

Block, colourless

$0.05 \times 0.05 \times 0.03$ mm

Data collection

Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector diffractometer

Radiation source: Rotating anode, Rigaku 007 HF

Detector resolution: 10 pixels mm⁻¹
profile data from ω -scans

Absorption correction: multi-scan
(*CrysAlis PRO*; Rigaku OD, 2017)

$T_{\min} = 0.848$, $T_{\max} = 1.000$

12857 measured reflections

1345 independent reflections

1325 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.038$

$\theta_{\max} = 68.2^\circ$, $\theta_{\min} = 4.8^\circ$

$h = -13 \rightarrow 13$

$k = -19 \rightarrow 19$

$l = -4 \rightarrow 4$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.033$ $wR(F^2) = 0.088$ $S = 1.08$

1345 reflections

102 parameters

1 restraint

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0559P)^2 + 0.1277P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$ Absolute structure: Refined as a perfect
inversion twin.

Absolute structure parameter: 0.5

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refined as a 2-component perfect inversion twin.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O13	0.45979 (12)	0.42019 (8)	-0.4095 (4)	0.0311 (4)
H13	0.443792	0.464123	-0.506595	0.047*
O21	0.69333 (12)	0.21322 (8)	0.0350 (4)	0.0303 (4)
N12	0.56325 (15)	0.43011 (9)	-0.2166 (5)	0.0265 (4)
C1	0.72029 (18)	0.35490 (12)	0.0488 (5)	0.0260 (5)
C2	0.76289 (18)	0.27720 (11)	0.1386 (6)	0.0266 (5)
C3	0.86797 (18)	0.26830 (12)	0.3172 (6)	0.0296 (5)
H3	0.895987	0.215571	0.375132	0.036*
C4	0.93229 (19)	0.33707 (13)	0.4113 (6)	0.0316 (5)
H4	1.004194	0.331184	0.535045	0.038*
C5	0.89200 (18)	0.41434 (12)	0.3255 (6)	0.0310 (5)
H5	0.936169	0.461141	0.390355	0.037*
C6	0.78780 (18)	0.42272 (12)	0.1462 (6)	0.0296 (5)
H6	0.761062	0.475674	0.087081	0.035*
C11	0.61087 (18)	0.36135 (11)	-0.1461 (6)	0.0275 (5)
H11	0.573609	0.312951	-0.223553	0.033*
C21	0.7290 (2)	0.13337 (11)	0.1388 (7)	0.0319 (5)
H21A	0.668356	0.093725	0.069500	0.048*
H21B	0.736939	0.132217	0.381231	0.048*
H21C	0.805963	0.119570	0.036670	0.048*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O13	0.0346 (7)	0.0199 (6)	0.0389 (9)	-0.0001 (6)	-0.0067 (6)	0.0040 (6)
O21	0.0346 (7)	0.0172 (7)	0.0391 (9)	0.0000 (5)	-0.0025 (7)	0.0018 (6)
N12	0.0300 (8)	0.0215 (8)	0.0279 (9)	0.0012 (6)	0.0017 (7)	-0.0006 (7)
C1	0.0310 (10)	0.0206 (9)	0.0264 (11)	0.0008 (7)	0.0041 (9)	-0.0005 (8)

C2	0.0323 (9)	0.0194 (9)	0.0281 (11)	-0.0011 (7)	0.0048 (9)	-0.0007 (8)
C3	0.0348 (10)	0.0230 (9)	0.0310 (11)	0.0017 (8)	0.0036 (9)	0.0015 (9)
C4	0.0316 (10)	0.0307 (11)	0.0324 (12)	0.0001 (8)	0.0009 (9)	0.0006 (9)
C5	0.0350 (11)	0.0243 (10)	0.0337 (12)	-0.0043 (7)	0.0015 (9)	-0.0025 (10)
C6	0.0362 (11)	0.0203 (9)	0.0322 (11)	0.0003 (8)	0.0023 (9)	-0.0007 (9)
C11	0.0352 (11)	0.0181 (8)	0.0292 (11)	-0.0008 (7)	0.0030 (9)	-0.0011 (8)
C21	0.0390 (11)	0.0158 (9)	0.0408 (12)	0.0028 (7)	-0.0004 (10)	0.0028 (9)

Geometric parameters (Å, °)

O13—N12	1.402 (2)	C3—H3	0.9500
O13—H13	0.8400	C4—C5	1.390 (3)
O21—C2	1.372 (2)	C4—H4	0.9500
O21—C21	1.433 (2)	C5—C6	1.377 (3)
N12—C11	1.280 (3)	C5—H5	0.9500
C1—C6	1.401 (3)	C6—H6	0.9500
C1—C2	1.409 (3)	C11—H11	0.9500
C1—C11	1.456 (3)	C21—H21A	0.9800
C2—C3	1.384 (3)	C21—H21B	0.9800
C3—C4	1.391 (3)	C21—H21C	0.9800
N12—O13—H13	109.5	C6—C5—C4	119.69 (18)
C2—O21—C21	117.07 (16)	C6—C5—H5	120.2
C11—N12—O13	111.27 (15)	C4—C5—H5	120.2
C6—C1—C2	117.80 (19)	C5—C6—C1	121.50 (18)
C6—C1—C11	123.00 (17)	C5—C6—H6	119.3
C2—C1—C11	119.18 (17)	C1—C6—H6	119.3
O21—C2—C3	123.87 (17)	N12—C11—C1	122.16 (18)
O21—C2—C1	115.11 (18)	N12—C11—H11	118.9
C3—C2—C1	121.01 (17)	C1—C11—H11	118.9
C2—C3—C4	119.58 (18)	O21—C21—H21A	109.5
C2—C3—H3	120.2	O21—C21—H21B	109.5
C4—C3—H3	120.2	H21A—C21—H21B	109.5
C5—C4—C3	120.4 (2)	O21—C21—H21C	109.5
C5—C4—H4	119.8	H21A—C21—H21C	109.5
C3—C4—H4	119.8	H21B—C21—H21C	109.5
C21—O21—C2—C3	4.2 (3)	C2—C3—C4—C5	-0.4 (3)
C21—O21—C2—C1	-176.10 (18)	C3—C4—C5—C6	0.0 (3)
C6—C1—C2—O21	-179.70 (18)	C4—C5—C6—C1	0.4 (3)
C11—C1—C2—O21	-1.2 (3)	C2—C1—C6—C5	-0.4 (3)
C6—C1—C2—C3	0.0 (3)	C11—C1—C6—C5	-178.8 (2)
C11—C1—C2—C3	178.5 (2)	O13—N12—C11—C1	179.25 (17)
O21—C2—C3—C4	-179.9 (2)	C6—C1—C11—N12	-6.4 (3)
C1—C2—C3—C4	0.4 (3)	C2—C1—C11—N12	175.2 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O13—H13 \cdots N12 ⁱ	0.84	1.93	2.764 (2)	170
C3—H3 \cdots O13 ⁱⁱ	0.95	2.50	3.442 (2)	174
C21—H21C \cdots O13 ⁱⁱⁱ	0.98	2.57	3.506 (3)	160

Symmetry codes: (i) $-x+1, -y+1, z-1/2$; (ii) $x+1/2, -y+1/2, z+1$; (iii) $x+1/2, -y+1/2, z$.

2,3-Dimethoxy-benzaldehyde oxime (2)

Crystal data

$C_9H_{11}NO_3$

$M_r = 181.19$

Orthorhombic, $P2_12_12_1$

$a = 4.6775$ (2) Å

$b = 13.0996$ (5) Å

$c = 14.1984$ (5) Å

$V = 869.98$ (6) Å³

$Z = 4$

$F(000) = 384$

$D_x = 1.383$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54178$ Å

Cell parameters from 2093 reflections

$\theta = 4.6$ – 67.5°

$\mu = 0.87$ mm⁻¹

$T = 100$ K

Needle, colourless

$0.30 \times 0.05 \times 0.02$ mm

Data collection

Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector diffractometer

Radiation source: Rotating anode, Rigaku 007 HF

Detector resolution: 10 pixels mm⁻¹

profile data from ω -scans

Absorption correction: multi-scan (CrysAlis PRO; Rigaku OD, 2017)

$T_{\min} = 0.507$, $T_{\max} = 1.000$

7835 measured reflections

1596 independent reflections

1371 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.095$

$\theta_{\max} = 68.2^\circ$, $\theta_{\min} = 4.6^\circ$

$h = -5 \rightarrow 5$

$k = -15 \rightarrow 15$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.058$

$wR(F^2) = 0.151$

$S = 1.03$

1596 reflections

124 parameters

0 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.1064P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.35$ e Å⁻³

$\Delta\rho_{\min} = -0.20$ e Å⁻³

Absolute structure: Flack x determined using

474 quotients $[(F^-)-(I)]/[(F^+)+(I)]$ (Parsons *et al.*, 2013)

Absolute structure parameter: 0.2 (3)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O13	−0.3161 (6)	0.81289 (19)	0.42174 (19)	0.0376 (7)
H13	−0.382 (10)	0.799 (3)	0.485 (3)	0.045 (12)*
O21	0.3629 (5)	0.73710 (18)	0.18056 (18)	0.0331 (6)
O31	0.6683 (6)	0.57595 (18)	0.1198 (2)	0.0370 (7)
N12	−0.1250 (7)	0.7305 (2)	0.4137 (2)	0.0328 (7)
C1	0.1671 (8)	0.6379 (3)	0.3072 (3)	0.0311 (8)
C2	0.3440 (8)	0.6450 (2)	0.2283 (3)	0.0309 (8)
C3	0.5135 (8)	0.5622 (2)	0.1997 (3)	0.0315 (8)
C4	0.5069 (9)	0.4725 (3)	0.2525 (2)	0.0331 (9)
H4	0.620619	0.415756	0.234431	0.040*
C5	0.3334 (9)	0.4664 (3)	0.3318 (3)	0.0344 (8)
H5	0.332576	0.405472	0.368039	0.041*
C6	0.1631 (9)	0.5468 (3)	0.3588 (3)	0.0338 (9)
H6	0.042859	0.540487	0.412457	0.041*
C11	−0.0229 (8)	0.7232 (2)	0.3309 (3)	0.0332 (8)
H11	−0.069638	0.772807	0.284541	0.040*
C21	0.2048 (9)	0.7399 (3)	0.0942 (3)	0.0374 (9)
H21A	0.230310	0.806598	0.064099	0.056*
H21B	0.001542	0.728827	0.107410	0.056*
H21C	0.274481	0.686165	0.052004	0.056*
C31	0.8511 (10)	0.4933 (3)	0.0919 (3)	0.0390 (10)
H31A	0.952242	0.511808	0.033893	0.059*
H31B	0.735578	0.432080	0.080781	0.059*
H31C	0.990297	0.479650	0.141905	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O13	0.0411 (16)	0.0292 (13)	0.0423 (16)	0.0088 (12)	0.0053 (13)	0.0027 (11)
O21	0.0339 (14)	0.0250 (12)	0.0405 (14)	−0.0027 (11)	0.0009 (12)	0.0020 (10)
O31	0.0340 (14)	0.0291 (12)	0.0478 (15)	0.0042 (11)	0.0072 (13)	−0.0015 (11)
N12	0.0294 (16)	0.0244 (14)	0.0446 (18)	0.0025 (13)	−0.0001 (15)	−0.0001 (12)
C1	0.0277 (18)	0.0277 (17)	0.038 (2)	−0.0008 (15)	−0.0021 (17)	−0.0004 (14)
C2	0.0297 (17)	0.0244 (17)	0.0387 (19)	−0.0018 (15)	−0.0012 (17)	0.0017 (14)
C3	0.0285 (18)	0.0262 (16)	0.040 (2)	0.0016 (15)	0.0002 (16)	−0.0029 (15)
C4	0.0315 (18)	0.0236 (17)	0.044 (2)	0.0017 (16)	−0.0025 (17)	−0.0036 (14)
C5	0.036 (2)	0.0286 (17)	0.039 (2)	0.0000 (16)	−0.0054 (18)	0.0019 (15)
C6	0.034 (2)	0.0298 (18)	0.038 (2)	0.0001 (17)	−0.0004 (17)	0.0019 (15)
C11	0.0353 (19)	0.0238 (16)	0.040 (2)	0.0010 (17)	0.0001 (18)	0.0007 (15)
C21	0.040 (2)	0.0324 (18)	0.040 (2)	−0.0016 (17)	0.0010 (17)	0.0037 (15)
C31	0.035 (2)	0.0296 (18)	0.052 (2)	0.0034 (16)	0.009 (2)	−0.0068 (16)

Geometric parameters (Å, °)

O13—N12	1.405 (4)	C4—C5	1.389 (5)
O13—H13	0.97 (4)	C4—H4	0.9500
O21—C2	1.386 (4)	C5—C6	1.375 (5)
O21—C21	1.432 (4)	C5—H5	0.9500
O31—C3	1.357 (4)	C6—H6	0.9500
O31—C31	1.435 (4)	C11—H11	0.9500
N12—C11	1.273 (5)	C21—H21A	0.9800
C1—C2	1.396 (5)	C21—H21B	0.9800
C1—C6	1.401 (5)	C21—H21C	0.9800
C1—C11	1.467 (5)	C31—H31A	0.9800
C2—C3	1.404 (5)	C31—H31B	0.9800
C3—C4	1.394 (5)	C31—H31C	0.9800
N12—O13—H13	97 (3)	C5—C6—C1	119.9 (4)
C2—O21—C21	114.1 (3)	C5—C6—H6	120.1
C3—O31—C31	116.6 (3)	C1—C6—H6	120.1
C11—N12—O13	111.8 (3)	N12—C11—C1	119.7 (3)
C2—C1—C6	119.0 (3)	N12—C11—H11	120.1
C2—C1—C11	119.5 (3)	C1—C11—H11	120.1
C6—C1—C11	121.4 (3)	O21—C21—H21A	109.5
O21—C2—C1	119.2 (3)	O21—C21—H21B	109.5
O21—C2—C3	119.7 (3)	H21A—C21—H21B	109.5
C1—C2—C3	121.0 (3)	O21—C21—H21C	109.5
O31—C3—C4	125.0 (3)	H21A—C21—H21C	109.5
O31—C3—C2	116.1 (3)	H21B—C21—H21C	109.5
C4—C3—C2	118.9 (3)	O31—C31—H31A	109.5
C5—C4—C3	119.8 (3)	O31—C31—H31B	109.5
C5—C4—H4	120.1	H31A—C31—H31B	109.5
C3—C4—H4	120.1	O31—C31—H31C	109.5
C6—C5—C4	121.4 (3)	H31A—C31—H31C	109.5
C6—C5—H5	119.3	H31B—C31—H31C	109.5
C4—C5—H5	119.3		
C21—O21—C2—C1	-103.3 (4)	C1—C2—C3—C4	-1.2 (5)
C21—O21—C2—C3	80.0 (4)	O31—C3—C4—C5	-178.1 (4)
C6—C1—C2—O21	-175.9 (3)	C2—C3—C4—C5	0.2 (5)
C11—C1—C2—O21	7.9 (5)	C3—C4—C5—C6	1.1 (6)
C6—C1—C2—C3	0.8 (5)	C4—C5—C6—C1	-1.4 (6)
C11—C1—C2—C3	-175.4 (4)	C2—C1—C6—C5	0.5 (6)
C31—O31—C3—C4	-4.0 (5)	C11—C1—C6—C5	176.6 (3)
C31—O31—C3—C2	177.6 (4)	O13—N12—C11—C1	-176.2 (3)
O21—C2—C3—O31	-6.0 (5)	C2—C1—C11—N12	-161.0 (3)
C1—C2—C3—O31	177.3 (3)	C6—C1—C11—N12	22.8 (6)
O21—C2—C3—C4	175.5 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C1–C6 ring.

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O13—H13 \cdots N12 ⁱ	0.97 (4)	1.87 (5)	2.805 (4)	161 (4)
C4—H4 \cdots O21 ⁱⁱ	0.95	2.63	3.284 (4)	126
C21—H21B \cdots O31 ⁱⁱⁱ	0.98	2.54	3.323 (5)	136
C31—H31B \cdots O13 ^{iv}	0.98	2.51	3.448 (5)	161
C31—H31C \cdots Cg1 ^v	0.98	2.73	3.599 (5)	148

Symmetry codes: (i) $x-1/2, -y+3/2, -z+1$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $x-1, y, z$; (iv) $-x, y-1/2, -z+1/2$; (v) $x+1, y, z$.**2,4-Dimethoxybenzaldehyde oxime** (3)*Crystal data*C₉H₁₁NO₃ $M_r = 181.19$ Triclinic, $P\bar{1}$ $a = 4.9441$ (2) \AA $b = 8.2188$ (4) \AA $c = 12.1308$ (3) \AA $\alpha = 108.849$ (3) $^\circ$ $\beta = 92.288$ (3) $^\circ$ $\gamma = 106.273$ (4) $^\circ$ $V = 443.17$ (3) \AA^3 $Z = 2$ $F(000) = 192$ $D_x = 1.358$ Mg m⁻³Cu $K\alpha$ radiation, $\lambda = 1.54178$ \AA

Cell parameters from 3758 reflections

 $\theta = 3.9\text{--}69.9^\circ$ $\mu = 0.86$ mm⁻¹ $T = 100$ K

Block, colourless

 $0.20 \times 0.10 \times 0.05$ mm*Data collection*

Rigaku 007HF equipped with Varimax confocal mirrors and an AFC11 goniometer and HyPix 6000 detector diffractometer

Radiation source: Rotating anode, Rigaku 007 HF

Detector resolution: 10 pixels mm⁻¹profile data from ω -scans

Absorption correction: multi-scan

(CrysAlis PRO; Rigaku OD, 2017)

 $T_{\min} = 0.802, T_{\max} = 1.000$

7618 measured reflections

1594 independent reflections

1462 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.033$ $\theta_{\max} = 68.2^\circ, \theta_{\min} = 3.9^\circ$ $h = -5 \rightarrow 5$ $k = -9 \rightarrow 9$ $l = -14 \rightarrow 13$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.101$ $S = 0.88$

1594 reflections

124 parameters

0 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0775P)^2 + 0.1273P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.20$ e \AA^{-3} $\Delta\rho_{\min} = -0.19$ e \AA^{-3} *Special details***Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O13	-0.22417 (16)	0.06653 (12)	0.08642 (7)	0.0311 (2)
O21	0.29524 (16)	0.56560 (10)	0.41358 (7)	0.0261 (2)
O41	1.14895 (16)	0.88605 (11)	0.31075 (7)	0.0273 (2)
N12	0.05891 (19)	0.17759 (13)	0.09919 (8)	0.0255 (3)
C1	0.3951 (2)	0.45758 (15)	0.22073 (10)	0.0231 (3)
C2	0.4791 (2)	0.58860 (15)	0.33490 (9)	0.0228 (3)
C3	0.7331 (2)	0.72870 (15)	0.36234 (9)	0.0237 (3)
H3	0.789071	0.815115	0.439846	0.028*
C4	0.9063 (2)	0.74213 (15)	0.27541 (10)	0.0238 (3)
C5	0.8276 (2)	0.61490 (15)	0.16177 (10)	0.0247 (3)
H5	0.945380	0.624026	0.102665	0.030*
C6	0.5741 (2)	0.47475 (15)	0.13657 (10)	0.0245 (3)
H6	0.520758	0.387495	0.059294	0.029*
C11	0.1194 (2)	0.31641 (15)	0.19252 (10)	0.0242 (3)
H11	-0.017316	0.328624	0.244952	0.029*
C21	0.3513 (2)	0.70704 (15)	0.52560 (9)	0.0281 (3)
H21A	0.196509	0.679099	0.571095	0.042*
H21B	0.364539	0.821605	0.514202	0.042*
H21C	0.531398	0.717369	0.568289	0.042*
C41	1.3260 (2)	0.90933 (16)	0.22272 (10)	0.0278 (3)
H41A	1.490351	1.018155	0.257704	0.042*
H41B	1.216391	0.922054	0.158580	0.042*
H41C	1.392155	0.803696	0.191690	0.042*
H13	-0.235 (4)	-0.031 (2)	0.0244 (16)	0.049 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O13	0.0234 (4)	0.0320 (5)	0.0287 (5)	-0.0008 (3)	0.0076 (3)	0.0062 (4)
O21	0.0251 (4)	0.0298 (4)	0.0202 (4)	0.0037 (3)	0.0072 (3)	0.0081 (3)
O41	0.0227 (4)	0.0309 (5)	0.0241 (4)	0.0022 (3)	0.0068 (3)	0.0089 (3)
N12	0.0212 (5)	0.0278 (5)	0.0252 (5)	0.0032 (4)	0.0044 (4)	0.0100 (4)
C1	0.0229 (6)	0.0248 (6)	0.0230 (6)	0.0079 (4)	0.0033 (4)	0.0099 (4)
C2	0.0222 (5)	0.0285 (6)	0.0216 (6)	0.0092 (4)	0.0060 (4)	0.0123 (5)
C3	0.0235 (6)	0.0279 (6)	0.0196 (5)	0.0072 (5)	0.0034 (4)	0.0086 (4)
C4	0.0198 (6)	0.0269 (6)	0.0257 (6)	0.0063 (4)	0.0029 (4)	0.0114 (5)
C5	0.0236 (6)	0.0306 (6)	0.0229 (6)	0.0100 (5)	0.0077 (4)	0.0113 (5)
C6	0.0252 (6)	0.0271 (6)	0.0208 (5)	0.0083 (5)	0.0036 (4)	0.0077 (4)
C11	0.0242 (6)	0.0288 (6)	0.0221 (6)	0.0080 (5)	0.0058 (4)	0.0118 (4)
C21	0.0295 (6)	0.0324 (6)	0.0195 (6)	0.0065 (5)	0.0077 (4)	0.0075 (5)
C41	0.0250 (6)	0.0333 (6)	0.0258 (6)	0.0059 (5)	0.0085 (4)	0.0131 (5)

Geometric parameters (Å, °)

O13—N12	1.4112 (12)	C3—H3	0.9500
O13—H13	0.893 (18)	C4—C5	1.3939 (16)
O21—C2	1.3648 (13)	C5—C6	1.3869 (16)
O21—C21	1.4294 (13)	C5—H5	0.9500
O41—C4	1.3632 (13)	C6—H6	0.9500
O41—C41	1.4339 (13)	C11—H11	0.9500
N12—C11	1.2728 (15)	C21—H21A	0.9800
C1—C6	1.3931 (16)	C21—H21B	0.9800
C1—C2	1.4107 (15)	C21—H21C	0.9800
C1—C11	1.4634 (15)	C41—H41A	0.9800
C2—C3	1.3864 (16)	C41—H41B	0.9800
C3—C4	1.3971 (16)	C41—H41C	0.9800
N12—O13—H13	103.5 (11)	C1—C6—C5	122.23 (10)
C2—O21—C21	117.44 (8)	C1—C6—H6	118.9
C4—O41—C41	116.82 (9)	C5—C6—H6	118.9
C11—N12—O13	111.40 (9)	N12—C11—C1	121.37 (10)
C6—C1—C2	117.84 (10)	N12—C11—H11	119.3
C6—C1—C11	122.31 (10)	C1—C11—H11	119.3
C2—C1—C11	119.74 (10)	O21—C21—H21A	109.5
O21—C2—C3	123.79 (10)	O21—C21—H21B	109.5
O21—C2—C1	115.34 (10)	H21A—C21—H21B	109.5
C3—C2—C1	120.88 (10)	O21—C21—H21C	109.5
C2—C3—C4	119.64 (10)	H21A—C21—H21C	109.5
C2—C3—H3	120.2	H21B—C21—H21C	109.5
C4—C3—H3	120.2	O41—C41—H41A	109.5
O41—C4—C3	115.11 (10)	O41—C41—H41B	109.5
O41—C4—C5	124.27 (10)	H41A—C41—H41B	109.5
C3—C4—C5	120.62 (10)	O41—C41—H41C	109.5
C6—C5—C4	118.78 (10)	H41A—C41—H41C	109.5
C6—C5—H5	120.6	H41B—C41—H41C	109.5
C4—C5—H5	120.6		
C21—O21—C2—C3	8.36 (15)	C2—C3—C4—O41	179.03 (9)
C21—O21—C2—C1	-171.96 (9)	C2—C3—C4—C5	-0.70 (17)
C6—C1—C2—O21	179.72 (9)	O41—C4—C5—C6	-179.69 (9)
C11—C1—C2—O21	3.38 (15)	C3—C4—C5—C6	0.01 (17)
C6—C1—C2—C3	-0.59 (16)	C2—C1—C6—C5	-0.11 (17)
C11—C1—C2—C3	-176.93 (9)	C11—C1—C6—C5	176.12 (9)
O21—C2—C3—C4	-179.34 (9)	C4—C5—C6—C1	0.40 (17)
C1—C2—C3—C4	1.00 (16)	O13—N12—C11—C1	-175.95 (9)
C41—O41—C4—C3	-177.21 (9)	C6—C1—C11—N12	17.32 (17)
C41—O41—C4—C5	2.51 (16)	C2—C1—C11—N12	-166.51 (10)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C1–C6 ring.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O13—H13···N12 ⁱ	0.893 (18)	1.995 (19)	2.8124 (13)	151.5 (15)
C41—H41A···O13 ⁱⁱ	0.98	2.63	3.0680 (15)	107
C41—H41C···Cg1 ⁱⁱⁱ	0.98	2.60	3.4479 (13)	144

Symmetry codes: (i) $-x, -y, -z$; (ii) $x+2, y+1, z$; (iii) $x+1, y, z$.**2,5-Dimethoxybenzaldehyde oxime (4)***Crystal data*C₉H₁₁NO₃ $M_r = 181.19$ Monoclinic, $P2_1/c$ $a = 7.6480$ (1) Å $b = 21.3380$ (4) Å $c = 10.9421$ (2) Å $\beta = 90.555$ (2)° $V = 1785.59$ (5) Å³ $Z = 8$ $F(000) = 768$ $D_x = 1.348$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71075$ Å

Cell parameters from 21005 reflections

 $\theta = 2.1$ – 32.1 ° $\mu = 0.10$ mm⁻¹ $T = 100$ K

Block, colourless

 $0.20 \times 0.15 \times 0.13$ mm*Data collection*

Rigaku FRE+ equipped with VHF Varimax confocal mirrors and an AFC12 goniometer and HyPix 6000 detector diffractometer

Radiation source: Rotating Anode, Rigaku FRE+

Confocal mirrors, VHF Varimax monochromator

Detector resolution: 10 pixels mm⁻¹ profile data from ω -scans

Absorption correction: multi-scan (CrysAlis PRO; Rigaku OD, 2017)

 $T_{\min} = 0.935$, $T_{\max} = 1.000$

38753 measured reflections

4082 independent reflections

3761 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.020$ $\theta_{\max} = 27.5$ °, $\theta_{\min} = 1.9$ ° $h = -9 \rightarrow 9$ $k = -27 \rightarrow 27$ $l = -14 \rightarrow 14$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.031$ $wR(F^2) = 0.086$ $S = 1.06$

4082 reflections

247 parameters

0 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0477P)^2 + 0.5056P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.32$ e Å⁻³ $\Delta\rho_{\min} = -0.19$ e Å⁻³*Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O121	1.05737 (9)	0.46081 (3)	0.23188 (6)	0.01640 (15)
O221	0.33772 (9)	0.09728 (3)	0.32374 (6)	0.01705 (15)
O151	0.88735 (9)	0.21358 (3)	0.13343 (6)	0.01629 (14)
O213	0.63115 (9)	0.03239 (3)	0.60782 (6)	0.01852 (15)
H213	0.6028 (19)	-0.0064 (7)	0.5898 (13)	0.036 (4)*
O251	0.47197 (9)	0.34248 (3)	0.46029 (6)	0.01810 (15)
O113	0.84907 (9)	0.45697 (3)	0.56752 (6)	0.01890 (15)
H113	0.905 (2)	0.4914 (7)	0.5868 (13)	0.036 (4)*
N112	0.92134 (10)	0.44408 (4)	0.45300 (7)	0.01543 (16)
N212	0.53640 (10)	0.06589 (4)	0.51894 (7)	0.01507 (16)
C11	0.92057 (11)	0.36690 (4)	0.29191 (8)	0.01265 (17)
C12	1.01564 (11)	0.40037 (4)	0.20387 (8)	0.01345 (17)
C13	1.06269 (12)	0.37068 (4)	0.09534 (8)	0.01520 (18)
H13	1.125790	0.393201	0.035247	0.018*
C14	1.01792 (11)	0.30839 (4)	0.07452 (8)	0.01529 (18)
H14	1.051935	0.288508	0.000789	0.018*
C15	0.92359 (11)	0.27498 (4)	0.16103 (8)	0.01354 (17)
C16	0.87441 (11)	0.30442 (4)	0.26854 (8)	0.01321 (17)
H16	0.808499	0.281887	0.327125	0.016*
C21	0.47308 (11)	0.17198 (4)	0.45403 (8)	0.01312 (17)
C22	0.36691 (11)	0.15892 (4)	0.35049 (8)	0.01327 (17)
C23	0.29914 (11)	0.20820 (4)	0.28236 (8)	0.01541 (18)
H23	0.229476	0.199464	0.212053	0.018*
C24	0.33136 (12)	0.27053 (4)	0.31516 (8)	0.01597 (18)
H24	0.284202	0.303707	0.267240	0.019*
C25	0.43233 (11)	0.28369 (4)	0.41782 (8)	0.01448 (18)
C26	0.50301 (11)	0.23437 (4)	0.48551 (8)	0.01430 (17)
H26	0.573634	0.243537	0.555138	0.017*
C111	0.86582 (11)	0.39220 (4)	0.40987 (8)	0.01383 (17)
H111	0.784786	0.368751	0.456649	0.017*
C121	1.16522 (14)	0.49466 (5)	0.14906 (9)	0.0233 (2)
H12A	1.188699	0.536566	0.181970	0.035*
H12B	1.105118	0.498388	0.069891	0.035*
H12C	1.275896	0.472247	0.138464	0.035*
C151	0.79088 (13)	0.17920 (4)	0.22231 (9)	0.01864 (19)
H15A	0.769266	0.136629	0.192162	0.028*
H15B	0.679003	0.200181	0.236846	0.028*
H15C	0.858216	0.177225	0.298868	0.028*
C211	0.55768 (11)	0.12487 (4)	0.53198 (8)	0.01470 (18)
H211	0.632715	0.139016	0.595943	0.018*
C221	0.24158 (15)	0.08339 (5)	0.21389 (9)	0.0238 (2)
H22A	0.233090	0.037865	0.203679	0.036*
H22B	0.302042	0.101530	0.143698	0.036*
H22C	0.123897	0.101296	0.219223	0.036*
C251	0.41248 (14)	0.39453 (4)	0.38939 (9)	0.0225 (2)

H25A	0.453866	0.433561	0.426991	0.034*
H25B	0.284384	0.394557	0.386171	0.034*
H25C	0.458345	0.391238	0.306319	0.034*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O121	0.0203 (3)	0.0116 (3)	0.0174 (3)	-0.0029 (2)	0.0052 (2)	-0.0001 (2)
O221	0.0220 (3)	0.0133 (3)	0.0157 (3)	-0.0018 (2)	-0.0063 (2)	-0.0012 (2)
O151	0.0188 (3)	0.0128 (3)	0.0173 (3)	-0.0015 (2)	0.0015 (2)	-0.0033 (2)
O213	0.0230 (3)	0.0136 (3)	0.0189 (3)	0.0017 (3)	-0.0077 (3)	0.0021 (2)
O251	0.0208 (3)	0.0111 (3)	0.0223 (3)	-0.0006 (2)	-0.0048 (3)	0.0004 (2)
O113	0.0263 (4)	0.0160 (3)	0.0145 (3)	-0.0053 (3)	0.0077 (3)	-0.0043 (2)
N112	0.0186 (4)	0.0148 (4)	0.0129 (3)	-0.0001 (3)	0.0039 (3)	-0.0015 (3)
N212	0.0165 (4)	0.0146 (4)	0.0140 (4)	0.0020 (3)	-0.0027 (3)	0.0020 (3)
C11	0.0114 (4)	0.0128 (4)	0.0137 (4)	0.0010 (3)	-0.0013 (3)	-0.0004 (3)
C12	0.0123 (4)	0.0122 (4)	0.0158 (4)	0.0007 (3)	-0.0006 (3)	0.0003 (3)
C13	0.0151 (4)	0.0163 (4)	0.0142 (4)	0.0007 (3)	0.0015 (3)	0.0019 (3)
C14	0.0152 (4)	0.0173 (4)	0.0133 (4)	0.0021 (3)	0.0003 (3)	-0.0024 (3)
C15	0.0116 (4)	0.0126 (4)	0.0164 (4)	0.0013 (3)	-0.0030 (3)	-0.0015 (3)
C16	0.0117 (4)	0.0133 (4)	0.0146 (4)	0.0001 (3)	-0.0004 (3)	0.0010 (3)
C21	0.0125 (4)	0.0141 (4)	0.0128 (4)	-0.0005 (3)	0.0012 (3)	0.0007 (3)
C22	0.0126 (4)	0.0133 (4)	0.0139 (4)	-0.0012 (3)	0.0012 (3)	-0.0007 (3)
C23	0.0140 (4)	0.0175 (4)	0.0147 (4)	0.0001 (3)	-0.0017 (3)	-0.0001 (3)
C24	0.0145 (4)	0.0153 (4)	0.0180 (4)	0.0020 (3)	-0.0010 (3)	0.0026 (3)
C25	0.0128 (4)	0.0127 (4)	0.0179 (4)	-0.0009 (3)	0.0017 (3)	-0.0003 (3)
C26	0.0133 (4)	0.0159 (4)	0.0137 (4)	-0.0010 (3)	-0.0008 (3)	-0.0002 (3)
C111	0.0139 (4)	0.0135 (4)	0.0141 (4)	-0.0004 (3)	0.0016 (3)	0.0010 (3)
C121	0.0312 (5)	0.0160 (4)	0.0228 (5)	-0.0063 (4)	0.0096 (4)	0.0015 (4)
C151	0.0245 (5)	0.0131 (4)	0.0183 (4)	-0.0024 (3)	0.0002 (4)	-0.0005 (3)
C211	0.0145 (4)	0.0160 (4)	0.0135 (4)	-0.0011 (3)	-0.0018 (3)	-0.0005 (3)
C221	0.0321 (5)	0.0188 (4)	0.0204 (5)	-0.0043 (4)	-0.0119 (4)	-0.0020 (4)
C251	0.0280 (5)	0.0131 (4)	0.0263 (5)	0.0018 (4)	-0.0035 (4)	0.0027 (4)

Geometric parameters (Å, °)

O121—C12	1.3628 (10)	C21—C26	1.3935 (12)
O121—C121	1.4275 (11)	C21—C22	1.4151 (12)
O221—C22	1.3653 (10)	C21—C211	1.4651 (12)
O221—C221	1.4339 (11)	C22—C23	1.3866 (12)
O151—C15	1.3721 (10)	C23—C24	1.3988 (12)
O151—C151	1.4291 (11)	C23—H23	0.9500
O213—N212	1.4029 (9)	C24—C25	1.3858 (12)
O213—H213	0.877 (15)	C24—H24	0.9500
O251—C25	1.3706 (10)	C25—C26	1.3929 (12)
O251—C251	1.4268 (11)	C26—H26	0.9500
O113—N112	1.4017 (9)	C111—H111	0.9500
O113—H113	0.875 (16)	C121—H12A	0.9800

N112—C111	1.2747 (11)	C121—H12B	0.9800
N212—C211	1.2767 (12)	C121—H12C	0.9800
C11—C16	1.4020 (12)	C151—H15A	0.9800
C11—C12	1.4072 (12)	C151—H15B	0.9800
C11—C111	1.4639 (12)	C151—H15C	0.9800
C12—C13	1.3962 (12)	C211—H211	0.9500
C13—C14	1.3908 (12)	C221—H22A	0.9800
C13—H13	0.9500	C221—H22B	0.9800
C14—C15	1.3921 (12)	C221—H22C	0.9800
C14—H14	0.9500	C251—H25A	0.9800
C15—C16	1.3887 (12)	C251—H25B	0.9800
C16—H16	0.9500	C251—H25C	0.9800
C12—O121—C121	118.13 (7)	C23—C24—H24	120.1
C22—O221—C221	117.40 (7)	O251—C25—C24	125.45 (8)
C15—O151—C151	116.42 (7)	O251—C25—C26	115.32 (8)
N212—O213—H213	101.5 (10)	C24—C25—C26	119.23 (8)
C25—O251—C251	117.38 (7)	C25—C26—C21	121.89 (8)
N112—O113—H113	100.6 (10)	C25—C26—H26	119.1
C111—N112—O113	111.63 (7)	C21—C26—H26	119.1
C211—N212—O213	111.12 (7)	N112—C111—C11	123.34 (8)
C16—C11—C12	119.24 (8)	N112—C111—H111	118.3
C16—C11—C111	115.96 (8)	C11—C111—H111	118.3
C12—C11—C111	124.79 (8)	O121—C121—H12A	109.5
O121—C12—C13	123.99 (8)	O121—C121—H12B	109.5
O121—C12—C11	116.60 (8)	H12A—C121—H12B	109.5
C13—C12—C11	119.41 (8)	O121—C121—H12C	109.5
C14—C13—C12	120.53 (8)	H12A—C121—H12C	109.5
C14—C13—H13	119.7	H12B—C121—H12C	109.5
C12—C13—H13	119.7	O151—C151—H15A	109.5
C13—C14—C15	120.42 (8)	O151—C151—H15B	109.5
C13—C14—H14	119.8	H15A—C151—H15B	109.5
C15—C14—H14	119.8	O151—C151—H15C	109.5
O151—C15—C16	124.23 (8)	H15A—C151—H15C	109.5
O151—C15—C14	116.36 (8)	H15B—C151—H15C	109.5
C16—C15—C14	119.40 (8)	N212—C211—C21	123.80 (8)
C15—C16—C11	120.99 (8)	N212—C211—H211	118.1
C15—C16—H16	119.5	C21—C211—H211	118.1
C11—C16—H16	119.5	O221—C221—H22A	109.5
C26—C21—C22	118.53 (8)	O221—C221—H22B	109.5
C26—C21—C211	116.18 (8)	H22A—C221—H22B	109.5
C22—C21—C211	125.29 (8)	O221—C221—H22C	109.5
O221—C22—C23	123.76 (8)	H22A—C221—H22C	109.5
O221—C22—C21	116.92 (8)	H22B—C221—H22C	109.5
C23—C22—C21	119.32 (8)	O251—C251—H25A	109.5
C22—C23—C24	121.27 (8)	O251—C251—H25B	109.5
C22—C23—H23	119.4	H25A—C251—H25B	109.5
C24—C23—H23	119.4	O251—C251—H25C	109.5

C25—C24—C23	119.74 (8)	H25A—C251—H25C	109.5
C25—C24—H24	120.1	H25B—C251—H25C	109.5
C121—O121—C12—C13	-4.21 (13)	C211—C21—C22—O221	1.91 (13)
C121—O121—C12—C11	175.24 (8)	C26—C21—C22—C23	1.31 (12)
C16—C11—C12—O121	-179.74 (7)	C211—C21—C22—C23	-177.99 (8)
C111—C11—C12—O121	-0.50 (13)	O221—C22—C23—C24	179.06 (8)
C16—C11—C12—C13	-0.26 (13)	C21—C22—C23—C24	-1.05 (13)
C111—C11—C12—C13	178.98 (8)	C22—C23—C24—C25	-0.22 (13)
O121—C12—C13—C14	178.77 (8)	C251—O251—C25—C24	-3.63 (13)
C11—C12—C13—C14	-0.67 (13)	C251—O251—C25—C26	175.86 (8)
C12—C13—C14—C15	0.80 (13)	C23—C24—C25—O251	-179.32 (8)
C151—O151—C15—C16	0.74 (12)	C23—C24—C25—C26	1.21 (13)
C151—O151—C15—C14	179.85 (8)	O251—C25—C26—C21	179.53 (8)
C13—C14—C15—O151	-179.15 (8)	C24—C25—C26—C21	-0.94 (13)
C13—C14—C15—C16	0.01 (13)	C22—C21—C26—C25	-0.32 (13)
O151—C15—C16—C11	178.13 (8)	C211—C21—C26—C25	179.03 (8)
C14—C15—C16—C11	-0.95 (13)	O113—N112—C111—C11	-179.60 (8)
C12—C11—C16—C15	1.08 (13)	C16—C11—C111—N112	168.18 (8)
C111—C11—C16—C15	-178.23 (8)	C12—C11—C111—N112	-11.08 (14)
C221—O221—C22—C23	4.54 (13)	O213—N212—C211—C21	-178.99 (8)
C221—O221—C22—C21	-175.35 (8)	C26—C21—C211—N212	176.46 (8)
C26—C21—C22—O221	-178.79 (8)	C22—C21—C211—N212	-4.24 (14)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C11—C16 and C21—C26 rings, respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O113—H113...O121 ⁱ	0.875 (16)	2.247 (15)	2.8944 (9)	130.7 (12)
O113—H113...N112 ⁱ	0.875 (16)	1.965 (16)	2.7567 (10)	149.9 (13)
O213—H213...O221 ⁱⁱ	0.877 (15)	2.204 (15)	2.8758 (9)	133.1 (12)
O213—H213...N212 ⁱⁱ	0.877 (15)	2.034 (15)	2.8160 (10)	147.9 (13)
C111—H111...O251	0.95	2.46	3.2458 (11)	140
C121—H12C...N212 ⁱⁱⁱ	0.98	2.53	3.4400 (13)	155
C151—H15A...O113 ^{iv}	0.98	2.50	3.3947 (11)	152
C14—H14...Cg2 ⁱⁱⁱ	0.95	2.98	3.6656 (9)	130
C151—H15B...Cg2	0.98	2.72	3.5973 (10)	149
C24—H24...Cg1 ^v	0.95	2.67	3.4281 (10)	137
C211—H211...Cg1 ^{vi}	0.95	2.78	3.6272 (9)	149

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$; (iii) $x+1, -y+1/2, z-1/2$; (iv) $x, -y+1/2, z-1/2$; (v) $x-1, y, z$; (vi) $x, -y+1/2, z+1/2$.