

Crystal structure of a hydrogen-bonded 2:1 co-crystal of 4-nitrophenol and 4,4'-bipyridine


Angela Gotingco, Merary Villanueva Contreras, Jeanette A. Wolfarth, S. Chantal E. Stieber and Zoe Y. Marr*

Department of Chemistry & Biochemistry, California State Polytechnic University, Pomona, 3801 W. Temple Ave., Pomona, CA 91768, USA. *Correspondence e-mail: zymarr@cpp.edu

In the title compound, $C_{10}H_8N_2 \cdot 2C_6H_5NO_3$, 4-nitrophenol and 4,4'-bipyridine crystallized together in a 2:1 ratio in the space group $P2_1/n$. There is a hydrogen-bonding interaction between the nitrogen atoms on the 4,4'-bipyridine molecule and the hydrogen atom on the hydroxyl group on the 4-nitrophenol, resulting in trimolecular units. This structure is a polymorph of a previously reported structure [Nayak & Pedireddi (2016). *Cryst. Growth Des.* **16**, 5966–5975], which differs mainly due to a twist in the 4,4'-bipyridine molecule.

1. Chemical context

Co-crystals are a growing field of science as crystalline solids can be engineered to have improved physical-chemical properties such as better solubility, stability, and bioavailability (Karimi-Jafari *et al.*, 2018). Co-crystals are defined as crystalline solids composed of two or more different molecular and/or ionic compounds in a specific stoichiometric ratio and are neither solvates nor simple salts (Aitipamula *et al.*, 2012). They are held together by intermolecular interactions such as hydrogen bonding, halogen bonding, and π - π stacking (Wang *et al.*, 2022). Generally, co-crystals are high yielding, making them an appealing candidate for crystal engineering in the realm of pharmaceutical purposes (Chettri *et al.*, 2024).



The chemicals used in this co-crystal consist of 4,4'-bipyridine and 4-nitrophenol. 4,4'-Bipyridine can be found in a multitude of crystal structures due the pyridyl groups being suitable for both coordination polymers and co-crystals (Richard *et al.*, 2021). 4-Nitrophenol is commonly found as a drug manufacturing and synthesis intermediate in the pharmaceutical industry. Specifically, this compound has been used in the production of compounds such as acetaminophen, a drug used for pain relief, where it is nitrated and converted to 4-aminophenol, an intermediate for acetaminophen (Abdollahi *et al.*, 2014). This is a highly appealing synthesis process for its greener approach to science in comparison to other intermediates used. Additionally, since 4-nitrophenol takes on the color of a yellow crystalline solid, it is also a suitable candidate that is used to produce pigments/dyes such as leather darkener (National Center for Biotechnology Information, 2024).




Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O3-H3\cdots N1^i$	0.86 (3)	1.84 (3)	2.6921 (19)	174 (3)

Symmetry code: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

2. Structural commentary

4,4'-Bipyridine and 4-nitrophenol co-crystallized in a 1:2 ratio with the asymmetric unit containing one molecule of 4-nitrophenol and half of a 4,4'-bipyridine molecule in the $P2_1/n$ space group. Half of the atoms from the 4,4'-bipyridine molecule sit on Wyckoff position 4e, and the other half is generated by the center of inversion (0, 1/2, 0) (Fig. 1). The nitro groups exhibit a trigonal-planar geometry with bond angles of 122.44 (16)° for $O1-N2-O2$, 118.51 (14)° for $O1-N2-C6$, and 119.06 (15)° for $O2-N2-C6$. The $N-O/N=O$ bonds have lengths of 1.232 (2) and 1.2346 (19) Å, which are in between the average values for $N-O$ and $N=O$ bonds, as expected due to resonance. The molecular geometry of the hydroxyl group is bent with with an angle of 110.4 (19)° for $H3-O3-C9$. The aromatic benzene ring has bond angles ranging between 119.32 (15) and 121.33 (15)°, as expected for sp^2 -hybridized carbons.

3. Supramolecular features

In the structure, each nitrogen on the 4,4'-bipyridine is hydrogen bonded to the hydroxyl group from the 4-nitrophenol, which results in the formation of trimolecular units that propagate along [001] (Fig. 2). There is a hydrogen-bonding interaction (Table 1) between the two molecules with a $D\cdots A$ distance between H3 and N1 of 1.84 (3) Å. In addition to hydrogen bonding, there are also $\pi-\pi$ interactions




Figure 1
A view of the structure containing 4,4'-bipyridine and 4-nitrophenol showing the atom-labeling scheme. Atoms that contain $_a$ in the label are symmetry generated ($-x + 2, -y + 1, -z$). The displacement ellipsoids are drawn at 50% probability.

Table 2
Comparison of unit-cell parameters (Å, °).

Parameter	Nayak & Pedireddi (2016)	Title compound
Crystal system	Monoclinic	Monoclinic
Space group	$P2_1/c$	$P2_1/n$
a	19.090 (4)	12.3711 (7)
b	3.8080 (10)	3.8255 (2)
c	27.3470 (10)	21.4175 (12)
β	98.38 (3)	104.195 (2)

between the ring systems of the adjacent 4,4'-bipyridine molecules having a plane centroid to plane centroid distance of 3.8255 (11) Å.

4. Database survey

The crystal structure that is reported herein is a polymorph of a co-crystal (refcode AWEVUV) published by Nayak & Pedireddi (2016). Similar synthesis and crystallization conditions were used but there are slight differences in the structure. The asymmetric unit of the previous structure contains three molecules, two 4-nitrophenol and one 4,4'-bipyridine, and the 4,4'-bipyridine does not lie on a symmetry operation. In the previously published structure, the pyridyl groups are rotated about the $C13-C22$ bond with a plane twist angle of 24.60 (8)° whereas the pyridyl rings sit in plane with one another in the structure reported above. The unit-cell parameters and space group for the two structures also differ (Table 2).

5. Synthesis and crystallization

The synthesis for the newly reported co-crystal is modified from the procedure published by Nayak & Pedireddi (2016) with a 2:1 rather than a 1:1 ratio of 4,4'-bipyridine and 4-nitrophenol used. In addition, the method of heating was changed from a warm water bath to gentle heating directly on a hot plate. These differences in the synthesis could contribute to the deviation of the packing of the molecules from the original structure.




Figure 2
Diagram of the packed unit cell highlighting the hydrogen-bonding interaction between the hydrogen on the 4-nitrophenol and the nitrogen atom on the 4,4'-bipyridine. View down [010].

A 2:1 molar ratio of 4,4'-bipyridine and 4-nitrophenol was used to synthesize the new co-crystal. 20.0 mg (0.128 mmol, 2 eq) of 4,4'-bipyridine and 8.9 mg (0.0640 mmol, 1 eq) of 4-nitrophenol were added to a 20 mL scintillation vial. 4.4 mL of methanol was added to the vial and then the solution was warmed up on a hot plate to dissolve the solids. Once fully dissolved, the solution was cooled to room temperature. The sample underwent slow evaporation and to control evaporation rate, small holes were punctured on the parafilm covering as the solution was left to evaporate for 2 weeks. The resulting crystals were clear, colorless prisms. These crystals were grown as part of class CHM 5720 'Current advances in Inorganic Chemistry – Introduction to Crystallography' at Cal Poly Pomona.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. All hydrogens except H3 were placed at calculated positions using AFIX commands and refined using a riding model. The position of H3 was determined using the Fourier difference map and refined freely.

Funding information

Funding for this research was provided by: Camille and Henry Dreyfus Foundation (award to S. Chantal E. Stieber); Office of Postsecondary Education (grant No. P031C210068 to Zoe Y. Marr, S. Chantal E. Stieber); National Science Foundation, Directorate for Mathematical and Physical Sciences (grant No. 1847926 to S. Chantal E. Stieber); U.S. Department of Defense, U.S. Army (award No. W911NF-17-1-0537 to S. Chantal E. Stieber).

References

- Abdollahi, M. & Mohammadirad, A. (2014). *Encyclopedia of Toxicology*, **3**, 575–577.
- Aitipamula, S., Banerjee, R., Bansal, A. K., Biradha, K., Cheney, M. L., Choudhury, A. R., Desiraju, G. R., Dikundwar, A. G., Dubey, R., Duggirala, N., Ghogale, P. P., Ghosh, S., Goswami, P. K., Goud, N. R., Jetti, R. R. K. R., Karpinski, P., Kaushik, P., Kumar, D., Kumar, V., Moulton, B., Mukherjee, A., Mukherjee, G., Myerson, A. S., Puri, V., Ramanan, A., Rajamannar, T., Reddy, C. M., Rodriguez-Hornedo, N., Rogers, R. D., Row, T. N. G., Sanphui, P., Shan, N., Shete, G., Singh, A., Sun, C. C., Swift, J. A., Thaimattam, R., Thakur, T. S., Kumar Thaper, R., Thomas, S. P., Tothadi, S., Vangala, V. R., Variankaval, N., Vishweshwar, P., Weyna, D. R. & Zaworotko, M. J. (2012). *Cryst. Growth Des.* **12**, 2147–2152.
- Bruker (2018). *APEX4* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chettri, A., Subba, A., Singh, G. P. & Bag, P. P. (2024). *J. Pharm. Pharmacol.* **76**, 1–12.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Karimi-Jafari, M., Padrela, L., Walker, G. M. & Croker, D. M. (2018). *Cryst. Growth Des.* **18**, 6370–6387.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.
- National Center for Biotechnology Information (2024). PubChem Compound Summary for CID 980, 4-nitrophenol. Retrieved August 22, 2024 from <https://pubchem.ncbi.nlm.nih.gov/compound/4-Nitrophenol>.
- Nayak, A. & Pedireddi, V. R. (2016). *Cryst. Growth Des.* **16**, 5966–5975.
- Richard, J., Joseph, J., Wang, C., Ciesielski, A., Weiss, J., Samorì, P., Mamane, V. & Wytko, J. A. (2021). *J. Org. Chem.* **86**, 3356–3366.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Wang, Y., Wang, L., Zhang, F., Wang, N., Gao, Y., Xiao, Y., Wang, Z. & Bao, Y. (2022). *J. Mol. Struct.* **1258**, 132665.

Table 3

Experimental details.

Crystal data	
Chemical formula	C ₁₀ H ₈ N ₂ ·2C ₆ H ₅ NO ₃
<i>M_r</i>	434.40
Crystal system, space group	Monoclinic, <i>P</i> ₂ ₁ / <i>n</i>
Temperature (K)	123
<i>a</i> , <i>b</i> , <i>c</i> (Å)	12.3711 (7), 3.8255 (2), 21.4175 (12)
β (°)	104.195 (2)
<i>V</i> (Å ³)	982.65 (9)
<i>Z</i>	2
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.11
Crystal size (mm)	0.47 × 0.07 × 0.03
Data collection	
Diffractometer	Bruker D8 Venture
Absorption correction	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)
<i>T_{min}</i> , <i>T_{max}</i>	0.501, 0.746
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	24119, 2957, 2302
<i>R_{int}</i>	0.076
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.711
Refinement	
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.057, 0.161, 1.10
No. of reflections	2957
No. of parameters	149
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	0.47, -0.37

Computer programs: *APEX4* and *SAINT* V8.40B (Bruker, 2018), *SHELXT2018/2* (Sheldrick, 2015a), *SHELXL* (Sheldrick, 2015b) and *OLEX2* (Dolomanov *et al.*, 2009).

supporting information

Acta Cryst. (2024). E80, 1135-1137 [https://doi.org/10.1107/S205698902400971X]

Crystal structure of a hydrogen-bonded 2:1 co-crystal of 4-nitrophenol and 4,4'-bipyridine

Angela Gotingco, Merary Villanueva Contreras, Jeanette A. Wolfarth, S. Chantal E. Stieber and Zoe Y. Marr

Computing details

4,4'-Bipyridine–4-nitrophenol (1/2)

Crystal data

$C_{10}H_8N_2 \cdot 2C_6H_5NO_2$

$M_r = 434.40$

Monoclinic, $P2_1/n$

$a = 12.3711$ (7) Å

$b = 3.8255$ (2) Å

$c = 21.4175$ (12) Å

$\beta = 104.195$ (2)°

$V = 982.65$ (9) Å³

$Z = 2$

$F(000) = 452$

$D_x = 1.468$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5585 reflections

$\theta = 2.2$ – 30.3 °

$\mu = 0.11$ mm⁻¹

$T = 123$ K

Prism, clear colourless

$0.47 \times 0.07 \times 0.03$ mm

Data collection

Bruker D8 Venture

diffractometer

φ and ω scans

Absorption correction: multi-scan
(SADABS; Krause *et al.*, 2015)

$T_{\min} = 0.501$, $T_{\max} = 0.746$

24119 measured reflections

2957 independent reflections

2302 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.076$

$\theta_{\max} = 30.4$ °, $\theta_{\min} = 2.0$ °

$h = -17 \rightarrow 17$

$k = -5 \rightarrow 5$

$l = -30 \rightarrow 30$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.057$

$wR(F^2) = 0.161$

$S = 1.10$

2957 reflections

149 parameters

0 restraints

Primary atom site location: dual

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0648P)^2 + 0.7817P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.47$ e Å⁻³

$\Delta\rho_{\min} = -0.37$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O3	0.36469 (10)	0.2681 (4)	0.20323 (6)	0.0241 (3)
O1	0.76718 (11)	1.0265 (4)	0.13543 (7)	0.0315 (3)
O2	0.66554 (13)	0.9310 (5)	0.03941 (6)	0.0372 (4)
N1	1.07530 (12)	0.6778 (4)	0.16823 (7)	0.0224 (3)
N2	0.68271 (12)	0.9075 (4)	0.09853 (7)	0.0243 (3)
C3	1.01552 (13)	0.5393 (4)	0.03500 (7)	0.0184 (3)
C6	0.60089 (14)	0.7357 (4)	0.12575 (8)	0.0196 (3)
C11	0.49848 (14)	0.6398 (5)	0.08606 (8)	0.0220 (3)
H11	0.482527	0.682639	0.041035	0.026*
C7	0.62601 (14)	0.6748 (4)	0.19178 (8)	0.0209 (3)
H7	0.696365	0.741876	0.218231	0.025*
C1	0.97805 (15)	0.5281 (5)	0.13953 (8)	0.0227 (4)
H1	0.928687	0.468694	0.165566	0.027*
C9	0.44362 (14)	0.4190 (4)	0.17947 (8)	0.0202 (3)
C8	0.54795 (14)	0.5162 (4)	0.21849 (8)	0.0211 (3)
H8	0.564758	0.472398	0.263503	0.025*
C2	0.94480 (14)	0.4545 (5)	0.07449 (8)	0.0223 (3)
H2	0.874671	0.347421	0.056842	0.027*
C10	0.42023 (14)	0.4814 (5)	0.11295 (8)	0.0228 (4)
H10	0.350120	0.414291	0.086214	0.027*
C5	1.14324 (14)	0.7645 (5)	0.13058 (8)	0.0238 (4)
H5	1.212254	0.874593	0.149718	0.029*
C4	1.11678 (14)	0.6999 (5)	0.06463 (8)	0.0230 (4)
H4	1.167421	0.764784	0.039720	0.028*
H3	0.386 (2)	0.254 (8)	0.2445 (15)	0.053 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O3	0.0195 (6)	0.0320 (7)	0.0194 (6)	−0.0049 (5)	0.0022 (5)	0.0017 (5)
O1	0.0236 (6)	0.0378 (8)	0.0325 (7)	−0.0049 (6)	0.0056 (5)	0.0044 (6)
O2	0.0364 (8)	0.0536 (10)	0.0226 (7)	−0.0013 (7)	0.0094 (6)	0.0087 (6)
N1	0.0234 (7)	0.0230 (7)	0.0203 (7)	0.0029 (6)	0.0044 (5)	−0.0015 (5)
N2	0.0222 (7)	0.0275 (8)	0.0238 (7)	0.0040 (6)	0.0066 (5)	0.0048 (6)
C3	0.0183 (7)	0.0169 (7)	0.0198 (7)	0.0014 (6)	0.0044 (6)	−0.0002 (6)
C6	0.0201 (7)	0.0202 (7)	0.0187 (7)	0.0029 (6)	0.0052 (6)	0.0017 (6)
C11	0.0240 (8)	0.0252 (8)	0.0153 (7)	0.0043 (7)	0.0018 (6)	0.0005 (6)
C7	0.0195 (7)	0.0220 (8)	0.0185 (7)	−0.0005 (6)	−0.0004 (6)	0.0004 (6)
C1	0.0234 (8)	0.0245 (8)	0.0213 (8)	0.0000 (6)	0.0077 (6)	−0.0009 (6)

C9	0.0196 (7)	0.0202 (8)	0.0196 (7)	0.0015 (6)	0.0028 (6)	-0.0009 (6)
C8	0.0232 (8)	0.0223 (8)	0.0159 (7)	-0.0005 (6)	0.0009 (6)	0.0003 (6)
C2	0.0200 (8)	0.0254 (8)	0.0212 (8)	-0.0030 (6)	0.0044 (6)	-0.0024 (6)
C10	0.0200 (8)	0.0279 (9)	0.0175 (7)	0.0010 (6)	-0.0010 (6)	-0.0015 (6)
C5	0.0190 (8)	0.0283 (9)	0.0227 (8)	0.0002 (6)	0.0023 (6)	-0.0025 (7)
C4	0.0182 (7)	0.0283 (9)	0.0224 (8)	-0.0015 (6)	0.0048 (6)	-0.0011 (7)

Geometric parameters (Å, °)

O3—C9	1.338 (2)	C11—C10	1.382 (2)
O3—H3	0.86 (3)	C7—H7	0.9500
O1—N2	1.232 (2)	C7—C8	1.378 (2)
O2—N2	1.2346 (19)	C1—H1	0.9500
N1—C1	1.338 (2)	C1—C2	1.381 (2)
N1—C5	1.341 (2)	C9—C8	1.405 (2)
N2—C6	1.444 (2)	C9—C10	1.403 (2)
C3—C3 ⁱ	1.484 (3)	C8—H8	0.9500
C3—C2	1.396 (2)	C2—H2	0.9500
C3—C4	1.400 (2)	C10—H10	0.9500
C6—C11	1.391 (2)	C5—H5	0.9500
C6—C7	1.391 (2)	C5—C4	1.392 (2)
C11—H11	0.9500	C4—H4	0.9500
C9—O3—H3	110.4 (19)	C2—C1—H1	117.9
C1—N1—C5	117.07 (15)	O3—C9—C8	122.52 (15)
O1—N2—O2	122.44 (16)	O3—C9—C10	118.15 (15)
O1—N2—C6	118.51 (14)	C10—C9—C8	119.32 (15)
O2—N2—C6	119.06 (15)	C7—C8—C9	120.32 (15)
C2—C3—C3 ⁱ	121.30 (18)	C7—C8—H8	119.8
C2—C3—C4	116.84 (15)	C9—C8—H8	119.8
C4—C3—C3 ⁱ	121.86 (18)	C3—C2—H2	120.4
C11—C6—N2	119.80 (15)	C1—C2—C3	119.19 (16)
C11—C6—C7	121.33 (15)	C1—C2—H2	120.4
C7—C6—N2	118.86 (15)	C11—C10—C9	120.44 (15)
C6—C11—H11	120.4	C11—C10—H10	119.8
C10—C11—C6	119.16 (15)	C9—C10—H10	119.8
C10—C11—H11	120.4	N1—C5—H5	118.7
C6—C7—H7	120.3	N1—C5—C4	122.70 (16)
C8—C7—C6	119.41 (15)	C4—C5—H5	118.7
C8—C7—H7	120.3	C3—C4—H4	120.0
N1—C1—H1	117.9	C5—C4—C3	119.98 (16)
N1—C1—C2	124.22 (16)	C5—C4—H4	120.0
O3—C9—C8—C7	-179.23 (16)	C3 ⁱ —C3—C4—C5	-179.13 (19)
O3—C9—C10—C11	179.35 (16)	C6—C11—C10—C9	0.2 (3)
O1—N2—C6—C11	171.57 (16)	C6—C7—C8—C9	-0.4 (3)
O1—N2—C6—C7	-7.4 (2)	C11—C6—C7—C8	0.0 (3)
O2—N2—C6—C11	-8.2 (2)	C7—C6—C11—C10	0.1 (3)

O2—N2—C6—C7	172.86 (16)	C1—N1—C5—C4	-1.0 (3)
N1—C1—C2—C3	-0.1 (3)	C8—C9—C10—C11	-0.6 (3)
N1—C5—C4—C3	0.2 (3)	C2—C3—C4—C5	0.6 (3)
N2—C6—C11—C10	-178.82 (16)	C10—C9—C8—C7	0.7 (3)
N2—C6—C7—C8	178.95 (15)	C5—N1—C1—C2	0.9 (3)
C3 ⁱ —C3—C2—C1	179.06 (19)	C4—C3—C2—C1	-0.7 (2)

Symmetry code: (i) $-x+2, -y+1, -z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3 \cdots N1 ⁱⁱ	0.86 (3)	1.84 (3)	2.6921 (19)	174 (3)

Symmetry code: (ii) $-x+3/2, y-1/2, -z+1/2$.