

Received 28 March 2025

Accepted 7 April 2025

Edited by W. T. A. Harrison, University of
Aberdeen, United Kingdom

Keywords: crystal structure; 4-aminoantipyrine;
cadmium(II); tetranuclear; six-coordinate
geometry index; hydrogen bonding.

CCDC reference: 2441688

Supporting information: this article has
supporting information at journals.iucr.org/e

Di- μ_3 -chlorido-1:2:3 κ^3 Cl;2:3:4 κ^3 Cl-di- μ_2 -chlorido-1:2 κ^2 Cl;3:4 κ^2 Cl-tetrakis[(4-amino-1,5-dimethyl-2-phenyl-2,3-dihydro-1H-pyrazol-3-one- κ^2 N⁴,O)-chloridocadmium(II)] 1.7-hydrate: a new six-coordinate geometry index, τ_6

Helen Stoeckli-Evans,^{a*} M. G. Shankar,^b R. Kumaravel,^c A. Subashini,^b T. Sabari Girisun,^d K. Ramamurthi,^e Monika Kučeráková,^f Michal Dušek^f and Aurélien Crochet^g


^aInstitute of Physics, University of Neuchâtel, Rue Emile-Argand 11, CH-2000 Neuchâtel, Switzerland, ^bPG and Research Department of Physics, Srimad Andavan Arts and Science College (Autonomous), Affiliated to Bharathidasan University, Tiruchirappalli 620005, Tamilnadu, India, ^cDepartment of Physics, Annappoorana Engineering College (Autonomous), Salem 636308, Tamilnadu, India, ^dNanophotonics Laboratory, Department of Physics, Bharathidasan University, Tiruchirappalli 620024, Tamilnadu, India, ^eCrystal Growth and Thin Film Laboratory, Department of Physics, Bharathidasan University, Tiruchirappalli 620024, Tamilnadu, India, ^fInstitute of Physics ASCR, Na Slovance 2, 182 21 Praha 8, Czech Republic, and ^gChemistry Department, University of Fribourg, Chemin du Musée 9, CH-1700 Fribourg, Switzerland.

*Correspondence e-mail: helen.stoeckli-evans@unine.ch


The title centrosymmetric tetranuclear cadmium(II) complex of 4-aminoantipyrine and chloride ions, $[\text{Cd}_4\text{Cl}_8(\text{C}_{11}\text{H}_{13}\text{N}_3\text{O})_4]\cdot 1.7\text{H}_2\text{O}$, was synthesized using methanol as solvent. The two independent Cd^{2+} ions in the asymmetric unit have different geometries; the outer Cd atoms have fivefold CdONCl_3 coordination spheres, while the inner Cd atoms have sixfold CdONCl_4 coordination spheres. The complex is consolidated by intramolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds. In the crystal, a combination of $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{O}_w-\text{H}\cdots\text{Cl}$ (w = water) hydrogen bonds link the components to form chains propagating along the a -axis direction. The chains are crosslinked by $\text{C}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds to form a three-dimensional structure. A new geometry index, τ_6 , is proposed to quantitatively estimate the geometry of a sixfold coordinated atom.

1. Chemical context

Antipyrine derivatives have gained interest as model compounds for functional materials due to their various properties, including antioxidant (Bashkatova *et al.*, 2005), antiputrefactive (Abd El Rehim *et al.*, 2001) and optical properties (Collado *et al.*, 2000; Coolen *et al.*, 1999). One such analogue is 4-amino-1,5-dimethyl-2-phenylpyrazol-3-one, $\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}$, known as 4-aminoantipyrine (4-AAP). Its crystal structure was first reported by Li *et al.* (2013), who also analysed its electronic structure and that of a number of its derivatives, including 4-(dimethylamino)antipyrine. A low-temperature structural analysis of 4-AAP has been reported by Mnguni & Lemmerer (2015). The structure features a five-membered lactam ring in the pyrazole unit and a free amino group. Pyrazolone-based ligands exhibit variable complexing behaviour and a variety of coordination possibilities to metal centres. Such complexes have applications in both chemistry and the pharmaceutical sciences (Raman *et al.*, 2014). Derivatives of 4-AAP have also emerged as important compounds in the fields of biology and medicine (Senthilkumar *et al.*, 2016). The presence of heteroatoms influences the electron



distribution, which in turn enhances its reactivity and chelating properties (Matczak & Domagała, 2017; Joule & Mills, 2008). Due to this excellent chelating effect it can form a wide variety of metal complexes with almost all transition-metal ions and lanthanides. They have applications in many fields of research, such as sensor development (Banasz & Wałęsa-Chorab, 2019), renewable energy materials (Zhang *et al.*, 2018), chemosensors (Selvan *et al.*, 2016), DNA binding (Sakthivel *et al.*, 2020), antipyretic (Turan-Zitouni *et al.*, 2001), antioxidant (Bashkatova *et al.*, 2005), anticancer (Bose *et al.*, 2005) and anti-inflammatory agents (Sondhi *et al.*, 1999).



Cadmium(II) complexes in general are notable because of their excellent optical and electronic properties (Venkataramanan *et al.*, 1997). They also have uses in environmental and analytical chemistry, and materials science (Adhikari *et al.*, 2020; Roccanova *et al.*, 2017; Cheng *et al.*, 2017). A search of the Cambridge Structural Database (CSD, Version 5.46, update February 2025; Groom *et al.*, 2016) for transition-metal complexes of 4-AAP gave 13 hits. Two of these are cadmium(II) complexes, *viz.* *catena*-[[4-aminoantipyrine]aqua-(μ_2 -5-nitroisophthalato)cadmium(II)] monohydrate] (CSD refcode CIXQIK; Wang *et al.*, 2008) and *catena*-[bis(μ -4-amino-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one)-tetradeca- μ -chloro-bis(ethanol)heptacadmium(II)] (IQATAY; Hu *et al.*, 2012). In both of these, 4-aminoantipyrine acts as a bidentate ligand, donating the lone pairs of electrons from the amino N atom and the carbonyl O atom to the cadmium ion.

Depending on the stoichiometry and reaction conditions, the structure of the complex can vary. For example, when reacting 4-aminoantipyrine (0.0203 g, 0.1 mmol) with $\text{CdCl}_2 \cdot 2.5\text{H}_2\text{O}$ (0.142 g, 0.06 mmol) in a mixture of ethanol and ethyl acetate (1:3 v/v) at 343 K, Hu *et al.* (2012) synthesized a mono-periodic coordination polymer, *catena*-[bis(μ -4-amino-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one)-tetradeca- μ -chloro-bis(ethanol)heptacadmium(II)]; structure

IQATAY mentioned above. The complex unit possesses inversion symmetry and the asymmetric unit consists of 3.5 Cd^{II} atoms coordinated to seven bridging chloride ions, one bridging 4-aminoantipyrine ligand and one ethanol molecule. In the present work, a new centrosymmetric tetranuclear cadmium(II) complex of 4-aminoantipyrine was synthesized when reacting an equimolar ratio of 4-AAP and $\text{CdCl}_2 \cdot 2.5\text{H}_2\text{O}$ using methanol as the solvent. Herein, we report on the structure and various properties of the title complex, $[\text{Cd}_4\text{Cl}_8(\text{C}_{11}\text{H}_{13}\text{N}_3\text{O})_4] \cdot 1.7\text{H}_2\text{O}$ (**I**) (Scheme 1), and compare them to those of the heptacadmium complex IQATAY.

2. τ_6 , a sixfold geometry index

The molecular structure of (**I**) was found to be centrosymmetric with two independent cadmium(II) atoms (see *Structural commentary* section). The two outer Cd^{II} atoms have fivefold coordination spheres, while the inner Cd^{II} atoms have sixfold coordination spheres. Previously, a number of authors have described methods to measure the size of distortions in polyhedra; some have included only distortions in the bond lengths from their average values (Robinson *et al.*, 1971; Muetterties & Guggenberger, 1974; Brown, 2006). Robinson *et al.* (1971) introduced the notion of quadratic elongation (QE) and this analysis is incorporated in *PLATON* (Spek, 2020). As of yet, no simple geometry index has been defined to describe the geometry of an octahedral coordination sphere. Deviations of the bond angles from their ideal values were largely ignored, but as shown below, when considered, they provide a simple method to calculate the various geometry indexes.

The fivefold geometry index τ_5 was proposed by Addison *et al.* (1984) and is illustrated in Scheme 2 ($\tau_5 = 0$ for a perfect square pyramid and $\tau_5 = 1$ for a trigonal bipyramid). A fourfold geometry index τ_4 was proposed by Yang *et al.* (2007) and is also illustrated in Scheme 2 ($\tau_4 = 0$ for a perfect square-planar geometry and $\tau_4 = 1$ for a pyramidal geometry). Following the reasoning of Yang *et al.* (2007), we propose a simple numerical estimation of the geometry of a sixfold coordination sphere; see Scheme 3 where $\tau_6 = [(3 \times 180^\circ) - (\alpha_1 + \alpha_2 + \alpha_3)]/180^\circ = 0$ for a perfect octahedron where $\alpha_1 = \alpha_2 = \alpha_3 = 180^\circ$. A value of 0.75 is obtained for a trigonal prismatic geometry, where $\alpha_1 = \alpha_2 = \alpha_3 = 135^\circ$ and a value of 1.00 is obtained for a pentagonal pyramidal geometry, where $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha_5 = 72^\circ$.


3. Structural commentary

The molecular structure of (**I**) is illustrated in Fig. 1. This tetranuclear complex possesses inversion symmetry and crystallizes as a 1.7 hydrate. Selected bond lengths and angles in the complex are given in Table 1. The two independent Cd^{II} atoms of the asymmetric unit have different geometries, as shown in Figs. 1 and 2.


The outer Cd atoms [Cd1 and Cd1ⁱ; symmetry code: (i) $-x + 2, -y + 2, -z + 1$] have fivefold CdONCl_3 coordination

spheres with a distorted shape, and according to the definition of Addison *et al.* (1984), the structural index $\tau_5 = (161.0^\circ - 142.63^\circ)/60^\circ = 0.31$. The inner Cd atoms (Cd2 and Cd2ⁱ) have sixfold CdONCl₄ coordination spheres. The three principal (*trans*) bond angles O2—Cd2—Cl2, N6—Cd2—Cl3ⁱ and Cl3—Cd2—Cl4 are 159.61 (4), 161.49 (4) and 170.34 (2)°, respectively, compared to 180° for a perfect octahedron. The τ_6 geometry index gives a value of $[540^\circ - (159.61^\circ + 161.49^\circ + 170.34^\circ)]/180^\circ = 0.27$, a significant distortion from the geometry of a perfect octahedron. The QE parameter for atom Cd2 is 1.027 (PLATON; Spek, 2020).


4-fold coordination



square planar $\tau_4 = [360^\circ - (\alpha 1 + \alpha 2)]/141^\circ = 0$




trigonal pyramidal $\tau_4 = [360^\circ - (\alpha 1 + \alpha 2)]/141^\circ = 0.85$




tetrahedral $\tau_4 = [360^\circ - (\alpha 1 + \alpha 2)]/141^\circ = 1.0$

5-fold coordination



$\tau_5 = (\alpha 1 - \alpha 2)/60^\circ = 0$

square pyramidal



$\tau_5 = (\alpha 1 - \alpha 2)/60^\circ = 1$

trigonal bipyramidal

Scheme 2

Comparing the coordination geometry of the four Cd atoms in IQATAY, it was found that the values of τ_6 for atoms Cd1, Cd2, Cd3 and Cd4 are 0, 0.06, 0.16 and 0.17, respectively. Hence, in this case, atom Cd1, which lies on an inversion centre, has a perfect octahedral geometry. Atom Cd2 has a very small distortion from a perfect octahedron. Atoms Cd3 and Cd4 have distorted geometries but less so than that observed for atom Cd2 in complex (I). The QE parameters for

Table 1


Selected geometric parameters (Å, °).

Cd1—O1	2.3498 (14)	Cd2—Cl2	2.5929 (5)
Cd1—N3	2.2863 (17)	Cd2—Cl3 ⁱ	2.6734 (5)
Cd1—Cl1	2.4486 (5)	Cd2—Cl3	2.7631 (5)
Cd1—Cl2	2.5396 (5)	Cd2—Cl4	2.5157 (5)
Cd1—Cl3	2.6908 (5)	Cd1...Cd2	3.7831 (5)
Cd2—O2	2.3107 (13)	Cd2...Cd2 ⁱ	4.0745 (4)
Cd2—N6	2.4346 (17)	Cd1...Cd2 ⁱ	4.6003 (4)
N3—Cd1—Cl1	142.63 (5)	Cd1—Cl2—Cd2	94.958 (15)
O1—Cd1—Cl3	161.00 (4)	Cd2 ⁱ —Cl3—Cd1	118.092 (16)
O2—Cd2—Cl2	159.61 (4)	Cd2 ⁱ —Cl3—Cd2	97.069 (14)
N6—Cd2—Cl3 ⁱ	161.49 (4)	Cd1—Cl3—Cd2	87.825 (14)
Cl4—Cd2—Cl3	170.343 (16)		


Symmetry code: (i) $-x + 2, -y + 2, -z + 1$.

atoms Cd1 to Cd4 in IQATAY are 1.002, 1.005, 1.013 and 1.018, respectively.


6-fold coordination



octahedral $\tau_6 = [540^\circ - (\alpha 1 + \alpha 2 + \alpha 3)]/180^\circ = 0$



trigonal prismatic $\tau_6 = [540^\circ - (\alpha 1 + \alpha 2 + \alpha 3)]/180^\circ = 0.75$



pentagonal pyramidal $\tau_6 = [540^\circ - (5 \times 72^\circ)]/180^\circ = 1$

Scheme 3

In (I), there are two terminal chlorido ligands (Cl1 and Cl4), one μ_2 -bridging Cl atom (Cl2) and one μ_3 -bridging Cl atom (Cl3); see Table 1. In IQATAY, there are three μ_2 -bridging Cl atoms and four μ_3 -bridging Cl atoms. The Cd—Cl bond lengths vary from 2.4486 (5) to 2.7631 (5) Å in (I), compared to a range of 2.551 (2)–2.779 (2) Å in IQATAY. The various Cd...Cd separations and Cd—Cl—Cd bridging angles in (I) are given in Table 1. These values are very similar to those observed for IQATAY.

The two independent 4-AAP molecules in (I) have slightly different conformations. A view of the molecular overlap of the two ligands [ligand 2 (involving atom O2) inverted over




Figure 1

A view of the molecular structure of (**I**), with displacement ellipsoids drawn at the 50% probability level. [Symmetry code: (i) $-x + 2, -y + 2, -z + 1$.]

ligand 1 (involving atom O1)] is shown in Fig. S1 of the supporting information. In (**I**), the ligands are bidentate, coordinating to one Cd atom each time, while in IQATAY, the ligand bridges two Cd atoms. In (**I**), arene ring C4–C9 is inclined to the pyrazole ring mean plane (N1/N2/C1–C3; r.m.s. deviation = 0.05 Å) by 56.7 (1)°, while arene ring C15–C20 is inclined to the pyrazole ring mean plane (N4/N5/C12–C14; r.m.s. deviation 0.044 Å) by 45.2 (1)°. In IQATAY, the corresponding dihedral angle is larger at 62.1 (4)°. In (**I**), the pyrazole ring of ligand 1 (N1/N2/C1–C3) is almost parallel to arene ring C15ⁱ–C20ⁱ of ligand 2, with a centroid–centroid separation of 3.749 (1) Å. The 4-AAP ligands in (**I**) coordinate to the Cd atoms *via* the carbonyl O atom and the 4-amino N atom. The Cd1–O1 bond length is 2.3498 (13) Å compared to 2.3107 (13) Å for Cd2–O2. In IQATAY, the corresponding Cd–O bond length is significantly shorter at 2.245 (6) Å. The Cd–N bond lengths in (**I**) are Cd1–N3 = 2.2863 (17) Å and Cd2–N6 = 2.4346 (17) Å, compared to 2.352 (7) Å for the corresponding bond in IQATAY.

Complex (**I**) is consolidated by intramolecular N–H···Cl and N–H···O hydrogen bonds, which are listed in Table 2 and illustrated in Fig. 1. The N6–H6NA···Cl1 hydrogen bond involves the 4-amino group of ligand 2, while the




Figure 2

A partial view of (**I**), showing the different polyhedra for atoms Cd1 and Cd2 (Mercury; Macrae *et al.*, 2020).

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N3–H3NA···O2 ⁱ	0.88 (3)	1.89 (3)	2.747 (2)	166 (3)
N6–H6NA···Cl1	0.87 (3)	2.54 (3)	3.4073 (19)	172 (2)
N6–H6NB···Cl1 ⁱⁱ	0.86 (3)	2.73 (3)	3.3718 (18)	133 (2)
O1W–H1WA···Cl4 ⁱⁱ	0.85 (4)	2.45 (4)	3.292 (2)	176 (4)
O1W–H1WB···Cl1	0.89 (4)	2.34 (4)	3.220 (2)	168 (3)
C11–H11A···Cl2 ⁱⁱⁱ	0.98	2.82	3.732 (2)	156
C16–H16···Cl2 ⁱ	0.95	2.82	3.589 (2)	138
C21–H21C···Cl4 ^{iv}	0.98	2.69	3.647 (2)	165
C22–H22C···O1 ⁱⁱ	0.98	2.51	3.417 (3)	154

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

N3–H3HA···O2ⁱ hydrogen bond involves the 4-amino group of ligand 1 and the carbonyl O atom of ligand 2.

4. Supramolecular features

In the crystal of (**I**), the molecules are linked by O–H···Cl hydrogen bonds (Table 2), involving the partially occupied water molecules (Cl···H–O–H···Cl). Together with N–H···Cl hydrogen bonds (Table 2), chains are formed propagating along the *a*-axis direction (Fig. 3). The chains enclose two ring motifs. The first, $R_4^2(8)$, involves the amine H atoms and chloride ion Cl1. The second, $R_3^2(8)$, involves an amine H atom, two chloride ions, the water H atoms and atom Cd1, as shown in Fig. 3. The chains are linked by a series of C–H···Cl and C–H···O hydrogen bonds to form a three-dimensional structure (Fig. 4 and Table 2). Atom N3HB does not partake in hydrogen bonding. This phenomenon is not unusual and has been observed previously (CSD; Groom *et al.*, 2016).

5. Hirshfeld surface analysis and fingerprint plots

The Hirshfeld surface (HS) analysis and the associated two-dimensional fingerprint plots were generated with *Crystal-Explorer17* (Spackman *et al.*, 2021) and interpreted following the protocol of Tan *et al.* (2019). The Hirshfeld surface (HS) of (**I**) is illustrated in Fig. 5. It is colour-mapped with the normalized contact distance d_{norm} in the colour range from




Figure 3

A partial view of the crystal packing of (**I**). Only the H atoms involved in the N–H···Cl and O–H···Cl hydrogen bonds (Table 1) have been included.




Figure 4

A view along the *a*-axis direction of the crystal packing of (I). Only the H atoms (grey spheres) involved in the various hydrogen bonds (Table 1) have been included.

0.00 to 1.41 a.u. There are a significant number of large red spots, indicating that in these regions the interatomic distances in the crystal are shorter than the sum of the van der Waals radii.

The full two-dimensional fingerprint plot for (I), and those delineated into H...H, Cl...H/H...Cl, C...H/H...C, O...H/H...O and N...H/H...N contacts, are given in Fig. 6. The H...H contacts have a major contribution to the HS of 43.1%. The second most significant contributions are from the Cl...H/H...Cl contacts at 26.1%, with sharp peaks at $d_i + d_e \simeq 2.2$ Å. This reflects the presence of the seven H...Cl interactions in the crystal structure (see Table 2). The C...H/H...C interactions contribute 15.0%, with relatively sharp spikes at $d_i + d_e \simeq 2.5$ Å. The O...H/H...O interactions contribute 10%, with sharp peaks at $d_i + d_e \simeq 2.35$ Å. The sharp pincer-




Figure 5

The Hirshfeld surface of (I) mapped over d_{norm}

like peaks for these three interatomic interactions indicate that they are significant. Finally, the N...H/H...N contacts contribute 2.5%, while other interatomic contacts contribute less than 1% to the overall HS.

6. Database survey

A search of the Cambridge Structural Database (CSD, Version 5.46, update February 2025; Groom *et al.*, 2016) for sixfold-coordinated cadmium(II) metal complexes was carried out with the following restrictions: three-dimensional structure, $R \leq 0.05$, no disorder, no ions, no polymers, single crystals only and no bidentate ligands with fewer than four atoms, such as nitrate or acetate. Over 720 hits were obtained.

Although the majority of the complexes have octahedral coordination spheres, there are a number of complexes with a




Figure 6

The full two-dimensional fingerprint plot for (I), and those delineated into H...H, Cl...H/H...Cl, C...H/H...C, O...H/H...O and N...H/H...N contacts.

trigonal prismatic geometry. One such complex is (1,5,5,9-13,13,20,20-octamethyl-3,7,11,15,18,22-hexaazabicyclo[7.7.7]-tricosane)cadmium(II) dinitrate dihydrate (CSD refcode ULESAH; Alcock *et al.*, 2016), illustrated in Figs. 7(a) and 7(b). Here the structural index $\tau_6 = 0.73$ and the QE parameter for atom Cd1 is 1.24.

Examples of pentagonal pyramidal geometry are rare. One example is (benzimidazole)(4,5,9,24-tetraethyl-10,23-dimethyl-13,20,25,26,27-pentaazapentacyclo[20.2.1.1^{3,6}.1^{8,11}.0^{14,19}]heptacosane-1,3,5,7,9,11(27),12,14,16,18,20,22(25),23-tridecaene)cadmium(II) nitrate chloroform solvate (VAPSAG; Sessler *et al.*, 1989), illustrated in Figs. 7(c) and 7(d); here the structural index $\tau_5 = 1.02$ (the QE parameter for atom Cd1 is 1.33). Another example is that of complex tetraaquabis(μ_2 -xanthurenato)dicadmium(II), with two Cd atoms related by a centre of symmetry (CIJBII; Tratar *et al.*, 2012). Here the structural index $\tau_6 = 1.0$ (QE parameter for atom Cd1 = 1.31).

7. Thermal analysis

To investigate the thermal characteristics of (**I**), a SQT-Q600 V20.9 Build 20 Universal Thermo Analytical system was used under a nitrogen atmosphere and with a sample weight of 2.044 mg in an alumina crucible over a temperature range from 25 to 800 °C, with a heating rate of 20 °C per minute. The TGA and DTA results are illustrated in Fig. S2 of the supporting information.

The melting point of the complex, observed from the DTA curve, is 245 °C (518 K). According to the TGA curve, the first weight loss step (2.3%) corresponds to the loss of water molecules present in the title complex. The second step

(34.7% loss) is probably due to the gradual decomposition of one 4-aminoantipyrine molecule, accompanied by the release of chlorine gas. The third step (48.9% loss) may correspond to the loss of one 4-aminoantipyrine molecule and one CdCl_2 molecule. The final step may be due to the degradation of the rest of the complex, leading to the formation of the final residue of CdO (observed 16.23%; calculated 16.63%)

8. Synthesis and spectroscopic data

An equimolar mixture of 4-aminoantipyrine (0.100 mmol) and cadmium(II) chloride (0.100 mmol) was dissolved in 20 ml methanol and refluxed at 363 K using an oil bath. After 6 h the solution was filtered and left aside at room temperature. Orange needle-like crystals of (**I**) were obtained by evaporation of the solvent over a period of one week. The melting point is 518 K (245 °C), as seen from the DTA curve (Fig. S2).

The FT-IR spectrum of (**I**) was recorded with a JASCO Infrared spectrometer (400–4000 cm^{-1}) using the KBr pellet technique (Fig. S3 of the supporting information). The FT-IR spectrum of the free ligand (Swaminathan *et al.*, 2009) was compared with that of the cadmium complex to verify the coordination of 4-AAP with the Cd^{2+} ions. For 4-AAP, prominent peaks are seen at 3432 and 3325 cm^{-1} , corresponding to the asymmetric and symmetric stretching modes of NH_2 , respectively, and at 1679 cm^{-1} for the carbonyl-group stretching band. In the FT-IR spectrum of (**I**), sharp peaks were observed around 3301, 3213 and 1631 cm^{-1} . It may be seen that the vibrational frequencies of NH_2 and $\text{C}=\text{O}$ are shifted to lower frequencies upon Cd^{2+} coordination, consistent with the crystal structure.




Figure 7

(a)/(b) The trigonal prismatic coordination sphere of ULESAH (redrawn from Alcock *et al.*, 2016) and (c)/(d) the pentagonal pyramidal coordination sphere of VAPSAG (redrawn from Sessler *et al.*, 1989).

9. Refinement details

Crystal data, data collection and structure refinement details are summarized in Table 3. The NH₂ and water H atoms were located in difference Fourier maps and freely refined. C-bound H atoms were included in calculated positions and treated as riding, with C–H = 0.95–0.98 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Acknowledgements

MK and MD acknowledge using the CzechNanoLab Research Infrastructure supported by MEYS CR (No. LM2018110) for crystallographic analysis. HSE is grateful to the University of Neuchâtel for their support over the years.

References

- Abd El Rehim, S. S., Ibrahim, M. A. M. & Khalid, K. F. (2001). *Mater. Chem. Phys.* **70**, 268–273.
- Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
- Adhikari, S., Bhattacharjee, T., Das, A., Roy, S., Daniliuc, C. G., Zaręba, J. K., Bauzá, A. & Frontera, A. (2020). *CrystEngComm*, **22**, 8023–8035.
- Alcock, L. J., Cavigliasso, G., Willis, A. C., Stranger, R. & Ralph, S. F. (2016). *Dalton Trans.* **45**, 9036–9040.
- Banasz, R. & Wałęsa-Chorab, M. (2019). *Coord. Chem. Rev.* **389**, 1–18.
- Bashkatova, N. V., Korotkova, E. I., Karbainov, Y. A., Yagovkin, A. Y. & Bakibaev, A. A. (2005). *J. Pharm. Biomed. Anal.* **37**, 1143–1147.
- Bose, R., Murty, D. S. R. & Chakrapani, G. (2005). *J. Radioanal. Nucl. Chem.* **265**, 115–122.
- Brown, I. D. (2006). *Acta Cryst.* **B62**, 692–694.
- Cheng, C. C., Shen, N. N., Du, C. F., Wang, Z., Li, J. R. & Huang, X. Y. (2017). *Inorg. Chem. Commun.* **85**, 21–25.
- Collado, M. S., Mantovani, V. E., Goicoechea, H. C. & Olivieri, H. C. (2000). *Talanta*, **52**, 909–920.
- Coolen, S. A., Ligor, T., Lieshout, M. & Huf, F. A. (1999). *J. Chromatogr. B Biomed. Sci. Appl.* **732**, 103–113.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hu, J., Qi, J., Luo, Y., Yin, T., Wang, J., Wang, C., Li, W. & Liang, L. (2012). *Arab. J. Chem.* **14**, 103117–103124.
- Joule, J. A. & Mills, K. (2008). In *Heterocyclic Chemistry*, 5th ed. Chichester: John Wiley & Sons.
- Li, Y., Liu, Y., Wang, H., Xiong, X., Wei, P. & Li, F. (2013). *Molecules*, **18**, 877–893.
- Macrae, C. F., Sovago, I., Cottrell, S. J., Galek, P. T. A., McCabe, P., Pidcock, E., Platings, M., Shields, G. P., Stevens, J. S., Towler, M. & Wood, P. A. (2020). *J. Appl. Cryst.* **53**, 226–235.
- Matczak, M. & Domagała, M. (2017). *J. Mol. Model.* **23**, 268–279.
- Mguni, M. J. & Lemmerer, A. (2015). *Acta Cryst.* **C71**, 103–109.
- Muettterties, E. L. & Guggenberger, L. J. (1974). *J. Am. Chem. Soc.* **96**, 1748–1756.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Petríček, V., Palatinus, L., Plášil, J. & Dušek, M. (2023). *Z. Kristallogr.* **238**, 271–282.
- Raman, N., Sakthivel, A. & Pravin, N. (2014). *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **125**, 404–413.
- Rigaku OD (2022). *CrysAlis PRO*. Rigaku Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Robinson, K., Gibbs, G. V. & Ribbe, P. H. (1971). *Science*, **172**, 567–570.
- Rocanova, R., Ming, W., Whiteside, V. R., McGuire, M. A., Sellers, I. R., Du, M. H. & Saparov, B. (2017). *Inorg. Chem.* **56**, 13878–13888.
- Sakthivel, A., Jeyasubramanian, M., Thangagiri, B. & Raja, J. D. (2020). *J. Mol. Struct.* **1222**, 128885.
- Selvan, G. T., Kumaresan, M., Sivaraj, R., Enoch, I. V. M. V. & Selvakumar, P. M. (2016). *Sens. Actuators B Chem.* **229**, 181–189.
- Senthilkumar, K., Thirumoorthy, K., Dragonetti, C., Marinotto, D., Righetto, S., Colombo, A., Haukka, M. & Palanisami, N. (2016). *Dalton Trans.* **45**, 11939–11943.
- Sessler, J. L., Murai, T. & Lynch, V. (1989). *Inorg. Chem.* **28**, 1333–1341.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Sondhi, S. M., Sharma, V. K., Verma, R. P., Singhal, N., Shukla, R., Raghubir, R. & Dubey, M. P. (1999). *Synthesis*, **1999**, 878–884.
- Spackman, P. R., Turner, M. J., McKinnon, J. J., Wolff, S. K., Grimwood, D. J., Jayatilaka, D. & Spackman, M. A. (2021). *J. Appl. Cryst.* **54**, 1006–1011.
- Spek, A. L. (2020). *Acta Cryst.* **E76**, 1–11.
- Swaminathan, J., Ramalingam, M., Sethuraman, V., Sundaraganesan, N. & Sebastian, S. (2009). *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **73**, 593–600.
- Tan, S. L., Jotani, M. M. & Tiekink, E. R. T. (2019). *Acta Cryst.* **E75**, 308–318.
- Tratar Pirc, E., Cer-Kerčmar, K., Modec, B. & Bukovec, P. (2012). *Monatsh. Chem.* **143**, 1643–1648.
- Turan-Zitouni, G., Sivaci, M., Kılıç, F. S. & Erol, K. (2001). *Eur. J. Med. Chem.* **36**, 685–689.
- Venkataramanan, V., Bhat, H. L., Srinivasan, M. R., Ayyub, P. & Multani, M. S. (1997). *J. Raman Spectrosc.* **28**, 779–784.

Table 3

Experimental details.

Crystal data	
Chemical formula	[Cd ₄ Cl ₈ (C ₁₁ H ₁₃ N ₃ O) ₄].1.7H ₂ O
M_r	1576.80
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	95
a, b, c (Å)	11.0702 (2), 13.5058 (2), 19.0700 (3)
β (°)	102.006 (2)
V (Å ³)	2788.82 (8)
Z	2
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	1.94
Crystal size (mm)	0.23 × 0.09 × 0.06
Data collection	
Diffractometer	Rigaku OD SuperNova Dual source diffractometer with an AtlasS2 detector
Absorption correction	Multi-scan (<i>CrysAlis PRO</i> ; Rigaku OD, 2022)
$T_{\text{min}}, T_{\text{max}}$	0.659, 1.000
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	46639, 7273, 6505
R_{int}	0.033
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.696
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.022, 0.051, 1.05
No. of reflections	7273
No. of parameters	362
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.60, -0.50

Computer programs: *CrysAlis PRO* (Rigaku OD, 2022), *SUPERFLIP* (Palatinus & Chapuis, 2007), *JANA2006* (Petríček *et al.*, 2023), *Mercury* (Macrae *et al.*, 2020), *SHELXL2019* (Sheldrick, 2015), *PLATON* (Spek, 2020) and *publCIF* (Westrip, 2010).

- Wang, Q., Wu, M.-J., Yang, E.-C., Wang, X.-G. & Zhao, X.-J. (2008). *J. Coord. Chem.* **61**, 595–604.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yang, L., Powell, D. R. & Houser, R. P. (2007). *Dalton Trans.* pp. 955–964.
- Zhang, J., Xu, L. & Wong, W. (2018). *Coord. Chem. Rev.* **355**, 180–198.

supporting information

Acta Cryst. (2025). E81, 393–400 [https://doi.org/10.1107/S2056989025003123]

Di- μ_3 -chlorido-1:2:3 κ^3 Cl;2:3:4 κ^3 Cl-di- μ_2 -chlorido-1:2 κ^2 Cl;3:4 κ^2 Cl-tetrakis[(4-amino-1,5-dimethyl-2-phenyl-2,3-dihydro-1*H*-pyrazol-3-one- κ^2 N⁴,O)chloridocadmium(II)] 1.7-hydrate: a new six-coordinate geometry index, τ_6

Helen Stoeckli-Evans, M. G. Shankar, R. Kumaravel, A. Subashini, T. Sabari Girisun, K. Ramamurthi, Monika Kučeráková, Michal Dušek and Aurélien Crochet

Computing details

Di- μ_3 -chlorido-1:2:3 κ^3 Cl;2:3:4 κ^3 Cl-di- μ_2 -chlorido-1:2 κ^2 Cl;3:4 κ^2 Cl-tetrakis[(4-amino-1,5-dimethyl-2-phenyl-2,3-dihydro-1*H*-pyrazol-3-one- κ^2 O,N⁴)chloridocadmium(II)] 1.7-hydrate

Crystal data

[Cd₄Cl₈(C₁₁H₁₃N₃O)₄]·1.7H₂O

$M_r = 1576.80$

Monoclinic, $P2_1/n$

$a = 11.0702$ (2) Å

$b = 13.5058$ (2) Å

$c = 19.0700$ (3) Å

$\beta = 102.006$ (2)°

$V = 2788.82$ (8) Å³

$Z = 2$

$F(000) = 1554$

$D_x = 1.878$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 23338 reflections

$\theta = 2.4$ – 29.4°

$\mu = 1.94$ mm⁻¹

$T = 95$ K

Needle, orange

$0.23 \times 0.09 \times 0.06$ mm

Data collection

Rigaku OD SuperNova Dual source
diffractometer with an AtlasS2 detector

Radiation source: micro-focus sealed X-ray tube

Mirror monochromator

Detector resolution: 5.2027 pixels mm⁻¹

ω scans

Absorption correction: multi-scan
(CrysAlis PRO; Rigaku OD, 2022)

$T_{\min} = 0.659$, $T_{\max} = 1.000$

46639 measured reflections

7273 independent reflections

6505 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 29.7^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -13 \rightarrow 15$

$k = -17 \rightarrow 17$

$l = -24 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.022$

$wR(F^2) = 0.051$

$S = 1.05$

7273 reflections

362 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0206P)^2 + 2.4714P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.004$

$$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.50 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cd1	0.69699 (2)	0.94027 (2)	0.38615 (2)	0.01331 (4)	
Cd2	0.87944 (2)	1.07542 (2)	0.54768 (2)	0.01079 (4)	
Cl1	0.55065 (4)	0.86942 (3)	0.45253 (2)	0.01487 (9)	
Cl2	0.73018 (4)	1.11865 (3)	0.42784 (2)	0.01186 (9)	
Cl3	0.90013 (4)	0.89352 (3)	0.48490 (2)	0.01128 (9)	
Cl4	0.89700 (4)	1.23560 (3)	0.61639 (2)	0.01465 (9)	
O1	0.55409 (13)	0.94942 (10)	0.27639 (7)	0.0146 (3)	
O2	0.98088 (12)	0.98270 (10)	0.64391 (7)	0.0141 (3)	
N1	0.56201 (15)	0.92585 (12)	0.15576 (9)	0.0133 (3)	
N2	0.65230 (15)	0.89099 (13)	0.12006 (9)	0.0146 (3)	
N3	0.81571 (16)	0.90156 (13)	0.30520 (9)	0.0136 (3)	
H3NA	0.877 (3)	0.944 (2)	0.3144 (15)	0.034 (8)*	
H3NB	0.849 (2)	0.843 (2)	0.3157 (13)	0.024 (7)*	
N4	0.93489 (15)	0.86164 (12)	0.72079 (9)	0.0128 (3)	
N5	0.82718 (15)	0.82851 (12)	0.74095 (9)	0.0127 (3)	
N6	0.71297 (16)	0.99943 (13)	0.59441 (9)	0.0131 (3)	
H6NA	0.665 (3)	0.967 (2)	0.5601 (15)	0.029 (7)*	
H6NB	0.668 (3)	1.044 (2)	0.6087 (14)	0.026 (7)*	
C1	0.61290 (18)	0.93024 (13)	0.22807 (10)	0.0119 (4)	
C2	0.76124 (18)	0.88714 (14)	0.16873 (10)	0.0135 (4)	
C3	0.73903 (18)	0.90621 (14)	0.23532 (10)	0.0127 (4)	
C4	0.43370 (18)	0.91110 (14)	0.12652 (10)	0.0123 (4)	
C5	0.39217 (19)	0.82220 (14)	0.09338 (10)	0.0143 (4)	
H5	0.449080	0.771716	0.087636	0.017*	
C6	0.26638 (19)	0.80834 (15)	0.06883 (10)	0.0166 (4)	
H6	0.236818	0.748111	0.045759	0.020*	
C7	0.1838 (2)	0.88162 (16)	0.07774 (11)	0.0197 (4)	
H7	0.097618	0.871308	0.061310	0.024*	
C8	0.2263 (2)	0.97020 (16)	0.11061 (11)	0.0202 (4)	
H8	0.169172	1.020341	0.116638	0.024*	
C9	0.35225 (19)	0.98602 (15)	0.13480 (10)	0.0159 (4)	
H9	0.381885	1.047031	0.156594	0.019*	
C10	0.6400 (2)	0.91614 (17)	0.04415 (11)	0.0202 (4)	
H10A	0.561577	0.890208	0.016781	0.030*	
H10B	0.708564	0.886756	0.026115	0.030*	
H10C	0.641657	0.988261	0.038802	0.030*	

C11	0.87995 (19)	0.86561 (16)	0.14715 (12)	0.0186 (4)	
H11A	0.872718	0.803415	0.120124	0.028*	
H11B	0.945901	0.859609	0.190050	0.028*	
H11C	0.899560	0.919594	0.117035	0.028*	
C12	0.90388 (18)	0.93145 (13)	0.66823 (10)	0.0115 (4)	
C13	0.72966 (18)	0.86983 (14)	0.69393 (10)	0.0122 (4)	
C14	0.77348 (18)	0.93480 (14)	0.65018 (10)	0.0117 (4)	
C15	1.05447 (18)	0.85200 (13)	0.76565 (10)	0.0115 (4)	
C16	1.15402 (18)	0.84218 (14)	0.73232 (10)	0.0134 (4)	
H16	1.141132	0.837617	0.681590	0.016*	
C17	1.27289 (19)	0.83913 (14)	0.77424 (11)	0.0167 (4)	
H17	1.341938	0.834798	0.751990	0.020*	
C18	1.2910 (2)	0.84240 (15)	0.84841 (12)	0.0189 (4)	
H18	1.372259	0.839435	0.876910	0.023*	
C19	1.1906 (2)	0.84998 (16)	0.88080 (11)	0.0201 (4)	
H19	1.203312	0.851183	0.931614	0.024*	
C20	1.07113 (19)	0.85588 (14)	0.83977 (10)	0.0157 (4)	
H20	1.002367	0.862426	0.862051	0.019*	
C21	0.82669 (19)	0.72630 (14)	0.76782 (11)	0.0152 (4)	
H21A	0.905166	0.712708	0.801017	0.023*	
H21B	0.816074	0.679988	0.727439	0.023*	
H21C	0.758496	0.718190	0.792885	0.023*	
C22	0.60025 (18)	0.84102 (15)	0.69320 (11)	0.0172 (4)	
H22A	0.595252	0.811728	0.739527	0.026*	
H22B	0.572689	0.792542	0.654988	0.026*	
H22C	0.547234	0.899748	0.684607	0.026*	
O1W	0.3775 (2)	0.67988 (14)	0.46049 (11)	0.0262 (4)	0.85
H1WA	0.308 (4)	0.703 (3)	0.439 (2)	0.057 (13)*	0.85
H1WB	0.425 (4)	0.734 (3)	0.465 (2)	0.050 (11)*	0.85

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.01297 (7)	0.01485 (7)	0.01124 (7)	−0.00084 (5)	0.00051 (5)	−0.00144 (5)
Cd2	0.01082 (7)	0.01089 (7)	0.01011 (7)	−0.00029 (5)	0.00088 (5)	0.00165 (5)
Cl1	0.0106 (2)	0.0169 (2)	0.0168 (2)	0.00061 (17)	0.00230 (17)	0.00041 (17)
Cl2	0.0138 (2)	0.0106 (2)	0.0098 (2)	0.00143 (17)	−0.00070 (16)	0.00113 (15)
Cl3	0.0095 (2)	0.0116 (2)	0.0125 (2)	−0.00016 (16)	0.00191 (16)	0.00141 (16)
Cl4	0.0149 (2)	0.0135 (2)	0.0168 (2)	−0.00281 (17)	0.00625 (18)	−0.00634 (17)
O1	0.0113 (7)	0.0193 (7)	0.0127 (7)	0.0002 (6)	0.0012 (5)	−0.0027 (5)
O2	0.0111 (7)	0.0178 (7)	0.0128 (7)	−0.0034 (6)	0.0009 (5)	0.0046 (5)
N1	0.0111 (8)	0.0164 (8)	0.0125 (8)	0.0003 (6)	0.0027 (6)	−0.0032 (6)
N2	0.0135 (8)	0.0186 (8)	0.0123 (8)	0.0022 (7)	0.0042 (6)	−0.0023 (6)
N3	0.0128 (8)	0.0116 (8)	0.0144 (8)	−0.0003 (7)	−0.0018 (7)	−0.0006 (6)
N4	0.0101 (8)	0.0154 (8)	0.0134 (8)	−0.0014 (6)	0.0033 (6)	0.0040 (6)
N5	0.0111 (8)	0.0134 (8)	0.0146 (8)	−0.0011 (6)	0.0050 (6)	0.0028 (6)
N6	0.0112 (8)	0.0135 (8)	0.0139 (8)	0.0014 (7)	0.0006 (7)	0.0017 (6)
C1	0.0132 (9)	0.0090 (9)	0.0127 (9)	−0.0021 (7)	0.0008 (7)	−0.0015 (7)

C2	0.0136 (9)	0.0101 (9)	0.0163 (9)	0.0001 (7)	0.0017 (8)	−0.0005 (7)
C3	0.0123 (9)	0.0086 (8)	0.0164 (10)	−0.0009 (7)	0.0013 (7)	−0.0004 (7)
C4	0.0125 (9)	0.0152 (9)	0.0089 (9)	−0.0003 (7)	0.0013 (7)	0.0005 (7)
C5	0.0181 (10)	0.0137 (9)	0.0103 (9)	0.0012 (8)	0.0013 (8)	0.0005 (7)
C6	0.0196 (10)	0.0187 (10)	0.0105 (9)	−0.0061 (8)	0.0005 (8)	−0.0006 (7)
C7	0.0147 (10)	0.0285 (11)	0.0149 (10)	−0.0024 (9)	0.0005 (8)	0.0005 (8)
C8	0.0163 (10)	0.0222 (11)	0.0217 (11)	0.0061 (9)	0.0034 (8)	0.0006 (8)
C9	0.0189 (10)	0.0138 (9)	0.0144 (9)	0.0023 (8)	0.0020 (8)	−0.0010 (7)
C10	0.0186 (11)	0.0301 (12)	0.0120 (10)	0.0017 (9)	0.0033 (8)	−0.0013 (8)
C11	0.0141 (10)	0.0195 (10)	0.0226 (11)	0.0018 (8)	0.0046 (8)	−0.0037 (8)
C12	0.0130 (9)	0.0116 (9)	0.0094 (9)	0.0002 (7)	0.0011 (7)	0.0000 (7)
C13	0.0120 (9)	0.0121 (9)	0.0124 (9)	0.0007 (7)	0.0024 (7)	−0.0016 (7)
C14	0.0108 (9)	0.0125 (9)	0.0114 (9)	0.0007 (7)	0.0011 (7)	−0.0016 (7)
C15	0.0113 (9)	0.0086 (8)	0.0138 (9)	0.0003 (7)	0.0007 (7)	0.0025 (7)
C16	0.0156 (10)	0.0103 (9)	0.0146 (9)	0.0006 (7)	0.0037 (8)	0.0011 (7)
C17	0.0136 (10)	0.0103 (9)	0.0267 (11)	0.0012 (8)	0.0051 (8)	0.0006 (8)
C18	0.0148 (10)	0.0133 (10)	0.0248 (11)	0.0008 (8)	−0.0044 (8)	0.0030 (8)
C19	0.0241 (11)	0.0197 (10)	0.0140 (10)	0.0000 (9)	−0.0014 (8)	0.0032 (8)
C20	0.0180 (10)	0.0159 (10)	0.0133 (9)	0.0014 (8)	0.0033 (8)	0.0022 (7)
C21	0.0173 (10)	0.0120 (9)	0.0172 (10)	−0.0002 (8)	0.0056 (8)	0.0046 (7)
C22	0.0122 (10)	0.0195 (10)	0.0205 (10)	−0.0018 (8)	0.0049 (8)	0.0020 (8)
O1W	0.0261 (11)	0.0187 (10)	0.0349 (11)	−0.0030 (9)	0.0087 (9)	0.0008 (8)

Geometric parameters (Å, °)

Cd1—O1	2.3498 (14)	C5—C6	1.386 (3)
Cd1—N3	2.2863 (17)	C5—H5	0.9500
Cd1—Cl1	2.4486 (5)	C6—C7	1.382 (3)
Cd1—Cl2	2.5396 (5)	C6—H6	0.9500
Cd1—Cl3	2.6908 (5)	C7—C8	1.387 (3)
Cd2—O2	2.3107 (13)	C7—H7	0.9500
Cd2—N6	2.4346 (17)	C8—C9	1.391 (3)
Cd2—Cl2	2.5929 (5)	C8—H8	0.9500
Cd2—Cl3 ⁱ	2.6734 (5)	C9—H9	0.9500
Cd2—Cl3	2.7631 (5)	C10—H10A	0.9800
Cd2—Cl4	2.5157 (5)	C10—H10B	0.9800
Cd1—Cd2	3.7831 (5)	C10—H10C	0.9800
Cd2—Cd2 ⁱ	4.0745 (4)	C11—H11A	0.9800
Cd1—Cd2 ⁱ	4.6003 (4)	C11—H11B	0.9800
O1—C1	1.261 (2)	C11—H11C	0.9800
O2—C12	1.259 (2)	C12—C14	1.413 (3)
N1—C1	1.378 (2)	C13—C14	1.367 (3)
N1—N2	1.403 (2)	C13—C22	1.482 (3)
N1—C4	1.428 (2)	C15—C20	1.388 (3)
N2—C2	1.361 (3)	C15—C16	1.388 (3)
N2—C10	1.465 (3)	C16—C17	1.391 (3)
N3—C3	1.425 (3)	C16—H16	0.9500
N3—H3NA	0.88 (3)	C17—C18	1.387 (3)

N3—H3NB	0.88 (3)	C17—H17	0.9500
N4—C12	1.367 (2)	C18—C19	1.383 (3)
N4—N5	1.400 (2)	C18—H18	0.9500
N4—C15	1.425 (2)	C19—C20	1.392 (3)
N5—C13	1.370 (2)	C19—H19	0.9500
N5—C21	1.473 (2)	C20—H20	0.9500
N6—C14	1.430 (2)	C21—H21A	0.9800
N6—H6NA	0.87 (3)	C21—H21B	0.9800
N6—H6NB	0.86 (3)	C21—H21C	0.9800
C1—C3	1.412 (3)	C22—H22A	0.9800
C2—C3	1.367 (3)	C22—H22B	0.9800
C2—C11	1.485 (3)	C22—H22C	0.9800
C4—C9	1.386 (3)	O1W—H1WA	0.85 (4)
C4—C5	1.390 (3)	O1W—H1WB	0.89 (4)
N3—Cd1—O1	77.62 (5)	C6—C5—H5	120.5
N3—Cd1—Cl1	142.63 (5)	C4—C5—H5	120.5
O1—Cd1—Cl1	94.93 (4)	C7—C6—C5	120.39 (19)
N3—Cd1—Cl2	111.23 (5)	C7—C6—H6	119.8
O1—Cd1—Cl2	104.82 (4)	C5—C6—H6	119.8
Cl1—Cd1—Cl2	106.087 (16)	C6—C7—C8	120.1 (2)
N3—Cd1—Cl3	84.54 (4)	C6—C7—H7	119.9
O1—Cd1—Cl3	161.00 (4)	C8—C7—H7	119.9
Cl1—Cd1—Cl3	95.335 (15)	C7—C8—C9	120.4 (2)
Cl2—Cd1—Cl3	87.599 (14)	C7—C8—H8	119.8
O2—Cd2—N6	76.15 (5)	C9—C8—H8	119.8
O2—Cd2—Cl4	94.60 (4)	C4—C9—C8	118.74 (19)
N6—Cd2—Cl4	98.87 (5)	C4—C9—H9	120.6
O2—Cd2—Cl2	159.61 (4)	C8—C9—H9	120.6
N6—Cd2—Cl2	91.46 (4)	N2—C10—H10A	109.5
Cl4—Cd2—Cl2	103.381 (16)	N2—C10—H10B	109.5
O2—Cd2—Cl3 ⁱ	87.33 (4)	H10A—C10—H10B	109.5
N6—Cd2—Cl3 ⁱ	161.49 (4)	N2—C10—H10C	109.5
Cl4—Cd2—Cl3	170.343 (16)	H10A—C10—H10C	109.5
Cl4—Cd2—Cl3 ⁱ	90.613 (15)	H10B—C10—H10C	109.5
O2—Cd2—Cl3	77.99 (4)	C2—C11—H11A	109.5
N6—Cd2—Cl3	85.47 (4)	C2—C11—H11B	109.5
Cl2—Cd2—Cl3	85.035 (14)	H11A—C11—H11B	109.5
Cl3 ⁱ —Cd2—Cl3	82.931 (14)	C2—C11—H11C	109.5
Cd1—Cl2—Cd2	94.958 (15)	H11A—C11—H11C	109.5
Cd2 ⁱ —Cl3—Cd1	118.092 (16)	H11B—C11—H11C	109.5
Cd2 ⁱ —Cl3—Cd2	97.069 (14)	O2—C12—N4	124.29 (18)
Cd1—Cl3—Cd2	87.825 (14)	O2—C12—C14	128.91 (18)
C1—O1—Cd1	106.61 (12)	N4—C12—C14	106.78 (16)
C12—O2—Cd2	109.64 (12)	C14—C13—N5	109.23 (17)
C1—N1—N2	108.41 (15)	C14—C13—C22	128.88 (18)
C1—N1—C4	124.28 (16)	N5—C13—C22	121.85 (17)
N2—N1—C4	121.00 (15)	C13—C14—C12	107.75 (17)

C2—N2—N1	107.55 (15)	C13—C14—N6	132.43 (18)
C2—N2—C10	124.72 (17)	C12—C14—N6	119.82 (17)
N1—N2—C10	118.20 (16)	C20—C15—C16	121.33 (18)
C3—N3—Cd1	107.89 (12)	C20—C15—N4	121.20 (17)
C3—N3—H3NA	116.3 (19)	C16—C15—N4	117.43 (17)
Cd1—N3—H3NA	104.2 (19)	C15—C16—C17	119.07 (18)
C3—N3—H3NB	113.2 (16)	C15—C16—H16	120.5
Cd1—N3—H3NB	108.9 (16)	C17—C16—H16	120.5
H3NA—N3—H3NB	106 (2)	C18—C17—C16	120.26 (19)
C12—N4—N5	108.87 (15)	C18—C17—H17	119.9
C12—N4—C15	124.31 (16)	C16—C17—H17	119.9
N5—N4—C15	122.94 (15)	C19—C18—C17	119.89 (19)
C13—N5—N4	106.86 (15)	C19—C18—H18	120.1
C13—N5—C21	123.36 (16)	C17—C18—H18	120.1
N4—N5—C21	117.39 (15)	C18—C19—C20	120.74 (19)
C14—N6—Cd2	104.86 (12)	C18—C19—H19	119.6
C14—N6—H6NA	111.8 (18)	C20—C19—H19	119.6
Cd2—N6—H6NA	109.1 (18)	C15—C20—C19	118.67 (19)
C14—N6—H6NB	113.7 (18)	C15—C20—H20	120.7
Cd2—N6—H6NB	110.6 (18)	C19—C20—H20	120.7
H6NA—N6—H6NB	107 (2)	N5—C21—H21A	109.5
O1—C1—N1	125.05 (18)	N5—C21—H21B	109.5
O1—C1—C3	128.77 (18)	H21A—C21—H21B	109.5
N1—C1—C3	106.15 (16)	N5—C21—H21C	109.5
N2—C2—C3	108.73 (17)	H21A—C21—H21C	109.5
N2—C2—C11	121.87 (17)	H21B—C21—H21C	109.5
C3—C2—C11	129.40 (19)	C13—C22—H22A	109.5
C2—C3—C1	108.52 (17)	C13—C22—H22B	109.5
C2—C3—N3	132.32 (18)	H22A—C22—H22B	109.5
C1—C3—N3	119.10 (17)	C13—C22—H22C	109.5
C9—C4—C5	121.40 (19)	H22A—C22—H22C	109.5
C9—C4—N1	118.13 (17)	H22B—C22—H22C	109.5
C5—C4—N1	120.42 (17)	H1WA—O1W—H1WB	101 (3)
C6—C5—C4	118.96 (19)		
C1—N1—N2—C2	−8.1 (2)	C6—C7—C8—C9	0.0 (3)
C4—N1—N2—C2	−161.26 (17)	C5—C4—C9—C8	1.4 (3)
C1—N1—N2—C10	−155.99 (17)	N1—C4—C9—C8	−175.99 (18)
C4—N1—N2—C10	50.8 (2)	C7—C8—C9—C4	−1.1 (3)
C12—N4—N5—C13	7.4 (2)	Cd2—O2—C12—N4	173.58 (15)
C15—N4—N5—C13	166.03 (17)	Cd2—O2—C12—C14	−8.5 (2)
C12—N4—N5—C21	150.92 (17)	N5—N4—C12—O2	172.46 (17)
C15—N4—N5—C21	−50.4 (2)	C15—N4—C12—O2	14.2 (3)
Cd1—O1—C1—N1	178.55 (15)	N5—N4—C12—C14	−5.9 (2)
Cd1—O1—C1—C3	0.8 (2)	C15—N4—C12—C14	−164.13 (17)
N2—N1—C1—O1	−172.91 (17)	N4—N5—C13—C14	−6.0 (2)
C4—N1—C1—O1	−20.8 (3)	C21—N5—C13—C14	−146.82 (17)
N2—N1—C1—C3	5.3 (2)	N4—N5—C13—C22	172.07 (17)

C4—N1—C1—C3	157.38 (17)	C21—N5—C13—C22	31.3 (3)
N1—N2—C2—C3	7.6 (2)	N5—C13—C14—C12	2.5 (2)
C10—N2—C2—C3	152.94 (19)	C22—C13—C14—C12	−175.43 (19)
N1—N2—C2—C11	−172.06 (17)	N5—C13—C14—N6	−177.43 (19)
C10—N2—C2—C11	−26.8 (3)	C22—C13—C14—N6	4.7 (4)
N2—C2—C3—C1	−4.4 (2)	O2—C12—C14—C13	−176.07 (19)
C11—C2—C3—C1	175.25 (19)	N4—C12—C14—C13	2.1 (2)
N2—C2—C3—N3	172.5 (2)	O2—C12—C14—N6	3.8 (3)
C11—C2—C3—N3	−7.8 (4)	N4—C12—C14—N6	−177.94 (16)
O1—C1—C3—C2	177.47 (19)	Cd2—N6—C14—C13	−177.09 (18)
N1—C1—C3—C2	−0.6 (2)	Cd2—N6—C14—C12	3.0 (2)
O1—C1—C3—N3	0.1 (3)	C12—N4—C15—C20	123.4 (2)
N1—C1—C3—N3	−178.00 (16)	N5—N4—C15—C20	−32.0 (3)
Cd1—N3—C3—C2	−177.59 (18)	C12—N4—C15—C16	−54.3 (3)
Cd1—N3—C3—C1	−0.9 (2)	N5—N4—C15—C16	150.31 (17)
C1—N1—C4—C9	68.2 (2)	C20—C15—C16—C17	−1.9 (3)
N2—N1—C4—C9	−142.99 (18)	N4—C15—C16—C17	175.82 (17)
C1—N1—C4—C5	−109.2 (2)	C15—C16—C17—C18	2.2 (3)
N2—N1—C4—C5	39.6 (3)	C16—C17—C18—C19	−0.8 (3)
C9—C4—C5—C6	−0.6 (3)	C17—C18—C19—C20	−1.0 (3)
N1—C4—C5—C6	176.71 (17)	C16—C15—C20—C19	0.2 (3)
C4—C5—C6—C7	−0.5 (3)	N4—C15—C20—C19	−177.46 (18)
C5—C6—C7—C8	0.8 (3)	C18—C19—C20—C15	1.3 (3)

Symmetry code: (i) $-x+2, -y+2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}$, $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3N <i>A</i> ⋯O2 ⁱ	0.88 (3)	1.89 (3)	2.747 (2)	166 (3)
N6—H6N <i>A</i> ⋯C11	0.87 (3)	2.54 (3)	3.4073 (19)	172 (2)
N6—H6N <i>B</i> ⋯C11 ⁱⁱ	0.86 (3)	2.73 (3)	3.3718 (18)	133 (2)
O1 <i>W</i> —H1 <i>WA</i> ⋯C14 ⁱⁱ	0.85 (4)	2.45 (4)	3.292 (2)	176 (4)
O1 <i>W</i> —H1 <i>WB</i> ⋯C11	0.89 (4)	2.34 (4)	3.220 (2)	168 (3)
C11—H11 <i>A</i> ⋯C12 ⁱⁱⁱ	0.98	2.82	3.732 (2)	156
C16—H16⋯C12 ⁱ	0.95	2.82	3.589 (2)	138
C21—H21 <i>C</i> ⋯C14 ^{iv}	0.98	2.69	3.647 (2)	165
C22—H22 <i>C</i> ⋯O1 ⁱⁱ	0.98	2.51	3.417 (3)	154

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x+1, -y+2, -z+1$; (iii) $-x+3/2, y-1/2, -z+1/2$; (iv) $-x+3/2, y-1/2, -z+3/2$.