


Stephen G. Addinall,<sup>a</sup> Kenneth A. Johnson,<sup>b</sup> Timothy Dafforn,<sup>c</sup> Corinne Smith,<sup>d</sup> Alison Rodger,<sup>e,f</sup> Raul Paco Gomez,<sup>d,f</sup> Katherine Sloan,<sup>a</sup> Anne Blewett,<sup>d</sup> David J. Scott<sup>g</sup> and David I. Roper<sup>d\*</sup>

<sup>a</sup>Faculty of Life Sciences, University of Manchester, Michael Smith Building, Oxford Road, Manchester M13 9PT, England, <sup>b</sup>Department of Biochemistry and Biophysics, The Arrhenius Laboratories for Natural Sciences, Stockholm University, SE-106 91 Stockholm, Sweden, <sup>c</sup>Department of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, England, <sup>d</sup>Department of Biological Sciences, University of Warwick, Coventry CV4 7AL, England, <sup>e</sup>Department of Chemistry, University of Warwick, Coventry CV4 7AL, England, <sup>f</sup>MOAC Doctoral Training Centre, University of Warwick, Coventry CV4 7AL, England, and <sup>g</sup>National Centre for Macromolecular Hydrodynamics, School of Biosciences, University of Nottingham, Sutton Bonington, Leicestershire LE12 5RD, England

Correspondence e-mail:  
david.roper@warwick.ac.uk

Received 21 December 2004  
Accepted 3 February 2005  
Online 24 February 2005



© 2005 International Union of Crystallography  
All rights reserved

## Expression, purification and crystallization of the cell-division protein YgfE from *Escherichia coli*

An open reading frame designated b2910 (*ygfE*) in the *Escherichia coli* K12-MG1655 genome sequence, identified as a possible homologue to the cell-division protein ZapA, was cloned into the high-expression plasmid pETDuet-1 and overexpressed in *E. coli* BL21 (DE3)-AI. The protein was purified in three steps to 99% purity. Crystals were obtained by the hanging-drop vapour-diffusion method at 291 K from a wide range of screened conditions, but principally from solutions containing 0.1 M HEPES pH 7.0, 18% PEG 6000, 5 mM CaCl<sub>2</sub>. Diffraction data to 1.8 Å were collected at the European Synchrotron Radiation Facility (ESRF). The crystals belong to space group *P*6<sub>1</sub>22 or *P*6<sub>5</sub>22, with unit-cell parameters  $a = 53.8$ ,  $b = 53.8$ ,  $c = 329.7$  Å,  $\alpha = \beta = 90$ ,  $\gamma = 120^\circ$ .

### 1. Introduction

The ancestral tubulin protein FtsZ has a crucial role in the process of bacterial cell division. FtsZ polymerizes into a ring (the 'Z-ring') around the middle of pre-divisional cells on the inner surface of the cytoplasmic membrane. The Z-ring then constricts, driving cell division (Addinall & Holland, 2002; Lutkenhaus & Addinall, 1997; Margolin, 2000). FtsZ is extensively conserved throughout bacterial species and plants (where it is required for plastid division). FtsZ has also been implicated in the division of some archaeal species and the mitochondria of some eukaryotes (Vaughan *et al.*, 2004). As a cytoskeletal element with prokaryotic origins the Z-ring represents a fascinating topic of study and as a major player in a fundamental process for bacterial life FtsZ represents an exciting potential target for novel antibacterial compounds.

The mechanisms by which Z-rings form and constrict are poorly characterized and the detailed *in vivo* structure of the Z-ring is unknown. *In vitro* studies reveal that FtsZ is a GTPase which can polymerize into linear polymers in a GTP-dependent manner (Romberg & Levin, 2003), and that these polymers form lateral associations to a greater or lesser degree depending on polymerization conditions. Measurements indicate that cells contain sufficient FtsZ for the Z-ring to be composed of multiple protofilaments (Lu *et al.*, 1998; Feucht *et al.*, 2001). This is likely to be the case since cells containing GFP-FtsZ fusion protein incorporate approximately 30% of the GFP fluorescence into the Z-ring (Anderson *et al.*, 2004). Therefore, lateral interactions between FtsZ polymers are likely to be important in Z-ring function.

Two FtsZ accessory proteins have been identified that promote lateral associations between FtsZ polymers. ZipA (from *Escherichia coli*) is a membrane-spanning protein with a cytoplasmic domain that interacts with the FtsZ C-terminus (Hale & de Boer, 1997). This domain of ZipA induces bundling of FtsZ polymers *in vitro* (Hale *et al.*, 2000; RayChaudhuri, 1999) and its structure has been determined in complex with a small FtsZ peptide (Moy *et al.*, 2000). ZapA from both *Bacillus subtilis* and *Pseudomonas aeruginosa* has been demonstrated to bind and to induce bundling between FtsZ polymers (Gueiros-Filho & Losick, 2002; Low *et al.*, 2004), and the crystal structure of tetrameric *P. aeruginosa* ZapA has been solved (Low *et al.*, 2004). We recently demonstrated that the *E. coli* YgfE protein binds to and bundles FtsZ polymers, inhibits the FtsZ GTPase and adopts a dimer-tetramer equilibrium in solution (Small *et al.*,



We are therefore investigating MIR and MAD strategies to obtain a solution to the structure of YgfE as part of a coordinated study investigating the interaction of YgfE, FtsZ and other cell-division proteins.

We are grateful for access and user support at the synchrotron facilities of the ESRF, Grenoble and SRS, Daresbury. We would like to thank Klaus Fütterer (University of Birmingham) for assistance with data collection. AB is funded by a BBSRC studentship; RPG is supported *via* the EPSRC-supported Molecular Organisation and Assembly in Cells (MOAC) doctoral training centre at Warwick, KAJ is supported by the Karl Trygger Foundation, SA and TD are supported by the Wellcome Trust and MRC Career Development fellowships, respectively.

### References

- Addinall, S. G. & Holland, B. (2002). *J. Mol. Biol.* **318**, 219–236.
- Anderson, D. E., Gueiros-Filho, F. J. & Erickson, H. P. (2004). *J. Bacteriol.* **186**, 5775–5781.
- Brzozowski, A. M. & Walton, J. (2001). *J. Appl. Cryst.* **34**, 97–101.
- Feucht, A., Lucet, I., Yudkin, M. D. & Errington, J. (2001). *Mol. Microbiol.* **40**, 115–125.
- Gueiros-Filho, F. J. & Losick, R. (2002). *Genes Dev.* **16**, 2544–2556.
- Hale, C. A. & de Boer, P. A. (1997). *Cell*, **88**, 175–185.
- Hale, C. A., Rhee, A. C. & de Boer, P. A. (2000). *J. Bacteriol.* **182**, 5153–5166.
- Jancarik, J. & Kim, S.-H. (1991). *J. Appl. Cryst.* **24**, 409–411.
- Low, H. H., Moncrieffe, M. C. & Löwe, J. (2004). *J. Mol. Biol.* **341**, 839–852.
- Lu, C., Stricker, J. & Erickson, H. P. (1998). *Cell Motil. Cytoskeleton*, **40**, 71–86.
- Lutkenhaus, J. & Addinall, S. G. (1997). *Annu. Rev. Biochem.* **66**, 93–116.
- Margolin, W. (2000). *FEMS Microbiol. Rev.* **24**, 531–548.
- Moy, F. J., Glasfeld, E., Mosyak, L. & Powers, R. (2000). *Biochemistry*, **39**, 9146–9156.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- RayChaudhuri, D. (1999). *EMBO J.* **18**, 2372–2383.
- Romberg, L. & Levin, P. A. (2003). *Annu. Rev. Microbiol.* **57**, 125–154.
- Vaughan, S., Wickstead, B., Gull, K. & Addinall, S. G. (2004). *J. Mol. Evol.* **58**, 19–29.