

Phillip J. Monaghan, David Leys  
and Nigel S. Scrutton\*

Department of Biochemistry, University of  
Leicester, University Road, Leicester LE1 7RH,  
England

Correspondence e-mail: nss4@le.ac.uk

Received 8 April 2005  
Accepted 29 June 2005  
Online 8 July 2005

## Crystallization and preliminary X-ray diffraction analysis of a flavoenzyme amine dehydrogenase/oxidase from *Pyrococcus furiosus* DSM 3638

A flavoprotein amine dehydrogenase/oxidase with subunit molecular weights of 54.8 kDa ( $\alpha$ -subunit) and 42.4 kDa ( $\beta$ -subunit) and specificity for L-proline was cloned from the genomic DNA of the hyperthermophilic marine archaeon *Pyrococcus furiosus* DSM 3638. The enzyme was overexpressed in *Escherichia coli* and purified to homogeneity. The enzyme was crystallized using the sitting-drop vapour-diffusion technique. Diffraction data from two different crystal forms were collected to 3.3 and 3.6 Å, respectively, using synchrotron radiation. Both crystals belonged to space group *P*1, with unit-cell parameters  $a = 91.3$ ,  $b = 136.3$ ,  $c = 203.8$  Å,  $\alpha = 94.5$ ,  $\beta = 99.4$ ,  $\gamma = 102.7^\circ$  and  $a = 93.7$ ,  $b = 116.3$ ,  $c = 126.9$  Å,  $\alpha = 97.3$ ,  $\beta = 99.9$ ,  $\gamma = 104.6^\circ$ .

### 1. Introduction

The mechanisms of amine oxidation catalysed by the quinoprotein amine dehydrogenase/oxidases are well established. Oxidation occurs through the formation of enzyme–substrate covalent adducts with topaquinone (TPQ), tryptophan tryptophylquinone (TTQ), cysteine tryptophylquinone (CTQ) and lysine tyrosyl quinone (LTQ) redox centres (Mure *et al.*, 2002; Datta *et al.*, 2001; Satoh *et al.*, 2002; Davidson, 2000; Wang *et al.*, 1996) and hydrogen transfer by quantum-mechanical tunnelling has been demonstrated for some quinoprotein enzymes (Masgrau *et al.*, 2004). The mechanism of amine oxidation by flavoproteins is less well understood (Scrutton, 2004). These enzymes catalyse the two-electron oxidation of amine substrates to the corresponding iminium ion. Mechanisms involving (i) proton abstraction by an active-site base to generate a carbanion species (Rohlf & Hille, 1994), (ii) an aminium radical cation species (Silverman, 1995), (iii) H-atom abstraction by an active-site radical species (Edmondson, 1995) and (iv) nucleophilic attack by the substrate nitrogen on the flavin C4a atom followed by proton abstraction by an active-site base or the flavin N5 atom (Kim *et al.*, 1993) have been considered over the years, but controversy remains in the field. Further structural elucidation of flavoprotein amine dehydrogenase/oxidases coupled with detailed spectroscopic and kinetic studies are required to unravel further the mechanistic details of amine oxidation by this class of enzyme.

We have identified a putative flavoprotein amine dehydrogenase/oxidase from *Pyrococcus furiosus* DSM 3638. The  $\beta$ -subunit of this putative amine oxidase is related to the active-site region of dimethylglycine oxidase (DMGO) of *Arthrobacter globiformis*, for which a crystal structure has been elucidated and a mechanism of substrate oxidation proposed (Leys *et al.*, 2003). The *P. furiosus* enzyme oxidizes sarcosine, L-pipecolic acid and L-proline, but is not reactive with dimethylglycine, glycine or betaine. Kinetic studies suggest that L-proline is the natural substrate, but the enzyme is not related in sequence to the structurally characterized and bifunctional proline dehydrogenase of *Escherichia coli* (Lee *et al.*, 2003). The flavoprotein amine dehydrogenase/oxidase from *P. furiosus* appears to represent a new class of proline-oxidizing flavoprotein. To further our understanding of the mechanism and structure of flavoprotein amine dehydrogenase/oxidases, we have initiated a structural study of this novel flavoprotein amine dehydrogenase/oxidase from *P. furiosus*. We have expressed the recombinant enzyme in *E. coli* and



© 2005 International Union of Crystallography  
All rights reserved

**Table 1**

 Data-collection and processing statistics for the form 1 and form 2 crystals of recombinant *P. furiosus* flavoprotein amine oxidase/dehydrogenase.


Crystal form	1	2
Synchrotron radiation	Beamline ID 14-3, ESRF	Beamline ID 14-3, ESRF
Space group	<i>P1</i>	<i>P1</i>
Wavelength (Å)	0.931	0.931
Unit-cell parameters		
<i>a</i> (Å)	93.7	91.3
<i>b</i> (Å)	116.3	136.3
<i>c</i> (Å)	126.9	203.8
$\alpha$ (°)	97.3	94.5
$\beta$ (°)	99.9	99.4
$\gamma$ (°)	104.6	102.7
Matthews coefficient (Å <sup>3</sup> Da <sup>-1</sup> )	3.34	3.12
Resolution (Å)	30–3.6 (3.8–3.6)	30–3.3 (3.4–3.3)
<i>R</i> <sub>merge</sub> (%)	0.120 (0.432)	0.084 (0.357)
Completeness (%)	96.4 (95.0)	96.0 (93.0)
Average <i>I</i> / $\sigma$ ( <i>I</i> )	6.2 (1.8)	8.2 (2.0)
Average redundancy	2.5 (2.3)	1.9 (1.8)
Unique reflections	55367	132489

purified the protein to homogeneity. Diffraction data are reported to 3.3 and 3.6 Å for two crystal forms both in space group *P1*.

## 2. Materials and methods

### 2.1. Cloning, overexpression and purification

Two open reading frames [gi|18977617 and gi|18977618; protein-extraction description and analysis tool (PEDANT) database] were identified in the genome of *P. furiosus* DSM 3638 that encode a putative flavoenzyme amine dehydrogenase/oxidase and were amplified by the polymerase chain reaction. Both genes encoding the  $\alpha$ - and  $\beta$ -subunits (492 and 382 amino-acid residues, respectively) were cloned into pET11d (Novagen) and the resultant construct transformed into *E. coli* strain Rosetta(DE3)pLysS for recombinant expression. Details of plasmid construction and propagation will be reported elsewhere. Selected transformed cells were grown in 12 l of 2×YT medium containing chloramphenicol (34 µg ml<sup>-1</sup>) and ampicillin (50 µg ml<sup>-1</sup>) at 310 K to an OD<sub>600</sub> ≈ 0.8 and were induced by adding isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) to a final concentration of 1 mM. Incubation took place for a further 8 h, after which cells were harvested and resuspended in 50 mM ice-cold potassium phosphate buffer pH 8.0 containing a Complete protease-inhibitor cocktail tablet (Roche). The cell suspension was incubated with lysozyme (10 µg ml<sup>-1</sup>) and sonicated to effect cell breakage.


**Figure 1**

Crystals (form 1) of the recombinant flavoprotein amine oxidase/dehydrogenase from *P. furiosus*.

**Table 2**

Peaks observed in the self-rotation analysis of the form 1 crystals.

The polar angles  $\theta$ ,  $\varphi$  and  $\chi$  are defined as follows:  $\theta$  is the angle between the rotation axis and  $z$ ,  $\varphi$  is the angle in the  $xy$  plane between  $x$  and the projection of the rotation axis and  $\chi$  is the rotation angle around the rotation axis.


Peak	$\theta$ (°)	$\varphi$ (°)	$\chi$ (°)	<i>R</i> / $\sigma$
1	86.6	-30.9	180	10.6
2	127.9	57.1	180	10.5
3	38.3	64.3	180	9.7

Deoxyribonuclease I (~10 µg) was added and the suspension was incubated for a further 30 min at 277 K. Cellular debris was removed by centrifugation and the supernatant was incubated at 353 K for 1 h followed by centrifugation to remove denatured host-cell proteins. The supernatant was dialysed exhaustively against 50 mM Tris-HCl pH 8.0 at 277 K, loaded onto a Q-Sepharose column (75 ml bed volume) and eluted using a 0.2–0.4 M gradient of NaCl. Fractions containing recombinant protein were pooled, concentrated and applied onto a Superdex 75 column (330 ml bed volume); protein was eluted at a flow rate of 0.75 ml min<sup>-1</sup>. Purified enzyme was exchanged into 10 mM Tris-HCl buffer pH 7.5 containing 100 mM KCl, concentrated to 12.2 mg ml<sup>-1</sup>, sterilized (0.22 µm Millex-GP Acrodisc filter, Millipore) and stored on ice at 277 K. Protein concentration was determined by the method of Bradford (1976). Purity was >95% as judged by SDS-PAGE.


### 2.2. Crystallization

Crystallization trials were performed using the sitting-drop vapour-diffusion technique. Drops were prepared by mixing 2 µl of 6.1 mg ml<sup>-1</sup> protein solution with 2 µl reservoir solution. After 3–4 d incubation at 292 K, yellow crystals were observed in two conditions containing 0.2 M potassium thiocyanate, 8% PEG 20K, 8% PEG 550 MME pH 7.5, 100 mM Tris (crystal form 1; Fig. 1) and 0.2 M sodium formate, 8% PEG 20K and 8% PEG 550 MME pH 8.5, 100 mM Tris (crystal form 2; Fig. 2) in the reservoir solution. The average dimensions of these crystals were 40 × 250 × 250 µm (form 1) and 50 × 50 × 200 µm (form 2).

Prior to data collection at 100 K, selected crystals were soaked in mother liquor with 10% PEG 200 as cryoprotectant and flash-cooled in liquid nitrogen.


**Figure 2**

Crystals (form 2) of the recombinant flavoprotein amine oxidase/dehydrogenase from *P. furiosus*.



**Figure 3**  
Self-rotation function calculated using crystal form 1 data between 30 and 3.6 Å with a 50.0 Å radius of integration. Twofold symmetry is evident from the very strong peaks in the  $\chi = 180^\circ$  plot.

### 2.3. X-ray diffraction analysis

Diffraction data were collected from cryocooled crystals (100 K) using a Quantum ADSC (Area Detector Systems Corporation) CCD detector on beamline ID 14-3 at the European Synchrotron Radiation Facility, Grenoble, France. Data were collected using  $1^\circ$  oscillations with the crystal-to-detector distance set to 230 mm. All data were indexed, integrated and scaled using the programs *DENZO* and *SCALEPACK* from the *HKL* software package (Otwinowski & Minor, 1997). Crystals of form 1 diffracted to a resolution of 3.6 Å and belonged to space group *P1*, with unit-cell parameters  $a = 93.7$ ,  $b = 116.3$ ,  $c = 126.9$  Å,  $\alpha = 97.3$ ,  $\beta = 99.9$ ,  $\gamma = 104.6^\circ$ . Crystals of form 2 diffracted to a resolution of 3.3 Å and belonged to space group *P1*, with unit-cell parameters  $a = 91.3$ ,  $b = 136.3$ ,  $c = 203.8$  Å,  $\alpha = 94.5$ ,  $\beta = 99.4$ ,  $\gamma = 102.7^\circ$ . Diffraction data-collection statistics are given in Table 1. Using the program *MOLREP* (Vagin, 1997), a general self-rotation function was computed for a number of  $\kappa$  angles (60, 90, 120 and  $180^\circ$ ) to test for the presence of non-crystallographic twofold, threefold, fourfold and sixfold axes. A self-rotation function of the form 1 data set indicates the presence of

three perpendicular twofold axes in the unit cell, which strongly suggests the protein is a heterooctamer, with heterodimers arranged in 222 non-crystallographic symmetry (Table 2, Fig. 3). Four heterodimers (one heterooctamer) per unit cell, totalling 388.8 kDa, corresponds to a Matthews coefficient of  $3.34 \text{ \AA}^3 \text{ Da}^{-1}$  and a solvent content of  $\sim 62.8\%$  (Matthews, 1968). Self-rotation functions of data from form 2 did not reveal any clear indication of the presence of non-crystallographic symmetry. However, two heterooctamers per unit cell, totalling 777.6 kDa, corresponds to a Matthews coefficient of  $3.12 \text{ \AA}^3 \text{ Da}^{-1}$  and a solvent content of  $\sim 60.2\%$ .

Molecular-replacement trials performed with the available structures of similar flavin oxidases as the starting model failed to identify a solution. Therefore, MIR (multiple isomorphous replacement)/MAD (multiwavelength anomalous diffraction) experiments are planned to provide sufficient experimental phasing information. A search for heavy-atom derivatives and the production of selenomethionine-substituted protein are in progress.

We are grateful for access to and user support at the ESRF, Grenoble, France. PJM thanks the BBSRC for the award of a studentship. DL is a Royal Society University Research Fellow and an EMBO Young Investigator.

### References

- Bradford, M. (1976). *Anal. Biochem.* **72**, 248–254.  
 Datta, S., Mori, Y., Takagi, K., Kawaguchi, K., Chen, Z. W., Okajima, T., Kuroda, S., Ikeda, T., Kano, K., Tanizawa, K. & Mathews, F. S. (2001). *Proc. Natl Acad. Sci. USA*, **98**, 14268–14273.  
 Davidson, V. L. (2000). *Subcell. Biochem.* **35**, 119–143.  
 Edmondson, D. E. (1995). *Xenobiotica*, **25**, 735–753.  
 Kim, J., Bogdan, M. & Mariano, P. S. (1993). *J. Am. Chem. Soc.* **115**, 10591–10595.  
 Lee, Y. H., Nadaraja, S., Gu, D., Becker, D. F. & Tanner, J. J. (2003). *Nature Struct. Biol.* **10**, 109–114.  
 Leys, D., Basran, J. & Scrutton, N. S. (2003). *EMBO J.* **22**, 4038–4048.  
 Masgrau, L., Basran, J., Hothi, P., Sutcliffe, M. J. & Scrutton, N. S. (2004). *Arch. Biochem. Biophys.* **428**, 41–51.  
 Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.  
 Mure, M., Mills, S. A. & Klinman, J. P. (2002). *Biochemistry*, **41**, 9269–9278.  
 Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.  
 Rohlfs, R. J. & Hille, R. (1994). *J. Biol. Chem.* **269**, 30869–30879.  
 Satoh, A., Kim, J. K., Miyahara, I., Devreese, B., Vandenberghe, I., Hacısalihoglu, A., Okajima, T., Kuroda, S., Adachi, O., Duine, J. A., Van Beeumen, J., Tanizawa, K. & Hirotsu, K. (2002). *J. Biol. Chem.* **277**, 2830–2834.  
 Scrutton, N. (2004). *Nat. Prod. Rep.* **21**, 722–730.  
 Silverman, R. B. (1995). *Acc. Chem. Res.* **28**, 335–342.  
 Wang, S. X., Mure, M., Medzihradszky, K. F., Burlingame, A. L., Brown, D. E., Dooley, D. M., Smith, A. J., Kagan, H. M. & Klinman, J. P. (1996). *Science*, **273**, 1078–1084.  
 Vagin, A. (1997). *J. Appl. Cryst.* **30**, 1022–1025.